

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	44 (1998)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	THREE DISTANCE THEOREMS AND COMBINATORICS ON WORDS
<b>Autor:</b>	Alessandri, Pascal / Berthé, Valérie
<b>Kapitel:</b>	4.1 Connectedness index
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-63900">https://doi.org/10.5169/seals-63900</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## REMARKS.

- Suppose that  $\alpha$  is an irrational number. By density of the sequence  $(\{n\alpha\})_{n \in \mathbb{N}}$ , this theorem still holds when considering the gaps between the successive integers  $k$  such that  $\{\alpha k\} \in I$ , where  $I$  denotes any interval of the unit circle of length  $\beta$ .
- Furthermore, the third gap, which is the largest, can have frequency 0, when  $\eta_k = \psi$ , with the above notation. This means that this gap does not appear at all, as a consequence of the uniform distribution of the sequence  $(\{n\alpha\})_{n \in \mathbb{N}}$  in the circle.
- The other two gaps do always appear (infinitely often, in fact, because of their positive frequencies) and are shown to be equal to the smallest positive integers  $l_1$  and  $l_2$  such that  $\{l_1\alpha\} < \beta$  and  $\{l_2\alpha\} > 1 - \beta$  (see [51]).
- The study of the rational case proves the equivalence between the three distance and the three gap theorems, as observed by Slater [51] in the case of an open interval and by Langevin, for any interval, in [35].

## 4.1 CONNECTEDNESS INDEX

Let  $u = (u_n)_{n \in \mathbb{N}}$  be a coding of a rotation by irrational angle  $0 < \alpha < 1$  with respect to the partition

$$\mathcal{P} = \{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[ \} .$$

We have seen in Section 2.1 that the sets  $I(w_1, \dots, w_n) = \bigcap_{j=0}^{n-1} R^{-j}(I_{w_{j+1}})$ , where  $I_k = [\beta_k, \beta_{k+1}[$ , for  $0 \leq j \leq p-1$ , are connected except for  $w_1 \dots w_n$  of the form  $a_K^n$ , where  $K$  denotes the index of the interval of  $\mathcal{P}$  (if such an interval exists) of length greater than  $\sup(\alpha, 1 - \alpha)$ .

Let us suppose that there exists an interval of  $\mathcal{P}$  of length  $L$  greater than  $1 - \alpha$  and index  $K$ , say. We deduce from the three gap theorem that the set of integers  $n$  such that  $a_K^n$  is a factor of the sequence  $u$  is bounded. More precisely, let us define  $n^{(1)}$  as the largest integer  $n$  such that  $a_K^n$  is a factor of the sequence  $u$ . We will call the integer  $n^{(1)}$  the *index of connectedness* of the sequence  $u$ . (If every interval of  $\mathcal{P}$  has length smaller than or equal to  $\sup(\alpha, 1 - \alpha)$  then the connectedness index of  $u$  is equal to 1.) The three gap theorem enables us to give an exact expression for the connectedness index. Indeed  $n^{(1)} + 1$  is the largest gap between the consecutive values of  $k$  for which  $0 < \{k\alpha\} < 1 - L$ . We thus have the following

**THEOREM 9.** *Let  $u = (u_n)_{n \in \mathbb{N}}$  be a coding of the rotation by irrational angle  $\alpha$ . Suppose that there exists an interval of  $\mathcal{P}$  of length  $L > \sup(\alpha, 1 - \alpha)$ . Let  $(\frac{p_k}{q_k})_{k \in \mathbb{N}}$  and  $(c_k)_{k \in \mathbb{N}}$  be the sequences of convergents and partial quotients associated to  $\alpha$  in its continued fraction expansion. Let  $\eta_k = (-1)^k(q_k\alpha - p_k)$ . Write*

$$1 - L = m\eta_k + \eta_{k+1} + \psi,$$

*with  $k \geq 1$ ,  $0 < \psi \leq \eta_k$  and  $1 \leq m \leq c_{k+1}$ . The connectedness index  $n^{(1)}$  of the sequence  $u$  satisfies*

$$\begin{aligned} n^{(1)} &= q_{k+1} - (m-1)q_k - 1, \text{ if } \psi \neq \eta_k, \\ n^{(1)} &= q_{k+1} - mq_k - 1, \text{ if } \psi = \eta_k \text{ and } m < c_{k+1}, \\ n^{(1)} &= q_k - 1, \text{ if } \psi = \eta_k \text{ and } m = c_{k+1}. \end{aligned}$$

## 4.2 APPLICATIONS

Precise knowledge of the connectedness index is useful, as shown by the following. Indeed Lemma 1 can be rephrased as follows.

**LEMMA 3.** *Let  $u$  be a coding of an irrational rotation on the unit circle with respect to the partition  $\{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[\}\}$ . The frequencies of factors of  $u$  of length  $n \geq n^{(1)}$ , where  $n^{(1)}$  denotes the connectedness index, are equal to the lengths of the intervals bounded by the points*

$$\{k(1 - \alpha) + \beta_i\}, \text{ for } 0 \leq k \leq n-1, \quad 0 \leq i \leq p-1.$$

The complexity of a coding on  $p$  letters of an irrational rotation ultimately has the form  $p(n) = an + b$ , where  $a \leq p$ , and depends on the algebraic relations between the angle and the lengths of the intervals of the coding. More precisely, we have the following theorem proved in [1].

**THEOREM 10.** *Let  $u = (u_n)_{n \in \mathbb{N}}$  be a coding of the irrational rotation  $R$  of irrational angle  $\alpha$  with respect to the partition*

$$\mathcal{P} = \{[\beta_0, \beta_1[, [\beta_1, \beta_2[, \dots, [\beta_{p-1}, \beta_p[\}\}.$$

*Let  $(k_n)_{n \in \mathbb{N}}$  be the sequence defined by*

$$k_0 = p = \text{card}(F),$$

$$k_n = \text{card} \{ \beta_i \in F; \forall k \in [1, \dots, n], R^{-k}(\beta_i) \notin F \}.$$