Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 44 (1998)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THREE DISTANCE THEOREMS AND COMBINATORICS ON WORDS
Autor: Alessandri, Pascal / Berthé, Valérie
Bibliographie
DOI: https://doi.org/10.5169/seals-63900

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Proof. Let us consider a coding of the rotation by angle α under the left-closed and right-open partition of the unit circle bounded by all the points of the form $\{n\alpha + \gamma_i\}$, for $0 \leq n < n_i$ and $1 \leq i \leq d$; let $\beta_0, \ldots, \beta_{p-1}$ denote these consecutive points. The letter associated with the interval $I_k = [\beta_k, \beta_{k+1}[$ has a unique right extension, except when I_k contains points of the form $\{\beta_i - \alpha\}$. Suppose there are $q \geq 2$ points of this form; the associated letter has $q+1$ right extensions. Since there are at most d points of this type, we obtain $p(2) - p(1) \leq d$. We deduce from Theorem 6 that there are at most $3d$ different frequencies for the letters of the coding, i.e., there are at most $3d$ different lengths for the intervals I_k.

REMARK. The start and finish intervals as introduced by Liang in his proof in [37] correspond exactly to the beginning of the branches in the graph of words. Indeed, Liang shows that any interval is associated either with a start point $\{\gamma_i\}$ (i.e., with one extension of a factor having more than one right extension) or with a finish point $\{(n_i - 1)\alpha + \gamma_i\}$ (i.e., with a factor having more than one left extension). Counting the finish and start points defined in [37] (there are $3d$ such points) is equivalent to counting the number of branches in the graph of words.

As in the remark of the previous section, we can consider a coding of the rotation by irrational angle $1 - \alpha$ under the partition $\{[\gamma_1, \gamma_2[, \ldots, [\gamma_d, \gamma_1[\}$. For such a coding, the $3d$ distance theorem can be rephrased as follows.

THEOREM 20. *The frequencies of the factors of given length $n \geq n^{(1)}$ of a coding of a rotation by irrational angle under a partition in d intervals take at most $3d$ values, where $n^{(1)}$ denotes the connectedness index.*

ACKNOWLEDGEMENTS. We would like to thank Arnoux, Berstel, Daudé, Dumont, Liardet and Shallit for their bibliographic help. Furthermore, we are greatly indebted to Allouche for his many useful comments and to Ferenczi and Jenkinson who read carefully a previous version of this paper.

REFERENCES

[1] ALESSANDRI, P. Codages de rotations et basses complexités. Université Aix-Marseille II. Thèse, 1996.

[2] ALLOUCHE, J.-P. Sur la complexité des suites infinies. *Bull. Belg. Math. Soc. Simon Stevin 1* (1994), 133–143.

[3] BERTHÉ, V. Fréquences des facteurs des suites sturmiennes. *Theoret. Comput. Sci. 165* (1996), 295–309.

[4] BERTHÉ, V. and L. VUILLON. Tilings and rotations. Prétirage 97–19, Institut de Mathématiques de Luminy.

[5] BERTHÉ, V., N. CHEKHOVA, and S. FERENCZI. Covering numbers: arithmetics and dynamics for rotations and interval exchanges. Prétirage 97–20, Institut de Mathématiques de Luminy.

[6] BERSTEL, J. Recent results in Sturmian words. *Developments in Language Theory II (Magdeburg 1995).* World Scientific (96), 13–24.

[7] BESSI, G. and J.-L. NICOLAS. Nombres 2-hautement composés. *J. Math. Pures Appl. 56* (1977), 307–326.

[8] BOSHERNITZAN, M. A condition for minimal interval exchange maps to be uniquely ergodic. *Duke Math. J. 52* (1985), 723–752.

[9] BROWN, T. C. Descriptions of the characteristic sequence of an irrational. *Canad. Math. Bull. 36* (1993), 15–21.

[10] CASSAIGNE, J. Special factors of sequences with linear subword complexity. *Developments in Language Theory II (Magdeburg 1995).* World Scientific (96), 25–34.

[11] —— Complexité et facteurs spéciaux. *Bull. Belg. Math. Soc. Simon Stevin 4* (1997), 67–88.

[12] CHEKHOVA, N. Covering numbers of rotations. *Theoret. Comput. Sci.*, to appear.

[13] CHEKHOVA, N., P. HUBERT AND A. MESSAOUDI. Propriétés combinatoires, ergodiques et arithmétiques de la substitution de Tribonacci. Preprint.

[14] CHEVALLIER, N. Distances dans la suite des multiples d'un point du tore à deux dimensions. *Acta Arith. 74* (1996), 47–59.

[15] —— Géométrie des suites de Kronecker. *Manuscripta Math. 94* (1997), 231–241.

[16] —— Meilleures approximations d'un élément du tore $\mathbf{T}^2$ et géométrie de la suite des multiples de cet élément. *Acta Arith. 78* (1996), 19–35.

[17] CHVÁTAL, V., D. A. KLARNER and D. E. KNUTH. Selected combinatorial research problems. *Stanford CS report 292* (1972).

[18] CHUNG, F. R. K. and R. L. GRAHAM. On the set of distances determined by the union of arithmetic progressions. *Ars Combin. 1* (1976), 57–76.

[19] COVEN, E. M. and G. A. HEDLUND. Sequences with minimal block growth. *Math. Systems Theory 7* (1973), 138–153.

[20] DEKKING, F. M. On the Thue-Morse measure. *Acta Univ. Carolin. Math. Phys. 33* (1992), 35–40.

[21] DELÉGLISE, M. Recouvrement optimal du cercle par les multiples d'un intervalle. *Acta Arith. 59* (1991), 21–35.

[22] DIDIER, G. Caractérisation des N-écritures et applications à l'étude des suites de complexité ultimement $n + c^{ste}$. *Theoret. Comput. Sci.*, to appear.

[23] —— Combinatoire des codages de rotations. *Acta Arith.*, to appear.

[24] FERENCZI, S. Les transformations de Chacon: combinatoire, structure géométrique, lien avec les systèmes de complexité $2n + 1$. *Bull. Soc. Math. France 123* (1995), 271–292.

[25] FLOREK, K. Une remarque sur la répartition des nombres $n\xi$ (mod 1). *Colloq. Math. 2* (1951), 323–324.

[26] FRAENKEL, A. S. and R. HOLZMAN. Gap problems for integer part and fractional part sequences. *J. Number Theory 50* (1995), 66–86.

[27] FRAENKEL, A. S., M. MUSHKIN, and U. TASSA. Determination of $[n\theta]$ by its sequence of differences. *Canad. Math. Bull. 21* (1978), 441–446.

[28] FRIED, E. and V. T. SÓS. A generalization of the three-distance theorem for groups. *Algebra Universalis 29* (1992), 136–149.

[29] GEELEN, A. S. and R. J. SIMPSON. A two dimensional Steinhaus theorem. *Australas. J. Combin. 8* (1993), 169–197.

[30] GOTTSCHALK, W. H. and G. A. HEDLUND. Topological dynamics. *American Math. Soc. Colloquium Publications 36* (1955).

[31] HALTON, J. H. The distribution of the sequence $\{n\xi\}$ $(n = 0, 1, 2, \ldots)$. *Proc. Cambridge Philos. Soc. 61* (1965), 665–670.

[32] HARDY, G. H. and E. M. WRIGHT. *An Introduction to the Theory of Numbers.* (Fifth ed.) Oxford University Press (1979).

[33] HARTMAN, S. Über die Abstände von Punkten $n\xi$ auf der Kreisperipherie. *Ann. Soc. Polon. Math. 25* (1952), 110–114.

[34] KNUTH, D. E. *The Art of Computer Programming.* Vol. 3, Ex. 6.4.10. Addison Wesley, New York, 1973.

[35] LANGEVIN, M. Stimulateur cardiaque et suites de Farey. *Period. Math. Hungar. 23* (1991), 75–86.

[36] LEFÈVRE, V. An algorithm that computes a lower bound on the distance between a segment and $\mathbf{Z}^2$. *Rapport de recherche LIP 18* (1997).

[37] LIANG, F. M. A short proof of the $3d$ distance theorem. *Discrete Math. 28* (1979), 325–326.

[38] MARZEC, C. and J. KAPPRAFF. Properties of maximal spacing on a circle related to phyllotaxis and to the golden mean. *J. Theor. Biol. 103* (1983), 201–226.

[39] MORSE, M. and G. A. HEDLUND. Symbolic dynamics. *Amer. J. Math. 60* (1938), 815–866.

[40] MORSE, M. and G. A. HEDLUND. Symbolic dynamics II: Sturmian trajectories. *Amer. J. Math. 62* (1940), 1–42.

[41] RAUZY, G. Suites à termes dans un alphabet fini. *Sém. de Théorie des Nombres de Bordeaux* (1983), 25-01–25-16.

[42] VAN RAVENSTEIN, T. On the discrepancy of the sequence formed from multiples of an irrational number. *Bull. Austral. Math. Soc. 31* (1985), 329–338.

[43] —— Optimal spacing of points on a circle. *Fibonacci Quart. 27* (1989), 18–24.

[44] —— The three gap theorem (Steinhaus conjecture). *J. Austral. Math. Soc. Ser. A 45* (1988), 360–370.

[45] VAN RAVENSTEIN, T. G. WINLEY and K. TOGNETTI. Characteristics and the three gap theorem. *Fibonacci Quart. 28* (1990), 204–214.

[46] ROTE, G. Sequences with subword complexity $2n$. *J. Number Theory 46* (1994), 196–213.

[47] SHALLIT, J. O. Automaticity IV: Sequences, sets, and diversity. *J. Théor. Nombres Bordeaux 8* (1996), 347–367.

[48] SIEGEL, A. Théorème des trois longueurs et suites sturmiennes: mots d'agencement des longueurs. Preprint.

[49] SLATER, N. B. The distribution of the integers N for which $\{\theta N\} < \phi$. *Proc. Cambridge Philos. Soc. 46* (1950), 525–534.

[50] —— Distribution problems and physical applications. *Compositio Math. 16* (1964), 176–183.

[51] —— Gaps and steps for the sequence $n\theta$ mod 1. *Proc. Cambridge Philos. Soc. 63* (1967), 1115–1123.

[52] SÓS, V. T. On the theory of diophantine approximations I. *Acta Math. Acad. Sci. Hungar. 8* (1957), 461–472.

[53] —— On the distribution mod 1 of the sequence $n\alpha$. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math. 1* (1958), 127–134.

[54] STOLARSKY, K. Beatty sequences, continued fractions, and certain shift operators. *Canad. Math. Bull. 19* (1976), 473–482.

[55] SURÁNYI, J. Über die Anordnung der Vielfachen einer reellen Zahl mod 1. *Ann. Univ. Sci. Budapest, Eötvös Sect. Math. 1* (1958), 107–111.

[56] ŚWIERCZKOWSKI, S. On successive settings of an arc on the circumference of a circle. *Fund. Math. 46* (1958), 187–189.

[57] VUILLON, L. Contribution à l'étude des pavages et des surfaces discrétisées. *Theoret. Comput. Sci.*, to appear.

[58] WOLFF, A. and J. PITMAN. Lattice points in strips. Tech. Rep., University of Adelaide, Australia, 1990.

(Reçu le 10 janvier 1997; version révisée reçue le 12 février 1998)

Pascal Alessandri
Valérie Berthé

I. M. L.
Case 907
163, av. de Luminy
F-13288 Marseille Cedex 09
France
e-mail: berthe@iml.univ-mrs.fr