

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 44 (1998)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE COMPLEX GEOMETRY OF THE LAGRANGE TOP
Autor: Gavrilov, Lubomir / ZHIVKOV, Angel
Kapitel: 5. The Lagrange top and the non-linear Schrödinger equation
DOI: <https://doi.org/10.5169/seals-63901>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. THE LAGRANGE TOP AND THE NON-LINEAR SCHRÖDINGER EQUATION

Our final remark concerns a previously unknown relation between the real solutions of the Lagrange top and the one-gap solutions of the nonlinear Schrödinger equation

$$(NLS^\pm) \quad u_{xx} = iu_t \pm 2|u|^2u.$$

In the physical applications both forms of (NLS) are of interest. Comparing Theorem 2.2 to the results of Previato [20] we note that the invariant manifolds of one-gap solutions of the NLS equation are isomorphic to the invariant manifolds of the Lagrange top. This relation can be made explicit if we compare the expressions for the solutions found in Theorem 3.4 to the well known formulae for $u(x, t)$ (cf. [5, 20]). We shall see that the S^\pm -real solutions of the Lagrange top give also one-gap solutions of NLS^\pm equation. Recall that, according to the preceding section, an S^- -real solution is a usual real solution of the Lagrange top (2), and that an S^+ -real solution is a real solution of the system (60).

Let X_E , X_{Ω_3} be the Hamiltonian vector fields (2) and (3) respectively and put

$$\frac{\partial}{\partial x} = \frac{1}{2}X_E, \quad \frac{\partial}{\partial t} = \frac{1}{4}(m-1)\Omega_3 X_E + \frac{1}{8}(2h_3 - (3m+1)\Omega_3^2)X_{\Omega_3}.$$

As $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial t}$ define translation invariant vector fields on the generalized Jacobian $J(C_h; \infty^\pm)$ then fixing an arbitrary point for origin we may introduce (x, t) coordinates on $J(C_h; \infty^\pm)$ (and hence on the complex invariant manifold T_h). If the real part T_h^R of T_h is not empty, then we shall choose for origin a real point. As the real vector fields $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial t}$ are tangent to the Liouville torus T_h^R , then (x, t) provide real affine coordinates on it. Denote, lastly, by $u^-(x, t)$ the restriction of the function $\bar{\epsilon}\Omega_1 + \epsilon\Omega_2$ on the Liouville torus T_h^R of the Lagrange top (2).

Similarly, let $u^+(x, t)$ be the restriction of the function $\bar{\epsilon}\Omega_1 + \epsilon\Omega_2$ on a connected component of the S^+ -real part of $J(C_h; \infty^\pm)$. If the origin belongs to this component too, then as above we conclude that $x, t \in \mathbf{R}$.

PROPOSITION 5.1. *The functions $u^+(x, t)$ and $u^-(x, t)$ satisfy NLS^+ and NLS^- respectively.*

The proof of the above Proposition is a straightforward computation (compare with [20], Theorem 2.2). From the definition of u^\pm we get

$\bar{u}^- = \bar{\epsilon}\Omega_2 + \epsilon\Omega_1$ and $\bar{u}^+ = -\bar{\epsilon}\Omega_2 - \epsilon\Omega_1$. It follows that $|u^\pm|^2 = \mp(\Omega_1^2 + \Omega_2^2)$ and it is easy to check that

$$u_{xx}^\pm = iu_t^\pm \pm 2|u^\pm|^2 u^\pm$$

is equivalent to the system

$$\begin{aligned} (\Omega_1)_{xx} + (\Omega_2)_t &= \pm 2\Omega_1(\Omega_1^2 + \Omega_2^2) \\ (\Omega_2)_{xx} - (\Omega_1)_t &= \pm 2\Omega_2(\Omega_1^2 + \Omega_2^2) \end{aligned}$$

where Ω_1, Ω_2 are defined on the S^\pm -real part of T_h respectively. Using (2) we get for the derivatives along X_E

$$\ddot{\Omega}_1 + (m-1)\Omega_3\dot{\Omega}_2 = -m\Omega_1\Omega_3^2 - \Omega_1\Gamma_3$$

and as

$$\Gamma_3 = \frac{1}{2}(\Omega_1^2 + \Omega_2^2 + (1+m)\Omega_3^2) - E,$$

then

$$\ddot{\Omega}_1 + (m-1)\Omega_3\dot{\Omega}_2 = -\frac{1}{2}\Omega_1(\Omega_1^2 + \Omega_2^2) + \Omega_1(E - \frac{3m+1}{2}\Omega_3^2).$$

Finally, as $X_{\Omega_3}\Omega_2 = -\Omega_1$ we conclude that

$$\begin{aligned} (\Omega_1)_{xx} + (\Omega_2)_t &= -2\Omega_1(\Omega_1^2 + \Omega_2^2) \\ (\Omega_2)_{xx} - (\Omega_1)_t &= -2\Omega_2(\Omega_1^2 + \Omega_2^2). \end{aligned}$$

This proves also that u^+ is a solution of NLS^+ (we just have to substitute $\Omega_1 \mapsto i\Omega_1$, $\Omega_2 \mapsto i\Omega_2$). \square