

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 45 (1999)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE CONSTRUCTION OF GENERALIZED JACOBIANS
Autor: Fu, Lei
Kapitel: 1. A THEOREM OF GROTHENDIECK
DOI: <https://doi.org/10.5169/seals-64440>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

1. A THEOREM OF GROTHENDIECK

The following theorem is a special case of Grothendieck's theorems, and the proof can be found in [Mu] §5, [H] §3.12, or [EGA] III, §7.7.5, 7.9.4.

THEOREM 1.1. *Let $q: V \rightarrow T$ be a proper flat morphism of noetherian schemes and let \mathcal{L} be an invertible sheaf on V . For each $t \in T$ denote the fiber $V \otimes_T \text{spec}(k(t))$ of q at t by V_t , where $k(t)$ is the residue field of T at t . Denote the inverse image of \mathcal{L} on V_t by \mathcal{L}_t .*

- (a) *The function $t \mapsto \chi(\mathcal{L}_t) = \sum_i (-1)^i \dim_{k(t)} H^i(V_t, \mathcal{L}_t)$ is locally constant on T .*
- (b) *For each i , the function $t \mapsto \dim_{k(t)} H^i(V_t, \mathcal{L}_t)$ on T is upper semicontinuous.*
- (c) *If T is reduced and connected and if $t \mapsto \dim_{k(t)} H^i(V_t, \mathcal{L}_t)$ is a constant function on T , then $R^i q_* \mathcal{L}$ is a locally free sheaf on T and the map $R^i q_* \mathcal{L} \otimes_{\mathcal{O}_T} k(t) \rightarrow H^i(V_t, \mathcal{L}_t)$ is an isomorphism.*
- (d) *If $H^1(V_t, \mathcal{L}_t) = 0$ for all $t \in T$, then $R^1 q_* \mathcal{L} = 0$ and $q_* \mathcal{L}$ is a locally free sheaf. Moreover the formation of $q_* \mathcal{L}$ commutes with any base change.*

2. RELATIVE EFFECTIVE CARTIER DIVISORS

Let $q: X \rightarrow T$ be a morphism of noetherian schemes. A *relative effective Cartier divisor* on X/T is an effective Cartier divisor on X that is flat over T when regarded as a closed subscheme of X . When $T = \text{spec}(R)$ is affine, a closed subscheme D of X is a relative effective Cartier divisor if and only if there exists an open affine covering $U_i = \text{spec}(R_i)$ of X and $g_i \in R_i$ such that

- (a) $D \cap U_i = \text{spec}(R_i/(g_i))$;
- (b) g_i is not a zero divisor;
- (c) $R_i/(g_i)$ is flat over R .

REMARK 2.1. Let D be an effective Cartier divisor on X/T , let $\mathcal{I}(D)$ be the sheaf of ideals defining D , and let $\mathcal{L}(D)$ be the invertible sheaf corresponding to D . We have $\mathcal{L}(D) = \mathcal{I}(D)^{-1}$. The inclusion $\mathcal{I}(D) \subset \mathcal{O}_X$ induces $\mathcal{O}_X \subset \mathcal{I}(D)^{-1} = \mathcal{L}(D)$, hence a section s_D of $\mathcal{L}(D)$.