Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 45 (1999)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HARMONIC ANALYSIS ON VECTOR BUNDLES OVER Sp(1,n)/Sp(1)xSp(n)

Autor: van Dijk, G. / PASQUALE, A.

Kapitel: 3. The τ_l -spherical functions

DOI: https://doi.org/10.5169/seals-64447

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 27.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. THE τ_l-SPHERICAL FUNCTIONS

Let $\mathfrak{U}(\mathfrak{g})$ be the universal enveloping algebra of the complexification $\mathfrak{g}_{\mathbf{C}}$ of $\mathfrak{g}=\mathfrak{sp}(1,n)$. The elements of $\mathfrak{U}(\mathfrak{g})$ are considered as left-invariant differential operators on $G=\mathrm{Sp}(1,n)$ acting on C^∞ functions on the right:

$$f(g;X):=\frac{d}{dt}f(g\exp tX)\Big|_{t=0}\qquad (f\in C^\infty(G),\ X\in\mathfrak{g},\ g\in G).$$

We adopt Harish-Chandra's notation $f(g;D)$ for the image of $f\in C^\infty(G)$ under the right action of $D\in\mathfrak{U}(\mathfrak{g})$. The set of K-invariant elements of $\mathfrak{U}(\mathfrak{g})$ is denoted by $\mathfrak{U}(\mathfrak{g})^K$.

3.1. THEOREM ([Go], Theorems 8, 10 and 14; [GaV], Theorems 1.3.14, 1.4.5 and Proposition 1.4.4). *Let $l\in\mathbf{N}/2$ be fixed. Let ζ be a complex-valued continuous function on G satisfying (2.9), (2.10) and $\zeta(e)=1$ (e is the unit element of G). The following statements are mutually equivalent.*

1. *The mapping $f\mapsto\int_G f(g)\zeta(g)\,dg$ is an algebra homomorphism of $\mathcal{D}(G;\chi_l)$ into $\mathbf{C}$.*
2. *ζ satisfies the functional equation*

$$\int_K \zeta(kg_1k^{-1}g_2)\,dk=\zeta(g_1)\zeta(g_2) \tag{3.15}$$

for all $g_1,g_2\in G$.

3. *ζ is a common eigenfunction of the elements of $\mathfrak{U}(\mathfrak{g})^K$.*

A function ζ satisfying the equivalent conditions of Theorem 3.1 is called a *spherical function of type τ_l (and height 1)* or briefly a *τ_l-spherical function*[1]).

Observe that Condition 3 implies in particular that every τ_l-spherical function is analytic on G because $\mathfrak{U}(\mathfrak{g})^K$ contains an elliptic differential operator.

---

[1]) Our definition of τ_l-spherical function ζ is formally less restrictive than Godement's, which also requires ζ to be quasi-bounded with respect to some seminorm on G (cf. [Go], p. 519). It is possible to show (e.g. [GaV], Theorem 1.3.14) that each of the conditions for ζ given by Theorem 3.1 is equivalent to the existence of an irreducible Fréchet representation $(T,\mathcal{H})$ – with a d_l-dimensional K-isotypic subspace $\mathcal{H}(\tau_l)$ of type τ_l – for which

$$(*)\qquad \zeta=\frac{1}{d_l}\operatorname{tr}[E(\tau_l)TE(\tau_l)],$$

$E(\tau_l)$ being the projection of $\mathcal{H}$ onto $\mathcal{H}(\tau_l)$. $(T,\mathcal{H})$ can be chosen to be a Banach representation of G if and only if ζ is quasi-bounded with respect to some seminorm on G. In Section 5 we will determine, for each τ_l-spherical function ζ, an irreducible *Hilbert* representation $(T,\mathcal{H})$ for which $(*)$ holds. It follows, in particular, that the condition of quasi-boundedness is, in our case, automatically satisfied.

For complex-valued functions f on G and F on $\mathbf{R}$, we set

$$f^*(g) = \overline{f(g^{-1})} \qquad (g \in G)$$
$$F^*(t) = \overline{F(-t)} \qquad (t \in \mathbf{R}).$$

The τ_l-*Abel transform* of $f \in \mathcal{D}(G;\chi_l)$ is the function $\mathcal{A}_l f$ on $\mathbf{R}$ defined by

$$\mathcal{A}_l f(t) = \frac{1}{d_l^2}\, e^{\rho t} \int_N f(a_t n)\, dn\,. \tag{3.16}$$

Its properties are summarized in the following proposition.

3.2. PROPOSITION. *For all $f \in \mathcal{D}(G;\chi_l)$, $\mathcal{A}_l f$ is a C^∞ function on $\mathbf{R}$ with compact support. If $f, f_1, f_2 \in \mathcal{D}(G;\chi_l)$ and $a_1, a_2 \in \mathbf{C}$, then*

$$(\mathcal{A}_l f)^* = \mathcal{A}_l(f^*)\,, \tag{3.17}$$

$$\mathcal{A}_l(a_1 f_1 + a_2 f_2) = a_1 \mathcal{A}_l f_1 + a_2 \mathcal{A}_l f_2\,, \tag{3.18}$$

$$\mathcal{A}_l(f_1 * f_2) = \mathcal{A}_l f_1 * \mathcal{A}_l f_2\,. \tag{3.19}$$

Formula (3.17) is equivalent to the fact that $\mathcal{A}_l f$ is an even function.

Proof. Formulas (3.17)–(3.19) are immediately proven by passing to $\mathcal{D}(G;\tau_l)$. For the last statement, recall that $f(g^{-1}) = f(g)$ for $f \in \mathcal{D}(G;\chi_l)$. □

The following lemma relates our definition of $\mathcal{A}_l$ to the definition often found in the literature (cf. e.g. [W2], p. 34).

3.3. LEMMA. *For $f \in \mathcal{D}(G;\chi_l)$ one has*

$$\int_N f(k a_t n)\, dn = \frac{1}{d_l}\chi_l(k) \int_N f(a_t n)\, dn\,.$$

Proof. Let $F \in \mathcal{D}(G;\tau_l)$. Then $\int_N F(a_t n)\, dn$ commutes with $\tau(m)$ $(m \in M)$ so with $\tau(k_1)$ $(k_1 \in K_1)$, hence is a scalar multiple of the identity. The lemma follows by taking traces. □

We now use the τ_l-Abel transform to construct τ_l-spherical functions. Because of Proposition 3.2, for any complex number s, the map

$$\lambda_s : f \longmapsto \int_{-\infty}^{\infty} \mathcal{A}_l f(t) e^{-st}\, dt \tag{3.20}$$

is an algebra homomorphism of $\mathcal{D}(G;\chi_l)$ into $\mathbf{C}$.

Set

$$\alpha_{l,s}(ka_tn) = \frac{1}{d_l}\chi_l(k)e^{-(s+\rho)t}\,. \tag{3.21}$$

Since $f = f * d_l\chi_l$ and $\chi_l(k^{-1}) = \chi_l(k)$ for $k \in K$, for every $f \in \mathcal{D}(G;\chi_l)$

$$\begin{aligned}\lambda_s(f) &= \frac{1}{d_l}\int_K\int_{-\infty}^{\infty}\int_N f(ka_tn)\chi_l(k)e^{(-s+\rho)t}\,dk\,dt\,dn\\ &= \int_G f(g)\alpha_{l,s}(g)\,dg\\ &= \int_G f(g)\int_K \alpha_{l,s}(kgk^{-1})\,dk\,dg\\ &= \int_G f(g)\zeta_{l,s}(g)\,dg \qquad (3.22)\end{aligned}$$

with

$$\zeta_{l,s} := \int_K \alpha_{l,s}(kgk^{-1})\,dk\,. \tag{3.23}$$

One easily checks that $\zeta_{l,s}$ satisfies $\zeta_{l,s} = \zeta_{l,s}{}^0$, $\zeta_{l,s}*d_l\chi_l = \zeta_{l,s}$ and $\zeta_{l,s}(e) = 1$. Thus $\zeta_{l,s}$ is a τ_l-spherical function. It will be shown in the next section that any τ_l-spherical function is of the form (3.24).

By Remark 2.3, we have

$$\zeta_{l,s}(g) = \frac{1}{d_l}\chi_l(k_1)\zeta_{l,s}(a_t) \quad \text{for } g = k_1k_2a_tk_2'\,, \tag{3.24}$$

so $\zeta_{l,s}$ is uniquely determined by its restriction to A.

4. The differential equation for the τ_l-spherical functions

For a subalgebra $\mathfrak{u}$ of $\mathfrak{g}$, let $\mathfrak{u}_{\mathbf{C}}$ denote the complex subalgebra of $\mathfrak{g}_{\mathbf{C}}$ generated by $\mathfrak{u}$. The universal enveloping algebra $\mathfrak{U}(\mathfrak{u})$ of $\mathfrak{u}_{\mathbf{C}}$ is considered as a subalgebra of $\mathfrak{U}(\mathfrak{g})$.

The representation τ_l of K_1 induces differentiated representations of the Lie algebra $\mathfrak{k}_1$ of K_1 and of the universal enveloping algebra $\mathfrak{U}(\mathfrak{k}_1)$ of $(\mathfrak{k}_1)_{\mathbf{C}}$. We indicate these representations with the same letter τ_l. Let $\mathfrak{k}_2$ be the Lie algebra of K_2. Every element $Y \in \mathfrak{k}_{\mathbf{C}}$ can be uniquely decomposed as