Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 45 (1999)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE CONSTRUCTION OF GENERALIZED JACOBIANS

Autor: Fu, Lei

Kapitel: 4. From birational groups to algebraic groups

DOI: https://doi.org/10.5169/seals-64440

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 01.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Note that Φ is a regular morphism defined on U and Θ is a regular morphism defined on V. Since

$$\Phi \Theta(D_1, D_2) = (D_1, D_2)$$
 and $\Theta \Phi(D_1, D_2) = (D_1, D_2)$

whenever the left-hand sides are defined, the maps Φ and Θ induce regular morphisms $\Phi \colon U \cap \Phi^{-1}(V) \to V \cap \Theta^{-1}(U)$ and $\Theta \colon V \cap \Theta^{-1}(U) \to U \cap \Phi^{-1}(V)$. To show that Φ and Θ are birational inverses to each other, it is enough to check that $U \cap \Phi^{-1}(V)$ and $V \cap \Theta^{-1}(U)$ are non-empty.

Note that $(D_1, D_2) \in U \cap \Phi^{-1}(V)$ if and only if $(D_1, D_2) \in U$ and

$$l_{\mathfrak{m}}(m(D_1, D_2) - D_1 + \pi P_0) = 1, \quad l(m(D_1, D_2) - D_1 + \pi P_0 - \mathfrak{m}) = 0.$$

Since $m(D_1, D_2) \sim_{\mathfrak{m}} D_1 + D_2 - \pi P_0$, the above equations are equivalent to

$$l_{\mathfrak{m}}(D_2) = 1, \quad l(D_2 - \mathfrak{m}) = 0.$$

Applying Lemma 3.3 to the divisor $D_0 = 0$, we conclude that the set

$$V_0 = \{ D \in (X - S)^{(\pi)} \mid l_{\mathfrak{m}}(D) = 0, \quad l(D - \mathfrak{m}) = 0 \}$$

is open and non-empty. Since $(X-S)^{(\pi)} \times (X-S)^{(\pi)}$ is irreducible, the set $U \cap ((X-S)^{(\pi)} \times V_0)$ is also open and non-empty. This set is exactly $U \cap \Phi^{-1}(V)$. So $U \cap \Phi^{-1}(V)$ is non-empty.

Similarly $V \cap \Theta^{-1}(U)$ is also non-empty. This completes the proof of the proposition.

4. From Birational Groups to algebraic groups

Let k be an algebraically closed field, let V be a connected nonsingular variety over k, and let $m\colon V\times V\to V$, $(a,b)\mapsto ab$ be a rational map satisfying (ab)c=a(bc). Assume the rational maps $\Phi(a,b)=(a,ab)$ and $\Psi(a,b)=(b,ab)$ are birational. Then there exist open subsets X_Φ , Y_Φ , X_Ψ and Y_Ψ in $V\times V$ such that Φ induces an isomorphism $X_\Phi\cong Y_\Phi$ and Ψ induces an isomorphism $X_\Psi\cong Y_\Psi$. Put $Z=X_\Phi\cap Y_\Phi\cap X_\Psi\cap Y_\Psi$.

It is convenient to write the formulae for Φ^{-1} and Ψ^{-1} as $\Phi^{-1}(a,b) = (a,a^{-1}b)$ and $\Psi^{-1}(a,b) = (ba^{-1},a)$.

LEMMA 4.1. Replacing V by an open subset, we may assume the two projections $p_i: Z \to V$ (i = 1, 2) are surjective.

Proof. Note that the two projections $p_i \colon V \times V \to V$, (i = 1, 2) are flat since $V \to \operatorname{spec}(k)$ is flat. So the p_i are open by [EGA] IV, §2.4.6. Hence the $p_i(Z)$ are open. Let $V' = p_1(Z) \cap p_2(Z)$. We claim V' has the property stated in the lemma. Let C = V - V' and let $A = (C \times V) \cup (V \times C)$. The subset X_{Φ}' of $V' \times V'$ corresponding to X_{Φ} is the complement in X_{Φ} of $S = (X_{\Phi} \cap A) \cup \Phi^{-1}(Y_{\Phi} \cap A)$. We claim that if the fiber of $p_1 \colon X_{\Phi} \to V$ at $v \in V$ is contained in S, then $v \in C$. Thus $p_1 \colon X_{\Phi}' \to V'$ is surjective.

Let us prove the claim. Assume $(v \times V) \cap X_{\Phi} \subset S$, but $v \notin C$. We have

$$(v \times V) \cap X_{\Phi} \subset S \subset A \cup \Phi^{-1}(A) \subset (C \times V) \cup (V \times C) \cup \Phi^{-1}(C \times V) \cup \Phi^{-1}(V \times C).$$

Since V is irreducible, we must have

$$(v \times V) \cap X_{\Phi} \subset C \times V$$
, $V \times C$, $\Phi^{-1}(C \times V)$, or $\Phi^{-1}(V \times C)$.

Since $v \notin C$, we have

$$(v \times V) \cap X_{\Phi} \not\subset C \times V, \quad \Phi^{-1}(C \times V).$$

So

$$(v \times V) \cap X_{\Phi} \subset V \times C \text{ or } \Phi^{-1}(V \times C)$$
.

Assume $(v \times V) \cap X_{\Phi} \subset V \times C$. Note that since $v \notin C$, we have $v \in V'$. Hence $(v \times V) \cap X_{\Phi}$ is not empty. So we have

$$\dim V = \dim((v \times V) \cap X_{\Phi}) = \dim(((v \times V) \cap X_{\Phi}) \cap (V \times C))$$

$$\leq \dim(v \times C) < \dim V,$$

that is, $\dim V < \dim V$. This is impossible.

Assume $(v \times V) \cap X_{\Phi} \subset \Phi^{-1}(V \times C)$. Then $\Phi((v \times V) \cap X_{\Phi}) \subset V \times C$. Since Φ is birational, we have

$$\dim V = \dim \Phi((v \times V) \cap X_{\Phi}) = \dim(\Phi((v \times V) \cap X_{\Phi}) \cap (V \times C))$$

$$\leq \dim(v \times C) < \dim V.$$

which is again impossible. So we must have $v \in C$.

Next we show that if the fiber of $p_2: X_{\Phi} \to V$ at $v \in V$ is contained in S, then $v \in C$, and hence $p_2: X_{\Phi}' \to V'$ is surjective.

Assume $(V \times v) \cap X_{\Phi} \subset S$ but $v \notin C$. As before we have

$$(V \times v) \cap X_{\Phi} \subset C \times V$$
, $V \times C$, $\Phi^{-1}(C \times V)$ or $\Phi^{-1}(V \times C)$.

Since $v \notin C$, we have $(V \times v) \cap X_{\Phi} \not\subset V \times C$. By counting dimensions, one can show $(V \times v) \cap X_{\Phi} \not\subset C \times V$. Since $\Phi^{-1}(C \times V) \subset C \times V$, we have $(V \times v) \cap X_{\Phi} \not\subset \Phi^{-1}(C \times V)$. So we can only have $(V \times v) \cap X_{\Phi} \subset \Phi^{-1}(V \times C)$. Then we have a rational map

$$V \xrightarrow{\iota_1} (V \times v) \cap X_{\Phi} \xrightarrow{\Phi} V \times C \xrightarrow{p_2} C$$

where $\iota_1(x) = (x, v)$. This map $p_2\Phi\iota_1 \colon V \to C$ is nothing but $x \mapsto xv$ and it is birational. (Its birational inverse is $p_1\Psi^{-1}\iota_2$, where $\iota_2(x) = (v, x)$.) So V is birational to C. This is impossible since dim $V \neq \dim C$. So we must have $v \in C$. This finishes the proof of the surjectivity of $p_2 \colon X_{\Phi}' \to V'$.

Similarly $p_i: X'_{\Phi}, Y_{\Phi'}, X'_{\Psi}, Y'_{\Psi} \to V'$ are surjective. Since the fibers of $p_i: V \times V \to V$ are irreducible, the projection $p_i: Z' = X'_{\Phi} \cap Y'_{\Phi} \cap X'_{\Psi} \cap Y'_{\Psi} \to V'$ is also surjective.

Having replaced V as in Lemma 4.1, we may assume V satisfies the following properties:

PROPERTY 4.2. There exists an open set $Z \subset V \times V$ such that Φ, Φ^{-1}, Ψ , and Ψ^{-1} are defined on Z, the restrictions $\Phi|_Z$ and $\Psi|_Z$ are open immersions, and the projections $p_i \colon Z \to V$ are surjective. Hence for every $v \in V$, the maps Φ , Φ^{-1} , Ψ and Ψ^{-1} are defined at (v,x) and at (x,v), provided x is generic, i.e. lies in an open set.

LEMMA 4.3. Assume 4.2 holds. Denote the closure of the graph of m in $V \times V \times V$ by Γ . Then the projections $p_{ij} \colon \Gamma \to V \times V$ $(1 \le i < j \le 3)$ are open immersions.

Proof. By [EGA] III, §4.4.9, it suffices to show that the maps p_{ij} are set-theoretically injective. Let x be a point of V. The two rational maps $\Gamma \to V$ defined by

$$(a, b, c) \mapsto (xa)b$$
 and $(a, b, c) \mapsto xc$

are equal by the associative law. Let (a,b,c), $(a,b,c') \in \Gamma$. Choose x so that (xa)b is defined and (x,c), $(x,c') \in Z$. Then xc = (xa)b = xc'. Hence $\Phi(x,c) = \Phi(x,c')$. Since Φ is an open immersion on Z, we have (x,c) = (x,c'). Hence c = c'. This shows that $p_{12} \colon \Gamma \to V \times V$ is injective. Similarly one can show the other projections are injective.

We will now expand V to the group we want by glueing translates of V. Let s be a point of V and let V_s be a copy of V thought of as the translate $V_s = \{vs \mid v \in V\}$. The subset $W_s = (V \times s \times V) \cap \Gamma$ is closed in $V \times s \times V \cong V \times V$, and the two projections $W_s \to V$ are open immersions because they are the base extensions of the open immersions $p_{ij} \colon \Gamma \to V \times V$ by the base changes $V \times s \to V \times V$ and $s \times V \to V \times V$, respectively. Therefore W_s defines glueing data and yields a separated scheme $V' = V \cup_{W_s} V_s$.

LEMMA 4.4. V is an open dense subset of V' and V' satisfies 4.2.

Proof. Since xs is defined for generic $x \in V$, the set $V \cap V_s$ is not empty. So V' is irreducible and V is dense in V'. We have

$$V' \times V' = (V \times V) \cup (V \times V_s) \cup (V_s \times V) \cup (V_s \times V_s).$$

For every point $v \in V$, denote by v_s the point v considered as a point in V_s . Note that if $(v,s) \in Z$, then $vs \in V$ and $v_s \in V_s$ are glued together in V'. Define $R_s \colon V \to V_s$ by $v \mapsto v_s$. Let

$$W_1 = \{(a, b) \in V \times V \mid (a, b), (s, a) \text{ and } (b, sa^{-1}) \text{ are all in } Z\}.$$

This is a non-empty open subset of Z. Take $Z_1 = (\mathrm{id} \times R_s)(W_1) \subset V \times V_s$. We define Φ , Ψ , Φ^{-1} and Ψ^{-1} on Z_1 by

$$\Phi(a, b_s) = (a, (ab)_s) \in V \times V_s,$$

$$\Psi(a, b_s) = (b_s, (ab)_s) \in V_s \times V_s,$$

$$\Phi^{-1}(a, b_s) = (a, (a^{-1}b)_s) \in V \times V_s,$$

$$\Psi^{-1}(a, b_s) = (b(sa^{-1}), a) \in V \times V$$

for any $(a, b_s) \in Z_1$. Let

$$W_2 = \{(a,b) \in V \times V$$

$$|(a,b), (s,b), (a,sb), (s,a^{-1}b) \text{ and } (bs^{-1},a) \text{ are all in } Z\}.$$

This is a non-empty open subset of Z. Take $Z_2 = (R_s \times id)(W_2) \subset V_s \times V$. We define Φ , Ψ , Φ^{-1} , and Ψ^{-1} on Z_2 by

$$\Phi(a_s, b) = (a_s, a(sb)) \in V_s \times V,$$

$$\Psi(a_s, b) = (b, a(sb)) \in V \times V,$$

$$\Phi^{-1}(a_s, b) = (a_s, s^{-1}(a^{-1}b)) \in V_s \times V,$$

$$\Psi^{-1}(a_s, b) = ((bs^{-1})a^{-1}, a_s) \in V \times V_s$$

for any $(a_s, b) \in \mathbb{Z}_2$.

Let $Z'=Z\cup Z_1\cup Z_2$. It is an open subset of $V'\times V'$, and Φ , Ψ , Φ^{-1} , Ψ^{-1} are defined on it. One can show that $\Phi|_{Z'}$ and $\Psi|_{Z'}$ are open

immersions. Given $v \in V'$, we need to show there exists $x \in V'$ such that (x, v) and (v, x) are in Z'. This is true if $v \in V$ by the property of Z. If $v \in V_s$, then $v = a_s$ for some $a \in V$. We leave it to the reader to show that $(x, a_s) \in Z_1$ and $(a_s, x) \in Z_2$ for generic x in V. This completes the proof of the lemma.

The above lemma allows us to replace V by V', hence to expand V whenever there exists a point s in V such that vs is not defined for all $v \in V$, and we can expand V' if there exists a point $s' \in V'$ such that v's' is not defined for all $v' \in V'$. Denote the result of finitely many such expansions also by V', and let $U \subset V \times V \times V'$ be the closure of Γ . By Lemma 4.3 applied to V', the projection $p_{12} \colon U \to V \times V$ is an open immersion. Its image is the set of points (a,b) such that $m \colon V \times V \to V'$ is defined at (a,b). If $V \times s \not\subset p_{12}(U)$ for some point s in V, then replacing V' by $V' \cup V_s'$ increases both V' and $p_{12}(U)$. Using noetherian induction on open subschemes of $V \times V$, we may assume that after finitely many expansions, $V \times s \subset p_{12}(U)$ for all points $s \in V$. Then we have $p_{12}(U) = V \times V$.

PROPOSITION 4.5. Let V, V', and U be as above. If $p_{12}(U) = V \times V$, then the operation $m: V' \times V' \to V'$ is everywhere defined on V' and makes V' an algebraic group.

Proof. Take (a',b') in $V' \times V'$. Choose a point x so that a'x and $x^{-1}b'$ are both defined and lie in V. Then we can define $m(a',b')=(a'x)(x^{-1}b')$. Similarly one can define $a'^{-1}b'$ and $b'a'^{-1}$. In this way we extend m, Φ , Ψ , Φ^{-1} and Ψ^{-1} to $V' \times V'$. The verification of the group axioms is routine and is omitted.

5. FUNDAMENTAL PROPERTIES OF GENERALIZED JACOBIANS

Keep the notations in §3. We have proved that there is a birational group structure on $(X-S)^{(\pi)}$. The algebraic group associated to this birational group is called the *generalized jacobian* of $X_{\mathfrak{m}}$ and is denoted by $J_{\mathfrak{m}}$. It is a commutative algebraic group.

Let D_0 be a divisor on X prime to S of degree 0. By Lemma 3.3, the set

$$V_{D_0} = \{ D \in (X - S)^{(\pi)} \mid l_{\mathfrak{m}}(D + D_0) = 1, \quad l(D + D_0 - \mathfrak{m}) = 0 \}$$

is a non-empty open subset of $(X - S)^{(\pi)}$. We have the following