

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 45 (1999)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HARMONIC ANALYSIS ON VECTOR BUNDLES OVER
Sp(1,n)/Sp(1)xSp(n)
Autor: van Dijk, G. / PASQUALE, A.
Kapitel: 4. The differential equation for the τ_I -spherical functions
DOI: <https://doi.org/10.5169/seals-64447>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

is an algebra homomorphism of $\mathcal{D}(G; \chi_l)$ into \mathbf{C} .

Set

$$(3.21) \quad \alpha_{l,s}(ka_t n) = \frac{1}{d_l} \chi_l(k) e^{-(s+\rho)t}.$$

Since $f = f * d_l \chi_l$ and $\chi_l(k^{-1}) = \chi_l(k)$ for $k \in K$, for every $f \in \mathcal{D}(G; \chi_l)$

$$\begin{aligned} \lambda_s(f) &= \frac{1}{d_l} \int_K \int_{-\infty}^{\infty} \int_N f(ka_t n) \chi_l(k) e^{(-s+\rho)t} dk dt dn \\ &= \int_G f(g) \alpha_{l,s}(g) dg \\ &= \int_G f(g) \int_K \alpha_{l,s}(kgk^{-1}) dk dg \\ (3.22) \quad &= \int_G f(g) \zeta_{l,s}(g) dg \end{aligned}$$

with

$$(3.23) \quad \zeta_{l,s} := \int_K \alpha_{l,s}(kgk^{-1}) dk.$$

One easily checks that $\zeta_{l,s}$ satisfies $\zeta_{l,s} = \zeta_{l,s}^0$, $\zeta_{l,s} * d_l \chi_l = \zeta_{l,s}$ and $\zeta_{l,s}(e) = 1$. Thus $\zeta_{l,s}$ is a τ_l -spherical function. It will be shown in the next section that any τ_l -spherical function is of the form (3.24).

By Remark 2.3, we have

$$(3.24) \quad \zeta_{l,s}(g) = \frac{1}{d_l} \chi_l(k_1) \zeta_{l,s}(a_t) \quad \text{for } g = k_1 k_2 a_t k_2' ,$$

so $\zeta_{l,s}$ is uniquely determined by its restriction to A .

4. THE DIFFERENTIAL EQUATION FOR THE τ_l -SPHERICAL FUNCTIONS

For a subalgebra \mathfrak{u} of \mathfrak{g} , let $\mathfrak{u}_{\mathbf{C}}$ denote the complex subalgebra of $\mathfrak{g}_{\mathbf{C}}$ generated by \mathfrak{u} . The universal enveloping algebra $\mathfrak{U}(\mathfrak{u})$ of $\mathfrak{u}_{\mathbf{C}}$ is considered as a subalgebra of $\mathfrak{U}(\mathfrak{g})$.

The representation τ_l of K_1 induces differentiated representations of the Lie algebra \mathfrak{k}_1 of K_1 and of the universal enveloping algebra $\mathfrak{U}(\mathfrak{k}_1)$ of $(\mathfrak{k}_1)_{\mathbf{C}}$. We indicate these representations with the same letter τ_l . Let \mathfrak{k}_2 be the Lie algebra of K_2 . Every element $Y \in \mathfrak{k}_{\mathbf{C}}$ can be uniquely decomposed as

$Y = Y^{(1)} + Y^{(2)}$ with $Y^{(j)} \in (\mathfrak{k}_j)_C$, $j = 1, 2$. The symbol χ_l will also be used for the \mathbf{C} -linear map on $\mathfrak{U}(\mathfrak{k})$ defined by

$$\chi_l(Y_1 \cdots Y_m) := \text{tr} \left[\tau_l(Y_1^{(1)}) \cdots \tau_l(Y_m^{(1)}) \right]$$

for $Y_1, \dots, Y_m \in \mathfrak{k}_C$.

The Iwasawa decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{a} + \mathfrak{n}$ gives $\mathfrak{U}(\mathfrak{g}) = \mathfrak{U}(\mathfrak{k})\mathfrak{U}(\mathfrak{a})\mathfrak{U}(\mathfrak{n}) \cong \mathfrak{U}(\mathfrak{k}) \otimes \mathfrak{U}(\mathfrak{a}) \oplus \mathfrak{U}(\mathfrak{g})\mathfrak{n}_C$. Let $P: \mathfrak{U}(\mathfrak{g}) \rightarrow \mathfrak{U}(\mathfrak{k}) \otimes \mathfrak{U}(\mathfrak{a})$ be the corresponding projection. For $s \in \mathbf{C}$, let e_s be the \mathbf{C} -linear map on $\mathfrak{U}(\mathfrak{a})$ defined by

$$e_s(L^m) := (-1)^m(s + \rho)^m \quad \text{for every integer } m \geq 0.$$

Define $p_{l,s}: \mathfrak{U}(\mathfrak{g}) \rightarrow \mathbf{C}$ to be the composition $p_{l,s} := \left(\frac{1}{d_l} \chi_l \otimes e_s \right) \circ P$, where as before $d_l = \dim \tau_l$.

4.1. PROPOSITION. *Let $\zeta_{l,s}$ be the function defined by Formula (3.23). For every $D \in \mathfrak{U}(\mathfrak{g})^K$ and $g \in G$*

$$(4.25) \quad \zeta_{l,s}(g; D) = p_{l,s}(D)\zeta_{l,s}(g).$$

Proof. Because of Theorem 3.1, $\zeta_{l,s}$ is an eigenfunction of every $D \in \mathfrak{U}(\mathfrak{g})^K$. The eigenvalue corresponding to $D \in \mathfrak{U}(\mathfrak{g})^K$ is $\zeta_{l,s}(e; D)$ because $\zeta_{l,s}(e) = 1$. Since D is K -invariant, $\zeta_{l,s}(e; D) = \alpha_{l,s}(e; D)$. Write $D = \sum_i y_i x_i + \sum_j n_j$ with $y_i \in \mathfrak{U}(\mathfrak{k})$, $x_i \in \mathfrak{U}(\mathfrak{a})$ and $n_j \in \mathfrak{U}(\mathfrak{g})\mathfrak{n}_C$. Then $\alpha_{l,s}(e; D) = \sum_i \alpha_{l,s}(e; y_i x_i)$ because $\alpha_{l,s}(gn) = \alpha_{l,s}(g)$ for $g \in G$ and $n \in N$. To compute $\alpha_{l,s}(e; y_i x_i)$, assume without loss of generality that $x_i = L^{m_i}$ and that $y_i = Y_1 \cdots Y_m$ with $Y_j \in \mathfrak{k}$. The definition of $\alpha_{l,s}$ gives

$$\alpha_{l,s}(e; y_i x_i) = \frac{1}{d_l} \chi_l(y_i) (-1)^m (s + \rho)^m = p_{l,s}(y_i x_i).$$

Thus $\zeta_{l,s}(e; D) = p_{l,s}(D)$. \square

Let $\delta_l(D)$ denote the τ_l -radial component on $A^+ := \{a_t : t > 0\}$ of the differential operator $D \in \mathfrak{U}(\mathfrak{g})$; that is, the unique differential operator on A^+ satisfying

$$f(a_t; \delta_l(D)) = f(a_t; D)$$

for all $f \in \mathcal{D}(G; \chi_l)$ and $t > 0$. Proposition 4.1 immediately implies

4.2. COROLLARY. *$\zeta_{l,s}$ is an eigenfunction of the τ_l -radial component on A^+ of every K -invariant differential operator on G :*

$$(4.26) \quad \zeta_{l,s}(a_t; \delta_l(D)) = p_{l,s}(D)\zeta_{l,s}(a_t) \quad (D \in \mathfrak{U}(\mathfrak{g})^K, t > 0).$$

We now write (4.26) explicitly in the case D is the Casimir operator ω of \mathfrak{g} . Let B denote the Cartan-Killing form of $\mathfrak{g}_{\mathbf{C}}$ ($\cong \mathfrak{sp}(1+n, \mathbf{C})$). If $X, Y \in \mathfrak{sp}(1, n)$, then

$$B(X, Y) = 4(n+2) \Re \operatorname{tr}(XY)$$

where \Re denotes the quaternionic real part: $\Re q = \frac{q + \bar{q}}{2}$ for $q \in \mathbf{H}$. The bilinear form $B_{\theta}(X, Y) := -B(X, \theta Y)$ is an inner product on \mathfrak{g} . Orthonormality will be considered with respect to B_{θ} .

Let $\{Z_j\}_{j=1}^m$ ($m := 2n^2 + n$) and $\{X_{\beta,j}\}_{j=1}^{m_{\beta}}$ ($\beta \in \{\alpha, 2\alpha\}$) be orthonormal bases in \mathfrak{m} and in \mathfrak{g}_{β} respectively. Define $X_{-\beta,j} = -\theta(X_{\beta,j})$ for $\beta \in \{\alpha, 2\alpha\}$ and $j = 1, \dots, m_{\beta}$. Then $\{X_{-\beta,j}\}_{j=1}^{m_{\beta}}$ is an orthonormal basis for $\mathfrak{g}_{-\beta}$, and $B(X_{\beta,i}, X_{-\beta,j}) = \delta_{ij}$. Moreover, for all $j = 1, \dots, m_{\beta}$, $H_{\beta} := [X_{\beta,j}, X_{-\beta,j}]$ is the unique element of \mathfrak{a} satisfying $B(H_{\beta}, L) = \beta(L)$, i.e.

$$H_{\beta} = \frac{h_{\beta}}{8(n+2)} L \quad \text{with} \quad h_{\beta} = \begin{cases} 1 & \text{if } \beta = \alpha \\ 2 & \text{if } \beta = 2\alpha. \end{cases}$$

Set $H_1 := \frac{L}{\sqrt{8(n+2)}}$, a unit vector in \mathfrak{a} . Then, if $D_{\beta,j} := X_{\beta,j}X_{-\beta,j} + X_{-\beta,j}X_{\beta,j}$, we have (cf. [GaV], p. 132)

$$\begin{aligned} (4.27) \quad \omega &= \omega_{\mathfrak{m}} + H_1^2 + \sum_{\beta \in \{\alpha, 2\alpha\}} \sum_{j=1}^{m_{\beta}} D_{\beta,j} \\ &= \omega_{\mathfrak{m}} + H_1^2 + \sum_{\beta \in \{\alpha, 2\alpha\}} m_{\beta} H_{\beta} + 2 \sum_{\beta \in \{\alpha, 2\alpha\}} \sum_{j=1}^{m_{\beta}} X_{\beta,j} X_{-\beta,j} \end{aligned}$$

where

$$(4.28) \quad \omega_{\mathfrak{m}} := - \sum_{j=1}^m Z_j^2.$$

Hence

$$P(\omega) = \omega_{\mathfrak{m}} + H_1^2 + \sum_{\beta \in \{\alpha, 2\alpha\}} m_{\beta} H_{\beta} = \omega_{\mathfrak{m}} + \frac{L^2 + 2\rho L}{B(L, L)},$$

from which we conclude

$$(4.29) \quad p_{l,s}(\omega) = p_{l,s}(\omega_{\mathfrak{m}}) + \frac{(s+\rho)^2 - 2\rho(s+\rho)}{B(L, L)} = \frac{1}{d_l} \chi_l(\omega_{\mathfrak{m}}) + \frac{s^2 - \rho^2}{8(n+2)}.$$

To compute $\delta_l(\omega)$ we use Formula (4.27). Observe first that if $f \in \mathcal{D}(G; \chi_l)$ and $Y \in \mathfrak{U}(\mathfrak{k})$, then $f(a_t; Y) = \frac{1}{d_l} \chi_l(Y)$. Hence $\delta_l(Y) = \frac{1}{d_l} \chi_l(Y)$. In particular,

$$(4.30) \quad \delta_l(\omega_{\mathfrak{m}}) = \frac{1}{d_l} \chi_l(\omega_{\mathfrak{m}}).$$

Write

$$(4.31) \quad X_{\beta,j} = Y_{\beta,j} + P_{\beta,j} \quad \text{with} \quad Y_{\beta,j} \in \mathfrak{k}, P_{\beta,j} \in \mathfrak{p}.$$

A standard computation (cf. e.g. [W2], p. 278) then gives for $f \in \mathcal{D}(G; \chi_l)$ and $t > 0$

$$f(a_t; D_{\beta,j}) = \coth(t\beta(L))f(a_t; H_{\beta}) + \frac{4}{d_l} \frac{1 - \cosh(t\beta(L))}{\sinh^2(t\beta(L))} \chi_l(Y_{\beta,j}^2)f(a_t)$$

i.e.

$$(4.32) \quad \delta_l(D_{\beta,j}) = \coth(t\beta(L))H_{\beta} + \frac{4}{d_l} \frac{1 - \cosh(t\beta(L))}{\sinh^2(t\beta(L))} \chi_l(Y_{\beta,j}^2).$$

Notice that $\chi_l(Y_{\alpha,j}^2) = 0$ for all $j = 1, \dots, m_{\alpha}$.

For $h = i, j, k$, let Y_h denote the tangent vector at e to the 1-parameter subgroup $t \mapsto \cos t + h \sin t$ in $\mathrm{Sp}(1)$. Explicit choices of the orthonormal bases in \mathfrak{m} and $\mathfrak{g}_{2\alpha}$ prove that

$$(4.33) \quad \chi_l(\omega_{\mathfrak{m}}) = -2 \sum_{j=1}^3 \chi_l(Y_{2\alpha,j}^2) = -\frac{1}{8(n+2)} \sum_{h \in \{i,j,k\}} \mathrm{tr} [\tau_l(Y_h)^2].$$

As shown in [T1], p. 381, there exists an orthonormal basis $\{v_p\}_{p=-l}^l$ in V_l such that

$$\begin{aligned} \tau_l(Y_i)v_p &= -2ipv_p \\ \tau_l(Y_j)v_p &= -i\alpha_{p+1}^l v_{p+1} \\ \tau_l(Y_k)v_p &= -\alpha_{p+1}^l v_{p+1} + \alpha_p^l v_{p-1} \end{aligned}$$

where

$$\alpha_p^l := [(l+p)(l-p+1)]^{1/2}.$$

It follows that for $h = i, j, k$

$$(4.34) \quad \mathrm{tr} [\tau_l(Y_h)^2] = -\frac{4}{3} l(l+1)(2l+1).$$

Identify A with \mathbf{R} and L with $\frac{d}{dt}$ under the isomorphism $t \mapsto \exp(tL) = a_t$. Formulas (4.27), (4.30) and (4.32)–(4.34) then prove the following proposition.

4.3. PROPOSITION. *Let τ_l be an irreducible unitary representation of K_1 of dimension $2l+1$. Then*

1. *The τ_l -radial component of the Casimir operator ω is*

$$\delta_l(\omega) = \frac{1}{8(n+2)} \left\{ \frac{d^2}{dt^2} + [(4n-1) \coth t + 3 \tanh t] \frac{d}{dt} + \frac{4l(l+1)}{\cosh^2 t} + 4l(l+1) \right\}.$$

2. *For every $s \in \mathbf{C}$*

$$(4.35) \quad p_{l,s}(\omega) = \frac{1}{8(n+2)} [4l(l+1) + s^2 - \rho^2].$$

3. *For every $s \in \mathbf{C}$, the function $\zeta_{l,s}(t) := \zeta_{l,s}(a_t)$ satisfies the differential equation on $(0, +\infty)$*

$$(4.36) \quad u'' + [(4n-1) \coth t + 3 \tanh t] u' + \frac{4l(l+1)}{\cosh^2 t} u = (s^2 - \rho^2) u.$$

The substitution $v(t) = (\cosh t)^{-2l} u(t)$ transforms the differential equation (4.36) into the Jacobi differential equation (cf. [K2], p. 6)

$$(4.37) \quad v'' + [(4n-1) \coth t + (4l+3) \tanh t] v' = (s^2 - \tilde{\rho}^2) v$$

with parameters $\alpha = 2n-1$ and $\beta = 2l+1$. Here $\tilde{\rho} := \alpha + \beta + 1 = \rho + 2l$.

The Jacobi function

$$(4.38) \quad \begin{aligned} \phi_{is}^{(2n-1, 2l+1)}(t) &:= F \left(\frac{\tilde{\rho} + s}{2}, \frac{\tilde{\rho} - s}{2}; 2n; -\sinh^2 t \right) \\ &= F \left(\frac{\rho + s}{2} + l, \frac{\rho - s}{2} + l; 2n; -\sinh^2 t \right) \end{aligned}$$

is the unique solution v to (4.37) satisfying $v(0) = 1$, $v'(0) = 0$. (In (4.38), $F(a, b; c; z)$ denotes the analytic branch on $\mathbf{C} \setminus [1, \infty)$ of the hypergeometric function.)

The τ_l -spherical function $\zeta_{l,s}$ is therefore explicitly given by

$$(4.39) \quad \begin{aligned} \zeta_{l,s}(t) &:= \zeta_{l,s}(a_t) = (\cosh t)^{2l} \phi_{is}^{(2n-1, 2l+1)}(t) \\ &= (\cosh t)^{2l} F \left(\frac{\rho + s}{2} + l, \frac{\rho - s}{2} + l; 2n; -\sinh^2 t \right). \end{aligned}$$

Formula (4.39) has been previously determined by Takahashi ([T2], Formula (7), p. 225) by direct integration of (3.23), using the following expression of χ_l in terms of Gegenbauer polynomials :

$$(4.40) \quad \chi_l(k_1) = C_{2l}^1(\Re k_1) = \frac{\sin((2l+1)\vartheta)}{\sin \vartheta} \quad \text{if } \Re k_1 = \cos \vartheta.$$

Formula (4.35) shows that $p_{l,s}(\omega)$ is an even function of s which assumes arbitrary complex values as s varies in \mathbf{C} . The following corollary can therefore be deduced from Theorem 3.1 and Proposition 4.3.

4.4. COROLLARY. *The τ_l -spherical functions are exactly the functions $\{\zeta_{l,s} : s \in \mathbf{C}\}$ given by Formulas (3.24) and (4.39). Further, $\zeta_{l,s}$ satisfies $\zeta_{l,s}(g) = \zeta_{l,s}(g^{-1})$ for all $g \in G$. Moreover, $\zeta_{l,s} = \zeta_{l,s'}$ if and only if $s = \pm s'$.*

The functional equation (3.15) with $g_1 = a_t$ and $g_2 = a_\tau$ becomes (cf. [T2], Théorème 1, p. 227)

$$(4.41) \quad \zeta_{l,s}(t)\zeta_{l,s}(\tau) = \int_0^\infty K_l(t, \tau, u)\zeta_{l,s}(u)\Delta(u) du$$

where Δ is as in (1.7) and the kernel $K_l(t, \tau, u)$ is defined as follows. Set

$$B := \frac{\cosh^2 t + \cosh^2 \tau + \cosh^2 u - 1}{2 \cosh t \cosh \tau \cosh u}.$$

Then

$$(4.42) \quad K_l(t, \tau, u) := \frac{2^{-2\rho}\Gamma(2n)}{\sqrt{\pi}\Gamma(2n - \frac{1}{2})} \frac{(\cosh t \cosh \tau \cosh u)^{2n-3}}{(\sinh t \sinh \tau \sinh u)^{4n-2}} (1 - B^2)^{2n-\frac{3}{2}} \\ \times F\left(2n + 2l, 2n - 2l - 2; 2n - \frac{1}{2}; \frac{1}{2}(1 - B)\right)$$

if $B < 1$, and $K_l(t, \tau, u) := 0$ if $B \geq 1$. Using (4.39) and Formula (7.11) in [K2], one can prove that (4.41) holds also outside our group-theoretical setting for all $l \in \mathbf{R}$ satisfying $2n - 1 > 2l \geq 0$.

5. THE POSITIVE DEFINITE τ_l -SPHERICAL FUNCTIONS

A continuous function ζ on a locally compact group G is said to be *positive definite* if for every $f \in C_c(G)$

$$\int_G \int_G \zeta(x^{-1}y)f(x)\overline{f(y)} dx dy \geq 0.$$

In this section we establish which among the $\zeta_{l,s}$ are positive definite.

Let us first introduce some notation and recall some definitions. Let G be a semisimple Lie group with finite center, and let K be a maximal compact subgroup of G . \mathfrak{g} and \mathfrak{k} ($\subset \mathfrak{g}$) are the Lie algebras of G and K , respectively. A (strongly continuous) representation T of G on a Banach space \mathcal{H} is denoted by (T, \mathcal{H}) . We may simply speak of the representation T if \mathcal{H} is understood. Irreducibility for T always means topological irreducibility (= no closed proper invariant subspaces). Let \widehat{K} denote the set of equivalence classes