

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** LOCAL-GLOBAL PRINCIPLE FOR NORMS FROM CYCLIC EXTENSIONS OF  $\mathbb{Q}(t)$  (A DIRECT, CONSTRUCTIVE AND QUANTITATIVE APPROACH)  
**Autor:** ZANNIER, Umberto  
**Kapitel:** 4. An example for the non-cyclic case  
**DOI:** <https://doi.org/10.5169/seals-64456>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$$N(r, b_1, \dots, b_d) = f(r)\mu$$

where  $\mu \in \mathbf{Q}_v$  is very close to 1; in fact  $f(r)$  is near to  $f(a_v)$ , which is nonzero. By the previous remarks,  $\mu^{-1}$  is in the image of  $N(r, x_1, \dots, x_d)$  on  $\mathbf{Q}_v^d$ , hence the same must be true for  $f(r)$ , by the basic multiplicative identity for  $N$ . In particular  $f(r)$  will be a norm from  $\mathbf{Q}_v(P_r)$  to  $\mathbf{Q}_v$ .

Let now  $S$  consist of the elements of  $\mathbf{Q}$  which are not poles or zeros of  $f$ , which satisfy  $[\mathbf{Q}(P_s) : \mathbf{Q}] = d$  and which are sufficiently close (in the mentioned sense) to  $a_v$ , for each  $v \in \Sigma$ . We have proved that  $f(s)$  is a norm from  $\mathbf{Q}_v(P_s)$ , for all  $s \in S$  and for all places  $v$ . By Hasse's theorem,  $f(s)$  is a norm from  $\mathbf{Q}(P_s)$ , so  $S \subset N_f$ . On the other hand  $S \cap \mathbf{Z}$  contains the complement of a thin set in an arithmetic progression, whence  $N_f$  cannot satisfy the conclusion of the Theorem (or of Corollary 1), as required.  $\square$

#### 4. AN EXAMPLE FOR THE NON-CYCLIC CASE

We show that assuming that  $L/K$  is cyclic is essential in the Theorem (as in the number-field case, as shown in [CF, Ex. 5]).

To describe a counterexample, define  $L = \mathbf{Q}(t, \sqrt{4t+3}, \sqrt{4t+7})$ ,  $f(t) = t^2$ . We proceed to show that  $\mathbf{N} \subset N_f$ . We have to show that for all large integers  $n$ ,  $n^2$  is a norm from  $L(n) := \mathbf{Q}(\sqrt{4n+3}, \sqrt{4n+7})$ . By [CF, Ex. 5.1 and 5.2, p.360] it is sufficient to show that the local degree  $[L(n)_w : \mathbf{Q}_p]$  is 4 for some prime  $p$ . Observe that the Jacobi symbol  $\left(\frac{4n+3}{4n+7}\right) = \left(\frac{-1}{4n+7}\right) = -1$ . Hence there exists some prime  $p$  dividing  $4n+7$  with an odd multiplicity and such that  $\left(\frac{4n+3}{p}\right) = -1$ . Then  $p$  ramifies in  $L(n)$  and the residual degree is 2, proving the claim. Observe that the first conclusion of Corollary 1 does not hold for  $N_f$ .

On the other hand,  $t^2$  is not a norm from  $L$  to  $K$ . Otherwise by [CF, Ex. 5.1] we could write  $t$  as the product of three norms from the three quadratic subfields of  $L$ . In other words we could write nontrivially

$$q^2(t)t = (a_1^2(t) - (4t+3)b_1^2(t))(a_2^2(t) - (4t+7)b_2^2(t))(a_3^2(t) - (4t+3)(4t+7)b_3^2(t)),$$

where  $q, a_i, b_j \in \mathbf{Q}[t]$ . We may suppose that  $a_i$  and  $b_i$  are coprime for each  $i$ , otherwise we can divide out a common factor. Now, putting  $t = 0$  we get a contradiction.