

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 45 (1999)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON GROUPS ACTING ON NONPOSITIVELY CURVED CUBICAL COMPLEXES
Autor: BALLMANN, Werner / WITKOWSKI, Jacek
Kapitel: 4. NONEXISTENCE OF FREE SUBGROUPS
DOI: <https://doi.org/10.5169/seals-64441>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proof. Let φ be an automorphism of X . If φ fixes a point p of Λ_i^* , then p can be chosen as a vertex or a midpoint of an edge. If p is a vertex, then the preimage X' of p under r_i is a closed and convex subcomplex of X . If p is the midpoint of an edge, X' is a hyperspace and as a union of walls, carries a natural cubical structure. In either case, X' is a closed, convex and φ -invariant subset of X , and therefore φ is semisimple if and only if the restriction $\varphi|_{X'}$ is semisimple. Since moreover X' is a simply connected folded cubical chamber complex of nonpositive curvature and of dimension lower than X , we can assume by induction on $\dim X$ that the action of φ on all the trees Λ_i^* is axial.

Let a_i be an axis of φ in Λ_i^* (unique up to parameter). Let $X_i = r_i^{-1}(a_i)$. Since r_i is surjective, X_i is non-empty. Furthermore, X_i is a closed, convex and φ -invariant subcomplex of X .

Set $Y_1 := X_1$. The image of Y_1 under r_2 is path connected and φ -invariant, hence contains a_2 . Let $Y_2 = Y_1 \cap X_2$. Then Y_2 is non-empty, closed, convex and φ -invariant. By induction we get that $Y = X_1 \cap \dots \cap X_n$ is a non-empty, closed, convex and φ -invariant subcomplex of X . It is then sufficient to prove semisimplicity for the restriction $\varphi|_Y$. Note that $Y = r^{-1}(F)$, where $F \cong \mathbf{R}^n$ is the flat

$$F = \{(a_1(t_1), \dots, a_n(t_n)) \mid t_i \in \mathbf{R}\}$$

in the product of trees. Now φ operates as a translation on F , hence the displacement of φ on F is constant, say $= \delta$. Since r is injective, we can consider Y as a closed subcomplex of F , namely a union of chambers. The metric on Y is the induced path metric. It follows easily that there are only finitely many possible values for the distance in Y from a point x to its image φx , if the location of x in its chamber is given. \square

4. NONEXISTENCE OF FREE SUBGROUPS

In this section we discuss the proof of Theorem 2 of the introduction. We assume throughout this section that X is a simply connected folded cubical chamber complex of nonpositive curvature and that $\Gamma \subset \text{Aut}(X)$ is a group that preserves the folding of X (this can be always assumed by passing to a finite index normal subgroup if necessary) and does not contain a free nonabelian subgroup acting freely on X . By equivariance of the maps r_i , the same holds for the actions of Γ on the trees Λ_i^* . Up to a subgroup of index two, there are three possibilities for each particular i [PV]:

- (0) Γ fixes a point of Λ_i^* ;
- (1) Γ fixes no point of Λ_i^* , but precisely one end of Λ_i^* ;
- (2) Γ fixes no point of Λ_i^* , but precisely two ends of Λ_i^* .

Thus by passing to a subgroup of Γ of index at most 2^n , we can assume that the above three alternatives hold for all i . Corresponding to the alternative, we say that i is an *index of type 0, 1 or 2* respectively.

We first construct a homomorphism $h = (h_1, \dots, h_n): \Gamma \rightarrow \mathbf{Z}^n$ as claimed. If Γ fixes a point of Λ_i^* , we define h_i to be the trivial homomorphism. If Γ does not fix a point of Λ_i^* , we let ω_i be the end or one of the two ends of Λ_i^* fixed by Γ . The Busemann function $b_i: \Lambda_i^* \rightarrow \mathbf{R}$ at ω_i is well defined up to an additive constant (see [Ba], Section 1 of Chapter II). Since Γ fixes ω_i ,

$$h_i(\phi) := b_i(\phi p) - b_i(p), \quad p \in \Lambda_i^*,$$

is a well defined homomorphism $h_i: \Gamma \rightarrow \mathbf{Z}$, called the *Busemann homomorphism*. Note that h_i is integer valued since Λ_i^* is a simplicial tree and Γ acts by automorphisms. This completes the definition of $h = (h_1, \dots, h_n)$. We set

$$\Delta_i = \ker h_i \quad \text{and} \quad \Delta = \bigcap \Delta_i = \ker h.$$

PROPOSITION 4.1. Δ consists precisely of the elliptic elements of Γ .

Proof. If the action of Γ on Λ_i^* has a fixed point, then any $\phi \in \Gamma$ is elliptic on Λ_i^* and $\Delta_i = \Gamma$. If Γ does not have a fixed point in Λ_i^* , but fixes a point $\xi_i \in \Lambda_i^*(\infty)$ and $\phi \in \Gamma$ is axial on Λ_i^* , then ξ_i is an end point of the axis of ϕ . Then $h_i(\phi) \neq 0$. Hence by Proposition 3.5, any $\phi \in \Delta$ is elliptic on X . Conversely, if $\phi \in \Gamma$ is elliptic on X , then $\phi \in \Delta$. \square

For the proof of the other assertions of Theorem 2 we need some more preparations.

LEMMA 4.2. Let Λ be a simplicial tree on which Γ acts by automorphisms. Suppose Δ fixes a point of Λ . Then either Γ fixes a point of Λ or exactly two points in $\Lambda(\infty)$.

Proof. Since Δ is a normal subgroup of Γ , the set Φ of fixed points of Δ is Γ -invariant. Now Φ is a subtree of Λ , hence we can assume $\Phi = \Lambda$. Then the quotient action by Γ/Δ on Λ is well defined.

Suppose that Γ/Δ contains an element ϕ which is axial on Λ . Since Γ/Δ is abelian, it leaves the unique axis of ϕ invariant and fixes the endpoints of the axis.

Suppose now that all elements of Γ/Δ are elliptic on Λ . Let ϕ_1, \dots, ϕ_k be a system of generators. The set of fixed points of ϕ_1 is a Γ/Δ -invariant subtree. Replacing Λ by this subtree, we can assume that $\phi_1 = \text{id}_\Lambda$. The quotient of Γ/Δ by the subgroup generated by ϕ_1 is abelian and has a system of $k-1$ generators. Induction on k shows that Γ has a fixed point. \square

If i is an index of type 0 and $p \in \Lambda_i^*$ a fixed point, then $X' := r_i^{-1}(p) \subset X$ is closed, convex and Γ -invariant. In particular, $X'(\infty) \subset X(\infty)$ is Γ -invariant. Although X' is not a subcomplex if p is not a vertex, it is parallel to the walls with label i in the chambers it intersects. Hence we obtain a natural cubical structure on X' with a folding onto an $(n-1)$ -cube, and Γ preserves this cubical structure and folding. Hence by passing to such subspaces if necessary, we can assume that no indices of type 0 occur.

Let i be an index of type 2. Let $\alpha_i, \omega_i \in \Lambda_i^*(\infty)$ be the fixed points of Γ and σ_i the unit speed geodesic from α_i to ω_i . Then σ_i is Γ -invariant and $\Delta_i = \text{Stab}(\sigma_i(t))$ for all $t \in \mathbf{R}$. Hence $X' = r_i^{-1}(\text{im } \sigma_i)$ is a closed, convex and Γ -invariant subcomplex of X . Hence by passing to such subspaces if necessary, we can assume that $\Lambda_i^* = \text{im } \sigma_i \cong \mathbf{R}$ for all indices i of type 2.

PROPOSITION 4.3. *If there are no indices of type 1, then there is a Γ -invariant convex subset $E \subset X$ isometric to a Euclidean space of dimension $k \in \{0, \dots, n\}$ and an exact sequence*

$$0 \rightarrow \Delta \rightarrow \Gamma \rightarrow \mathbf{Z}^k \rightarrow 0$$

such that Δ fixes E pointwise and such that the quotient $\Gamma/\Delta \cong \mathbf{Z}^k$ acts on E as a cocompact lattice of translations.

Proof. After reductions as above we can assume that all indices are of type 2, that $\Lambda_i^* \cong \mathbf{R}$ for all i and that Δ fixes each point of $\prod \Lambda_i^*$. Since r is an injection, Δ fixes each point of X .

The image $\text{im } h$ of the homomorphism h is a subgroup of the group \mathbf{Z}^n , hence it is isomorphic to \mathbf{Z}^k for some $k \leq n$. Thus we may identify the quotient group Γ/Δ with \mathbf{Z}^k . Consider the quotient action of $\mathbf{Z}^k = \Gamma/\Delta$ on X , which is well defined since Δ acts trivially on X . This action is free and the elements are semisimple by Proposition 3.6. Applying the Flat Torus Theorem, see [CE] and [BH], we get that there exists a \mathbf{Z}^k -invariant convex subspace $E \subset X$, isometric to k -dimensional Euclidean space, such that \mathbf{Z}^k acts on it as a cocompact lattice of translations. \square

We now discuss the more difficult case that indices of type 1 occur. As explained above, we can assume that no indices of type 0 occur and that $\Lambda_i^* \cong \mathbf{R}$ for all indices of type 2.

Choose a vertex $x_0 \in X$ as an origin. For indices of type 2 choose the parameter on the above geodesics σ_i such that $\sigma_i(0) = r_i(x_0)$. For indices of type 1 we denote by $\omega_i \in \Lambda_i^*(\infty)$ the corresponding fixed point. For these indices, we let $\sigma_i: [0, \infty) \rightarrow \Lambda_i^*$ be a unit speed geodesic ray with $\sigma_i(0) = r_i(x_0)$ and $\sigma_i(\infty) = \omega_i$.

We set $F = \text{im } \sigma_1 \times \cdots \times \text{im } \sigma_n$. Note that F is a closed and convex subspace of $\prod \Lambda_i^*$. We also define a geodesic ray

$$\sigma: [0, \infty) \rightarrow F \quad \text{by} \quad \sigma(t) = (\sigma_1(t), \dots, \sigma_n(t)).$$

By construction, $\sigma(0) = r(x_0)$.

LEMMA 4.4. *$\text{Stab}(\sigma_i(t)) \rightarrow \Delta_i$ and $\text{Stab}(\sigma(t)) \rightarrow \Delta$ as $t \rightarrow \infty$, where the limit of groups is understood as the union of increasing family.*

Proof. Let $\phi \in \Delta_i$. Then ϕ fixes $\omega_i = \sigma_i(\infty)$. Therefore $\phi \circ \sigma_i$ is asymptotic to σ_i . Now Λ_i^* is a tree, hence $\phi \circ \sigma_i(t) = \sigma_i(t + c)$ for all t sufficiently large, where c is some constant independent of t . Since $\phi \in \Delta_i$, $c = 0$ and therefore $\phi \in \text{Stab}(\sigma_i(t))$ for all t sufficiently large. \square

COROLLARY 4.5. *There exists a sequence (x_m) in X such that $\text{Stab}(x_m) \rightarrow \Delta$.*

Proof. We observe that $\text{Stab}(x) \subset \Delta$ for all $x \in X$. Now the assertion follows immediately from Proposition 3.5 and Lemma 4.4. \square

LEMMA 4.6. *If the group Γ fixes precisely one point $\omega_i \in \Lambda_i^*(\infty)$, then $\Delta \cap \text{Stab}(\sigma_i(t))$ has infinitely many jumps as $t \rightarrow \infty$.*

Proof. Let $\phi \in \Delta \subset \Delta_i$. By Lemma 4.4 there is $t_\phi \geq 0$ such that $\phi \in \text{Stab}(\sigma_i(t))$ for all $t \geq t_\phi$. Hence if $\Delta \cap \text{Stab}(\sigma_i(t)) = \Delta \cap \text{Stab}(\sigma_i(t'))$ for all t, t' sufficiently large, then $\Delta \subset \text{Stab}(\sigma_i(t))$ for all t sufficiently large. By Lemma 4.2, Γ either fixes a point of Λ_i^* , which is excluded by our reductions above, or Γ fixes exactly two points of $\Lambda_i^*(\infty)$, which is in contradiction to the assumption. \square

LEMMA 4.7. *Let (x_m) be a sequence in X such that $\text{Stab}(x_m) \rightarrow \Delta$ and $\gamma_m: [0, s_m] \rightarrow X$ be the unit speed geodesic from x_0 to x_m , where $s_m = d(x_0, x_m)$. Then given a constant $t_0 > 0$, there exists m_0 such that $s_m \geq t_0$ and $r \circ \gamma_m([0, t_0]) \in F$ for all $m \geq m_0$.*

Proof. For those i for which Γ fixes exactly one point $\omega_i \in \Lambda_i^*(\infty)$ we choose $\phi_i \in \Delta$ such that $\phi_i \notin \text{Stab}(\sigma_i(t))$ for $t \leq t_0$, see Lemma 4.6. By assumption, there is m_0 such that $\phi_i \in \text{Stab}(x_m)$ for all $m \geq m_0$ and all such i . Now $r_i \circ \gamma_m$ is a monotonic curve in Λ_i^* from $\sigma_i(0) = r_i(x_0)$ to $r_i(x_m)$. By equivariance of r_i , $\phi_i \in \text{Stab}(r_i(x_m))$ for all $m \geq m_0$. On the other hand, $r_i \circ \sigma$ has speed ≤ 1 , hence by the choice of t_0 , $s_m \geq t_0$ and $r_i(\gamma_m(t)) \in \sigma_i([0, t_0])$ for $0 \leq t \leq t_0$.

The claim follows since the image of r_i is σ_i for those i for which Γ fixes exactly two ends of Λ_i^* . \square

LEMMA 4.8. *Given $\phi \in \Gamma$, there is a constant $c = c_\phi$ such that $d(\phi(p), p) \leq c$ for all $p \in F$.*

Proof. We show that $d_i(\phi(p), p) \leq c_i$ for each point p in the image of σ_i . This is clear for those indices i for which Γ fixes exactly two ends of Λ_i^* . Consider some other index i . Then σ_i is defined on $[0, \infty)$.

If ϕ is elliptic on Λ_i^* , then $\phi \in \Delta_i$. By Lemma 4.4, there exists a constant t_ϕ such that ϕ fixes $\sigma_i(t)$ for all $t \geq t_\phi$. We conclude that $d_i(\phi(p), p) \leq 2t_\phi$ for each point p in the image of σ_i .

We assume now that ϕ is axial on Λ_i^* and let ρ be an axis of ϕ in Λ_i^* . We parametrize ρ such that $\rho(\infty) = \omega_i$. Since Λ_i^* is a tree and $\sigma_i(\infty) = \rho(\infty)$, we can actually choose the parameter such that $\sigma_i(t) = \rho(t)$ for all $t \geq t_\phi$, where t_ϕ is an appropriate constant. Now $\phi(\rho(t)) = \rho(t + \tau)$ for some constant τ independent of t . We conclude that $d_i(\phi(p), p) \leq 2t_\phi + \tau$ for each point p in the image of σ_i . \square

PROPOSITION 4.9. *Suppose that indices of type 1 occur. Then*

- (1) Δ does not fix a point of X ;
- (2) Γ fixes a point in $X(\infty)$. More precisely, if (x_m) is a sequence in X such that $\text{Stab}(x_m) \rightarrow \Delta$, then after passing to a subsequence if necessary, (x_m) converges to a fixed point $\xi \in X(\infty)$ of Γ .

Proof. The first assertion is an immediate consequence of Lemma 4.7. As for the proof of the second assertion, let (x_m) be a sequence in X with $\text{Stab}(x_m) \rightarrow \Delta$. Let $\gamma_m: [0, s_m] \rightarrow X$ be the unit speed geodesic from x_0 to x_m as in Lemma 4.7. Note that $r \circ \gamma_m$ is a sequence of unit speed curves (with respect to the metric $d_{(2)}$, for which r restricted to any chamber of X is an isometry) in $\prod \Lambda_i^*$. For each constant $t_0 > 0$, $r \circ \gamma_m([0, t_0])$ is contained in F for all m sufficiently large. Now F is locally compact, hence a subsequence of

the sequence of curves $r \circ \gamma_m$ converges locally uniformly. By Proposition 3.4, the corresponding subsequence of the sequence of unit speed geodesics γ_m converges locally uniformly. By definition, this means that the corresponding subsequence of (x_m) converges to a point $\xi \in X(\infty)$.

Let $\phi \in \Gamma$ and choose $c = c_\phi$ as in Lemma 4.8. Let $t_0 > 0$ be given. By Lemma 4.8 we have $r \circ \gamma_m(t_0) \in F$ for all $m \geq m_0$. By Proposition 3.4 and Lemma 4.8, we have $d(\phi(\gamma_m(t_0)), \gamma_m(t_0)) \leq \sqrt{n}c_\phi$ for all such m . Now c_ϕ is independent of t_0 , hence $\phi(\xi) = \xi$. \square

We now complete the proof of Theorem 2 of the introduction. By Proposition 4.1, $\Delta \cong \ker h$ consists precisely of the elliptic elements of Γ . If indices of type 1 do not occur, then Proposition 4.3 applies: If $k = 0$, then $\Gamma \cong \Delta$ fixes a point of X and possibility (1) holds. If $k > 0$, then possibility (2) holds. If indices of type 1 occur, then possibility (3) holds by Proposition 4.9 and Corollary 4.5. Note that $\text{Stab}(x) \neq \Delta$ for any $x \in X$ in this case since Δ would have a fixed point otherwise.

5. PARALLEL TRANSPORT IN A CUBICAL MANIFOLD AND THE PROOF OF THEOREM 3

Let X be a cubical manifold of dimension n . Given two chambers P and Q in X with a common face of dimension $n - 1$, we define $t_{PQ}: P \rightarrow Q$ to be the *translation* which moves each point p of P along the unit geodesic segment starting at p and orthogonal to the common $(n - 1)$ -face of P to the end point in Q . The map t_{PQ} is an isomorphism and isometry of P with Q . Given a gallery $\pi = (P_1, \dots, P_n)$ in X , the *parallel transport* along π is the isomorphism $t_\pi: P_1 \rightarrow P_n$ given by

$$t_\pi := t_{P_{n-1}P_n} \circ \dots \circ t_{P_2P_3} \circ t_{P_1P_2}.$$

LEMMA 5.1. *Let X be a simply connected cubical manifold and assume that the number of chambers adjacent to each face of codimension 2 in X is divisible by 4. Then for any two chambers P and Q in X , the parallel transport t_π along a gallery π connecting P and Q is independent of π .*

Proof. It is enough to show that the parallel transport along any closed gallery is the identity. Let π be such a gallery with initial and final chamber P .