**Zeitschrift:** L'Enseignement Mathématique
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique
**Band:** 45 (1999)
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** COUNTING PATHS IN GRAPHS
**Autor:** Bartholdi, Laurent
**Kapitel:** 7.2 Cycles
**DOI:** https://doi.org/10.5169/seals-64442

**Nutzungsbedingungen**
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

**Conditions d'utilisation**
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

**Terms of use**
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 15.01.2026

**ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch**

Solving the system, we obtain

$$F(u,t) = \frac{1+(1-u)t}{1-(v-2+u)t} \cdot \frac{1-(v-2)t+(1-u)(v-2+u)t^2}{1+t+(1-u)(v-2+u)t^2}.$$

We then compute

$$G(t) = F(1,t) = \frac{1-(v-2)t}{(1+t)(1-(v-1)t)},$$

$$F(0,t) = \frac{(1+t)(1-(v-2)t+(v-2)t^2)}{(1-(v-2)t)(1+t+(v-2)t^2)}.$$

## 7.2 Cycles

Let $\mathcal{X} = C_k$, the cycle on $k$ vertices. Here, as there are 2 proper circuits of length $n$ for all $n$ multiple of $k$ (except 0), we have

$$F(0,t) = \frac{1+t^k}{1-t^k}.$$

Obtaining a closed form for $G$ is much harder. The number of circuits of length $n$ is

$$g_n = \sum_{i\in\mathbf{Z}:\, i\equiv 0\ [k],\ i\equiv n\ [2]} \binom{n}{\frac{n+i}{2}},$$

from which, by [Gou72, 1.54], it follows that

$$G(t) = \frac{1}{k}\sum_{\zeta^k=1} \frac{1}{1-\left(\zeta+\zeta^{-1}\right)t} = \frac{1}{k}\sum_{j=0}^{k-1} \frac{1}{1-2\cos\left(\frac{2\pi j}{k}\right)t}.$$

It is not at all obvious how to simplify the above expression. A closed-form answer can be obtained from (2.3), namely

$$G(t) = \frac{(2t)^2+\left(1-\sqrt{1-4t^2}\right)^2}{(2t)^2-\left(1-\sqrt{1-4t^2}\right)^2} \cdot \frac{(2t)^k+\left(1-\sqrt{1-4t^2}\right)^k}{(2t)^k-\left(1-\sqrt{1-4t^2}\right)^k},$$

or, expanding,

$$G(t) = \frac{(2t)^k + \sum_{m=0}^{k/2}\left(1-4t^2\right)^m \binom{k}{2m}}{\sum_{m=1}^{(k+1)/2}\left(1-4t^2\right)^m \binom{k}{2m-1}}.$$

However in general this fraction is not reduced. To obtain reduced fractions for $F(u,t)$ (and thus for $G(t)$), we have to consider separately the cases where $k$ is odd or even.

For odd $k$, letting $k = 2\ell + 1$, we obtain

$$F(u,t) = \frac{1+(1-u)t}{1-(1+u)t} \cdot \frac{\sum_{m=0}^{\ell} \alpha_m^\ell (-t)^m (1+(1-u^2)t^2)^{\ell-m}}{\sum_{m=0}^{\ell} \alpha_m^\ell t^m (1+(1-u^2)t^2)^{\ell-m}},$$

$$G(t) = \frac{\sum_{m=0}^{\ell} \alpha_m^\ell (-t)^m}{(1-2t)\left(\sum_{m=0}^{\ell} \alpha_m^\ell t^m\right)},$$

where

$$\alpha_m^\ell = \begin{cases} (-)^{\frac{m}{2}} \binom{\ell - \frac{m}{2}}{\frac{m}{2}} & \text{if } m \equiv 0 \ [2], \\ (-)^{\frac{m-1}{2}} \binom{\ell - \frac{m+1}{2}}{\frac{m-1}{2}} & \text{if } m \equiv 1 \ [2]. \end{cases}$$

For even $k$, with $k = 2\ell$,

$$F(u,t) = \frac{\sum_{m=0}^{\ell/2} \frac{\ell}{\ell-m} \binom{\ell-m}{m} (-t^2)^m (1-(1-u^2)t^2)^{\ell-2m}}{(1-(1+u)^2t^2)\left(\sum_{m=0}^{(\ell-1)/2} \binom{\ell-1-m}{m} (-t^2)^m (1-(1-u^2)t^2)^{\ell-1-2m}\right)},$$

$$G(t) = \frac{\sum_{m=0}^{\ell/2} \frac{\ell}{\ell-m} \binom{\ell-m}{m} (-t^2)^m}{(1-4t^2)\left(\sum_{m=0}^{(\ell-1)/2} \binom{\ell-1-m}{m} (-t^2)^m\right)},$$

expressed as reduced fractions.

The first few values of $F$, where $\square$ stands for $1+(1-u^2)t^2$, are:

| $k$ | $F(u,t)$ | $k$ | $F(u,t)$ |
|---|---|---|---|
| 1 | $\dfrac{1+(1-u)t}{1-(1+u)t}$ | 2 | $\dfrac{\square}{1-(1+u)^2t^2}$ |
| 3 | $\dfrac{(1+(1-u)t)(\square-t)}{(1-(1+u)t)(\square+t)}$ | 4 | $\dfrac{\square^2-2t^2}{1-(1+u)^2t^2}$ |
| 5 | $\dfrac{(1+(1-u)t)(\square^2-\square t-t^2)}{(1-(1+u)t)(\square^2+\square t+t^2)}$ | 6 | $\dfrac{\square^2-3t^2}{(1-(1+u)^2t^2)(\square^2-t^2)}$ |
| 7 | $\dfrac{(1+(1-u)t)(\square^3-\square^2t-2\square t^2+t^3)}{(1-(1+u)t)(\square^3+\square^2t-2\square t^2-t^3)}$ | 8 | $\dfrac{\square^4-4\square^2t^2+2t^4}{(1-(1+u)^2t^2)(\square^2-2t^2)}$ |
| 9 | $\dfrac{(1+(1-u)t)(\square-t)(\square^3-3\square t^2-t^3)}{(1-(1+u)t)(\square+t)(\square^3-3\square t^2+t^3)}$ | 10 | $\dfrac{\square^4-5\square^2t^2+5t^4}{(1-(1+u)^2t^2)(\square^4-3\square^2t^2+t^4)}$ |

These rational expressions were computed and simplified using the computer algebra program *Maple*™.

## 7.3 Trees

Let $\mathcal{X}$ be the $d$-regular tree. Then

$$F(0,t)=1$$

as a tree has no proper circuit; while direct (i.e., without using Corollary 2.6) computation of $G$ is more complicated. It was first performed by Kesten [Kes59]; here we will derive the extended circuit series $F(u,t)$ and also obtain the answer using Corollary 2.6.

Let $\mathcal{T}$ be a regular tree of degree $d$ with a fixed root $\star$, and let $\mathcal{T}'$ be the connected component of $\star$ in the two-tree forest obtained by deleting in $\mathcal{T}$ an edge at $\star$. Let $F(u,t)$ and $F'(u,t)$ respectively count circuits at $\star$ in $\mathcal{T}$ and $\mathcal{T}'$. For instance if $d=2$ then $F'$ counts circuits in $\mathbf{N}$ and $F$ counts circuits in $\mathbf{Z}$. For a reason that will become clear below, we make the convention that the empty circuit is counted as '1' in $F$ and as '$u$' in $F'$. Then we have

$$F' = u + (d-1)\,tF't\frac{1}{1-(d-2+u)\,tF't},$$

$$F = 1 + dtF't\frac{1}{1-(d-1+u)\,tF't}.$$

Indeed a circuit in $\mathcal{T}'$ is either the empty circuit (counted as $u$), or a sequence of circuits composed of, first, a step in any of $d-1$ directions, then