

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TEICHMÜLLER SPACE AND FUNDAMENTAL DOMAINS OF FUCHSIAN GROUPS  
**Autor:** SCHMUTZ SCHALLER, Paul  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-64444>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 18.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## TEICHMÜLLER SPACE AND FUNDAMENTAL DOMAINS OF FUCHSIAN GROUPS

by Paul SCHMUTZ SCHALLER

### 1. INTRODUCTION

There are a number of ways to define the Teichmüller space of Riemann surfaces. In this paper I treat an approach which is less common than others. Let  $\Gamma$  be a Fuchsian group which uniformizes a closed Riemann surface of genus  $g$ . Then a fundamental domain for  $\Gamma$  is chosen in a canonical way, namely as a polygon with  $4g$  sides such that opposite sides are identified. The Teichmüller space  $T_g$  of closed Riemann surfaces of genus  $g$  is then constructed by varying these polygons.

This construction of  $T_g$  by polygons was first done by Coldewey and Zieschang in an annex in [17], see also [18]; the construction includes the proof that  $T_g$  is homeomorphic to  $\mathbf{R}^{6g-6}$ . In [2], Buser gave a different, however indirect proof. Here, I propose a new construction and a new proof which is, in my eyes, easier and more transparent than the original one of Coldewey and Zieschang.

The main idea is the following. Let  $P(g)$  be a canonical polygon of  $4g$  sides which is the fundamental domain of a Fuchsian group uniformizing a closed Riemann surfaces of genus  $g$  (the definition of  $P(g)$  will include some technical subtleties, to be discussed in Section 3). Then “triangulate”  $P(g)$  into  $4g - 4$  triangles and one quadrilateral  $S$ . This can be done in such a way that these triangles are determined by  $6g - 5$  positive real numbers (corresponding to the lengths of the sides of the triangles) with the condition that the different triangle inequalities hold. It turns out that these  $6g - 5$  lengths, *taken as homogeneous parameters*, provide a parametrization of the Teichmüller space  $T_g$ . Since the set of reals for which the different triangle




FIGURE 1

On the left hand side: usual identification

On the right hand side: identification chosen in this paper

inequalities hold is open and convex, this also proves that  $T_g$  is homeomorphic to  $\mathbf{R}^{6g-6}$ .

Let  $P$  be a polygon of  $4g$  sides which is the fundamental domain for a Fuchsian group  $\Gamma$  uniformizing a closed Riemann surface  $M$  of genus  $g$ . This means that we can write

$$M = \mathbf{H}/\Gamma$$

where  $\mathbf{H}$  is the upper halfplane. Usually,  $P$  is chosen such that the identification of the sides of  $P$  is that of the polygon on the left hand side in Figure 1. The construction described above would equally work for these polygons. For the following reasons I prefer to choose the identification (compare the polygon on the right hand side of Figure 1) such that opposite sides are identified. First the sides of  $P$  correspond to simple (this means with no selfintersections) closed curves in  $M$  and if opposite sides are identified, then these simple closed curves intersect transversally (which is not the case with the usual identification). Secondly, the vertices of  $P$  correspond to a (unique) point  $Q$  in  $M$ ; with the usual identification,  $Q$  is completely arbitrary while with the identification chosen here, there is a natural choice for  $Q$  in the case of hyperelliptic Riemann surfaces, namely, as a Weierstrass point. See Section 6 for details.

In this paper, I only treat the case of Fuchsian groups which uniformize closed Riemann surfaces. In a straightforward way, the construction and proof could be extended to all finitely generated Fuchsian groups. Note that concerning the original construction and proof in [17] (mentioned above) the corresponding generalization has been worked out by Coldewey in his thesis [3].

The paper is structured as follows. In Section 2 the basic definitions of hyperbolic geometry and Fuchsian groups are given. Section 3 defines the

canonical polygons. Section 4 provides the necessary material from hyperbolic trigonometry, it contains also some lemmas needed later. Section 5 contains the proof of the main theorem and Section 6 gives some applications, mainly concerning hyperelliptic Riemann surfaces. More precisely, I give a new proof of a geometric characterization of hyperelliptic Riemann surfaces which first appeared in [14] (I thank very much Feng Luo who, by his comments on [14], has contributed to the idea of this new proof). I also show (and this is a new result) that the Teichmüller space  $T_g$  for  $g = 2$  can be parametrized by 7 geodesic length functions, taken as homogeneous parameters. This is the optimum parametrization of Teichmüller space by geodesic length functions which one can expect.

I spoke about the content of this paper in lectures of the Troisième Cycle Romand de Mathématiques (Lausanne 1997); I thank the participants for their comments.

## 2. HYPERBOLIC GEOMETRY AND FUCHSIAN GROUPS

The material of this section and of parts of the following section is standard, see for example [1], [4], [5], [6], [7], [8], [15].

DEFINITION. (i)  $\mathbf{H} = \{z = (x, y) \in \mathbf{C} : y > 0\}$  denotes the *upper halfplane*. The *hyperbolic metric* on  $\mathbf{H}$  is given by

$$dz = \frac{1}{y}(dz)_E$$

where  $(dz)_E$  is the standard Euclidean metric on  $\mathbf{C}$  and  $y$  is the imaginary part of  $z$ .

(ii) Define

$$\mathrm{SL}(2, \mathbf{R}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : ad - bc = 1; a, b, c, d \in \mathbf{R} \right\}$$

and

$$\mathrm{PSL}(2, \mathbf{R}) = \mathrm{SL}(2, \mathbf{R})/\sim$$

with  $A \sim B$  if and only if  $A = \pm B$  for  $A, B \in \mathrm{SL}(2, \mathbf{R})$ . Let  $\gamma \in \mathrm{SL}(2, \mathbf{R})$ ,

$$\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$