

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TEICHMÜLLER SPACE AND FUNDAMENTAL DOMAINS OF FUCHSIAN GROUPS  
**Autor:** SCHMUTZ SCHALLER, Paul  
**Kapitel:** 3. FUNDAMENTAL DOMAINS AND CANONICAL POLYGONS  
**DOI:** <https://doi.org/10.5169/seals-64444>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**THEOREM 4.** *Let  $\Gamma$  be a Fuchsian group without elliptic elements (an element  $\gamma \in \text{PSL}(2, \mathbf{R})$  is elliptic if  $|\text{tr}(\gamma)| < 2$  where  $\text{tr}$  is the trace). Then  $\mathbf{H}/\Gamma$  is a complete connected orientable Riemannian manifold of dimension 2 with a metric of constant curvature  $-1$ .*

**DEFINITION.** A *hyperbolic surface* is a connected orientable manifold  $M = \mathbf{H}/\Gamma$  as in Theorem 4 (where  $\Gamma$  is a Fuchsian group without elliptic elements).  $M$  is called *closed* if  $M$  is compact and has no boundary.

### 3. FUNDAMENTAL DOMAINS AND CANONICAL POLYGONS

**DEFINITION** (Compare Figure 2). Let  $g \geq 2$  be an integer. A *canonical polygon*  $P(g)$  is a polygon with  $4g$  sides, denoted by  $a_1, \dots, a_{4g}$ , ordered clockwise, and angles  $\alpha_i$  between  $a_i$  and  $a_{i+1}$ ,  $i = 1, \dots, 4g$  (indices are taken modulo  $4g$ ), such that

- (I)  $a_i$  and  $a_{i+2g}$  have the same length,  $i = 1, \dots, 2g$ ;
- (II) the sum of the angles of  $P(g)$  is  $2\pi$ ;
- (III)  $0 < \alpha_i < \pi$ ,  $i = 1, \dots, 4g$ ;
- (IV)  $\alpha_1 = \alpha_{2g+1}$ ;

$$(V) \sum_{i=1}^g \alpha_{2i-1} + \sum_{i=g+1}^{2g} \alpha_{2i} = \sum_{i=1}^g \alpha_{2i} + \sum_{i=g+1}^{2g} \alpha_{2i-1}.$$

I shall speak of condition (I) (or (II) or (III) or (IV) or (V)) referring to this definition.




FIGURE 2  
A canonical polygon  $P(g)$  for  $g = 2$

REMARKS. (i) Note that, by condition (II), both sides of the equation in condition (V) equal  $\pi$ .

(ii) The terminology *canonical* polygon is not standard, one finds different objects called canonical polygons in the literature (see for example in [15]).

DEFINITION. Let  $\Gamma$  be a Fuchsian group. A *fundamental domain* for  $\Gamma$  is a measurable subset  $D$  of  $\mathbf{H}$  such that

- (i)  $\bigcup_{\gamma \in \Gamma} \gamma(D) = \mathbf{H}$ , and
- (ii)  $\text{int}(\bar{D}) \cap \text{int}(\gamma(\bar{D})) = \emptyset$  for  $\text{id} \neq \gamma \in \Gamma$ . Here,  $\text{int}(S)$  is the *interior* of a set  $S$  and  $\text{id}$  is the unit matrix.

THEOREM 5 (Poincaré). *A canonical polygon  $P = P(g)$  is the fundamental domain of a Fuchsian group  $\Gamma$  and  $\mathbf{H}/\Gamma$  is a closed hyperbolic surface of genus  $g$ . The group  $\Gamma$  is generated by the  $2g$  elements  $\gamma_i$  where  $\gamma_i$  is defined by the conditions  $\gamma_i(P) \cap \text{int}(P) = \emptyset$  and  $\gamma_i(a_i) = a_{i+2g}$  if  $i$  is odd and  $\gamma_i(a_{i+2g}) = a_i$  if  $i$  is even,  $i = 1, \dots, 2g$ .*

REMARKS. (i) For a proof see for example Poincaré [10], Siegel [15], Beardon [1], Iversen [5]. The theorem holds for much more general polygons. A general proof was first given by Maskit [9] and by de Rham [11].

(ii) Traditionally, the  $2g$  generators  $\gamma_i$  of a Fuchsian group corresponding to a closed hyperbolic surface of genus  $g$  are chosen such that the relation

$$\prod_{i=1}^{2g} [\gamma_{2i-1}, \gamma_{2i}] = \text{id}$$

holds where

$$[\gamma_{2i-1}, \gamma_{2i}] = \gamma_{2i-1} \gamma_{2i} (\gamma_{2i-1})^{-1} (\gamma_{2i})^{-1}.$$

With the choice made here, the relation

$$\gamma_1 \gamma_2 \dots \gamma_{2g} (\gamma_1)^{-1} (\gamma_2)^{-1} \dots (\gamma_{2g})^{-1} = \text{id}$$

holds. Compare the introduction for the reasons for this choice.

(iii) Let  $P(g)$  be a canonical polygon and  $M = \mathbf{H}/\Gamma$  be the corresponding closed hyperbolic surface. Then the vertices of  $P(g)$  correspond to a unique point  $Q$  in  $M$  and the side  $a_i$  (as well as  $a_{2g+i}$ ) of  $P(g)$  corresponds to a simple closed curve  $u_i$  in  $M$ ,  $i = 1, \dots, 2g$ . These curves all intersect transversally in  $Q$  and intersect in no other point. Moreover, these curves are geodesic loops based in  $Q$ , this means that the curves may have an angle  $\neq \pi$  in  $Q$ , but outside  $Q$ , they are geodesic. Further, condition (IV) and

condition (V) of canonical polygons are equivalent to the condition that  $u_1$  and  $u_2$  are simple closed geodesics in  $M$ .

#### 4. TRIGONOMETRY

REMARK. By abuse of notation a side of a polygon will often be identified with its length.

The following theorem is standard (for a proof see for example [1], [2]).




FIGURE 3  
The notation for a triangle

**THEOREM 6.** *Let  $T$  be a triangle with angles  $\alpha, \beta, \gamma$  and sides of length  $a, b, c$  with the notation of Figure 3. Then*

- (i)  $\frac{\sinh a}{\sin \alpha} = \frac{\sinh b}{\sin \beta} = \frac{\sinh c}{\sin \gamma}$ ;
- (ii)  $\cosh c = \cosh a \cosh b - \sinh a \sinh b \cos \gamma$ ;
- (iii)  $\cos \gamma = -\cos \alpha \cos \beta + \sin \alpha \sin \beta \cosh c$ .

**LEMMA 7.** *Let  $T$  be a triangle with the notation of Figure 3. Let  $T'$  be a triangle with sides of length  $a', b', c'$  and angles  $\alpha', \beta', \gamma'$ . Let  $a = a'$  and  $b = b'$ . Then*

$$c' > c \iff \gamma' > \gamma \iff \alpha' + \beta' < \alpha + \beta.$$

*Proof.* The first equivalence is a consequence of Theorem 6(ii).

Let  $Z$  be the centre of the side  $c$  and let  $u$  be the geodesic segment, of length  $d/2$  say, between  $Z$  and the vertex  $C$  of  $T$ . The segment  $u$  separates  $T$  into two triangles (compare Figure 4). Applying Theorem 6(ii) to them, we obtain

$$\cosh a = \cosh(c/2) \cosh(d/2) - \sinh(c/2) \sinh(d/2) \cos \delta$$