

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 45 (1999)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TEICHMÜLLER SPACE AND FUNDAMENTAL DOMAINS OF FUCHSIAN GROUPS  
**Autor:** SCHMUTZ SCHALLER, Paul  
**Kapitel:** 6. Applications  
**DOI:** <https://doi.org/10.5169/seals-64444>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 6. APPLICATIONS

LEMMA 13. *Let  $M$  be a closed hyperbolic surface of genus  $g$  which has  $2g - 2$  simple closed geodesics  $u_1, \dots, u_{2g-2}$  which all intersect in the same point  $Q$  and intersect in no other point. Then  $M$  has simple closed curves  $u_{2g-1}$  and  $u_{2g}$ , passing through  $Q$ , such that the curves  $u_i$  intersect in no other point than  $Q$ ,  $i = 1, \dots, 2g$ . Moreover,  $u_{2g-1}$  and  $u_g$  can be chosen such that*

$$M \setminus \bigcup_{i=1}^{2g} u_i$$

*is the interior of a canonical polygon  $P(g)$ .*

*Proof.* Cut  $M$  along  $u_1$ , the result is a hyperbolic surface  $M_1$  with boundary and genus  $g - 1$ , the boundary consists of two simple closed geodesics  $v_1$  and  $w_1$ . Cut  $M_1$  along  $u_2$ , the result is a hyperbolic surface  $M_2$  with one boundary component  $v_2$  and genus  $g - 1$ . Now cut  $M$  along all  $2g - 2$  simple closed geodesics  $u_1, \dots, u_{2g-2}$ . By induction, the result is a hyperbolic surface  $M_{2g-2}$  with one boundary component  $v$  and genus 1. More precisely, the boundary  $v$  is piecewise geodesic with  $4g - 4$  pieces and we may assume that the notation is chosen such that these pieces appear on  $v$  in the order (the pieces are called like the corresponding closed curves)  $u_1, u_2, \dots, u_{2g-2}, u_1, u_2, \dots, u_{2g-2}$  (note that closed geodesics intersect transversally). Denote by  $S$  and  $S'$  the two copies of  $Q$  on  $v$  between  $u_1$  and  $u_{2g-2}$ . Let  $u_{2g-1}$  be a simple geodesic in  $M_{2g-2}$  which joins  $S$  and  $S'$  such that  $u_{2g-1}$  is not homotopic to a part of  $v$ . Cut  $M_{2g-2}$  along  $u_{2g-1}$ . The result is a hyperbolic surface  $M_{2g-1}$  of genus zero with two boundary components  $w$  and  $w'$  which both consist of  $2g - 1$  geodesic pieces in the order  $u_1, u_2, \dots, u_{2g-2}, u_{2g-1}$ . Denote by  $R$  and  $R'$  the copies of  $Q$  between  $u_1$  and  $u_{2g-1}$  on  $w$  and  $w'$ , respectively. Let  $u_{2g}$  be a simple geodesic in  $M_{2g-1}$  which joins  $R$  and  $R'$ ,  $u_{2g}$  can be chosen such that when we cut  $M_{2g-1}$  along  $u_{2g}$ , then we obtain the interior of a canonical polygon as desired.  $\square$

DEFINITION. A *hyperelliptic surface* is a closed hyperbolic surface of genus  $g$  which has an isometry  $\phi$  with  $\phi^2 = id$  and with exactly  $2g + 2$  fixed points.

In [14], the equivalence of (i) and (ii) of the following theorem was first proved. With the approach chosen here, we can give a third equivalence and

a different proof.

**THEOREM 14.** *Let  $M$  be a closed hyperbolic surface  $M$  of genus  $g$ . Then the following conditions are equivalent.*

- (i)  $M$  is hyperelliptic.
- (ii)  $M$  has a set of at least  $2g-2$  simple closed geodesics which all intersect in the same point and intersect in no other point.
- (iii)  $M$  has a corresponding canonical polygon with equal opposite angles ( $\alpha_i = \alpha_{2g+i}$ ,  $i = 1, \dots, 2g$ ).

*Proof.* I shall prove (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (i).

Let  $M$  be hyperelliptic. Let  $R_i$ ,  $i = 1, \dots, 2g+2$ , be the fixed points of a hyperelliptic involution  $\phi$ . Let  $c_1$  be a simple geodesic segment from  $R_1$  to  $R_2$ . Then  $c_1 \cup \phi(c_1)$  is a simple closed geodesic  $u_1$  since  $\phi^2 = id$ . It also follows that on  $u_1$ , there are only two fixed points of  $\phi$  and that  $M_1 = M \setminus u_1$  is connected. Therefore, we can choose a simple geodesic segment  $c_2$  from  $R_1$  to  $R_3$  which intersects  $u_1$  only in  $R_1$ . By the same argument as above,  $c_2 \cup \phi(c_2)$  is a simple closed geodesic,  $M_2 = M \setminus (u_1 \cup u_2)$  is connected and on  $u_1 \cup u_2$ , there are only three fixed points of  $\phi$ . Continuing this construction we can find simple closed geodesics  $u_1, \dots, u_{2g-2}$  which all intersect in  $R_1$  and in no other point. This proves (i)  $\Rightarrow$  (ii).




FIGURE 6

The partition of a canonical polygon  $P(g)$  into two  $(2g-1)$ -gons and two quadrilaterals

Assume now that  $M$  has  $2g-2$  simple closed geodesics  $u_1, \dots, u_{2g-2}$  which all intersect in the same point  $Q$  and intersect in no other point. By Lemma 13 we then can find simple closed curves  $u_{2g-1}$  and  $u_{2g}$  such that

$$M \setminus \bigcup_{i=1}^{2g} u_i$$

is the interior of a canonical polygon  $P(g)$  with the usual notation. For  $i = 1, \dots, 4g$ , let  $\{Q_i\} = a_i \cap a_{i+1}$ . In  $P(g)$  let  $d_1$  be the geodesic segment from  $Q_{4g}$  to  $Q_{2g-2}$ ,  $d_2$  the geodesic segment from  $Q_{2g}$  to  $Q_{4g-2}$ , and  $d$  the geodesic segment from  $Q_{2g}$  to  $Q_{4g}$ , compare Figure 6. Then  $P(g) \setminus (d_1 \cup d_2 \cup d)$  has four connected components, two quadrilaterals  $W_j$  having  $d$  and  $d_j$ ,  $j = 1, 2$ , among the sides and two  $(2g - 1)$ -gons  $V_j$  having  $d_j$  among the sides,  $j = 1, 2$ . Since  $u_i$ ,  $i = 1, \dots, 2g - 2$ , are simple closed geodesics, it follows that  $\alpha_i = \alpha_{i+2g}$  for  $i = 1, \dots, 2g - 3$ . This implies that  $V_1$  and  $V_2$  are isometric and that  $d_1$  and  $d_2$  have the same length. Therefore,  $W_1$  and  $W_2$  are quadrilaterals with equal lengths of the four sides. Fix now  $W_1$  and try to vary  $W_2$  such that the lengths of the sides remain invariant and so that property (V) for canonical polygons holds. This is certainly the case if  $W_2$  and  $W_1$  are isometric. But then Corollary 8 implies that this is the unique possibility. Therefore,  $W_1$  and  $W_2$  must be isometric and hence  $\alpha_i = \alpha_{i+2g}$  for all  $i = 1, \dots, 2g$ , which proves (ii)  $\Rightarrow$  (iii).

Now assume that (iii) holds. Let  $d$  be the geodesic segment from  $Q_{2g}$  to  $Q_{4g}$ . Then  $d$  separates  $P(g)$  into two isometric  $(2g + 1)$ -gons and the  $\pi$ -rotation around the centre  $C$  of  $d$  induces an isometry  $\phi$  of  $M$  with  $\phi^2 = id$ . The fixed points of  $\phi$  are  $C$ , the point  $Q$  corresponding to the vertices of  $P(g)$  as well as the centres of the sides  $a_i$ ,  $i = 1, \dots, 2g$ . Therefore,  $\phi$  is a hyperelliptic involution which proves (iii)  $\Rightarrow$  (i).  $\square$

**COROLLARY 15.** *All closed hyperbolic surfaces of genus 2 are hyperelliptic.*

*Proof.* All closed hyperbolic surfaces have two simple closed geodesics which intersect in a unique point. The corollary follows by Theorem 14.  $\square$

**DEFINITION.** Let  $M_0$  be a closed hyperbolic surface in  $T_g$ . For every  $M \in T_g$  fix a homeomorphism  $\phi_M$ , homotopic to the identity, from  $M_0$  to  $M$  ( $\phi_M$  exists since closed surfaces of the same genus are homeomorphic). Let  $u$  be a simple closed geodesic in  $M_0$ . Then, in the homotopy class of  $\phi_M(u)$  there exists a unique simple closed geodesic which is denoted by  $\Phi_M(u)$ . The function

$$L(u): T_g \rightarrow \mathbf{R}$$

which associates to  $M$  the length of  $\Phi_M(u)$  is called a *geodesic length function*.

REMARK. It is well known that  $T_g$  can be parametrized by a finite number of geodesic length functions, see for example [12], [13] where it is shown that  $T_g$  can be parametrized by  $6g - 5$  geodesic length functions.

THEOREM 16. *The Teichmüller space  $T_g$  for  $g = 2$  can be parametrized by 7 (suitably chosen) geodesic length functions  $L(u_1), \dots, L(u_7)$ , taken as homogeneous parameters (which means that  $L(u_1)/L(u_7), \dots, L(u_6)/L(u_7)$  gives a parametrization of  $T_2$ ).*

*Proof.* Let  $P(2)$  be a canonical polygon corresponding to a closed hyperbolic surface  $M_0$  of genus 2. As usual let  $Q_i = a_i \cap a_{i+1}$ ,  $i = 1, \dots, 8$ , where the  $a_i$  are the sides of  $P(2)$ . Let  $b_i$  be the geodesic segment (in  $P(2)$ ) between  $Q_i$  and  $Q_{i+4}$ ,  $i = 1, \dots, 4$ . By Corollary 15,  $M_0$  is hyperelliptic, therefore (compare Theorem 14)  $b_i$  corresponds to a simple closed geodesic in  $M_0$ , denoted by  $B_i$ ,  $i = 1, \dots, 4$ . It also follows by Theorem 14 that  $a_i$  corresponds to a simple closed geodesic in  $M_0$ , denoted by  $A_i$ ,  $i = 1, \dots, 4$ .




FIGURE 7

A triangulation of a canonical polygon  $P(g)$  for  $g = 2$

I now prove that the 7 length functions, given by the simple closed geodesics  $A_i$ ,  $i = 1, 2, 3$ ,  $B_i$ ,  $i = 1, \dots, 4$ , taken as homogeneous parameters, give a parametrization of  $T_2$ . In order to do this, it is enough (by Theorem 11 and Corollary 12) to show that  $P(2)$  is uniquely determined by the lengths of  $a_i$ ,  $i = 1, 2, 3$ ,  $b_i$ ,  $i = 1, \dots, 4$ , taken as homogeneous parameters (in the sequel I shall refer to these lengths calling them “the seven lengths”). This can be done analogously as in the proof of Theorem 11. The geodesic segments  $b_i$ ,  $i = 1, \dots, 4$ , intersect in a point  $C$ , the “centre” of  $P(2)$ , and they separate

$P(2)$  into 8 triangles  $D_j$  so that  $a_j$  is a side of  $D_j$ ,  $j = 1, \dots, 8$ , compare Figure 7. Since  $M$  is hyperelliptic,  $D_j$  and  $D_{j+4}$  are isometric,  $j = 1, \dots, 4$ . Denote by  $\delta_i$  the angle of  $D_i$  in the vertex  $C$ ,  $i = 1, \dots, 4$ . The seven lengths determine the triangles  $D_i$ ,  $i = 1, 2, 3$ , as well as two sides and the angle  $\delta_4$  of  $D_4$  by the condition

$$(6) \quad \Delta := \sum_{j=1}^4 \delta_j = \pi,$$

so they determine also  $D_4$ . This shows that the seven lengths determine  $P(2)$ . Multiply the seven lengths by a positive real  $t$  and assume that the seven new lengths also determine a canonical polygon  $P_t(2)$ . If  $t > 1$ , then  $\delta_i$ ,  $i = 1, 2, 3$ , are smaller in  $P_t(2)$  than in  $P(2)$  by Lemma 9, therefore, by (6),  $\delta_4$  is larger in  $P_t(2)$  than in  $P(2)$ . It follows by Lemma 7 that the sum of the two other angles of  $D_4$  is smaller in  $P_t(2)$  than in  $P(2)$ . Since all angles in  $D_i$ ,  $i = 1, 2, 3$ , are smaller in  $P_t(2)$  than in  $P(2)$  by Lemma 9, it follows that

$$\sum_{i=1}^4 \alpha_i$$

is smaller in  $P_t(2)$  than in  $P(2)$ . But this contradicts condition (II) of canonical polygons. An analogous contradiction follows if  $t < 1$  proving thus that  $t = 1$  and therefore the theorem.  $\square$

REMARK. Theorem 16 is new. It is well known that  $6g-6$  length functions can never parametrize  $T_g$  so that the situation of Theorem 16 is the best we can expect. It is not known whether  $6g-5$  geodesic length functions, *taken as homogeneous parameters*, can parametrize  $T_g$  for  $g \geq 3$ .

## REFERENCES

- [1] BEARDON, A.F. *The Geometry of Discrete Groups*. Springer, 1983.
- [2] BUSER, P. *Geometry and Spectra of Compact Riemann Surfaces*. Birkhäuser, Boston, 1992.
- [3] COLDEWEY, H.-D. Kanonische Polygone endlich erzeugter Fuchsscher Gruppen. Dissertation, Bochum, 1971.
- [4] FORD, L. *Automorphic Functions*. Chelsea, New York, 1929.
- [5] IVERSEN, B. *Hyperbolic Geometry*. Cambridge University Press, 1992.
- [6] JOST, J. *Compact Riemann Surfaces*. Springer, 1997.
- [7] KATOK, S. *Fuchsian Groups*. The University of Chicago Press, 1992.