

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	46 (2000)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	GEOMETRIC K-THEORY FOR LIE GROUPS AND FOLIATIONS
Autor:	BAUM, Paul / CONNES, Alain
Kapitel:	4. Solvable simply connected Lie groups
DOI:	https://doi.org/10.5169/seals-64793

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

When G has torsion, the map $K_*^\tau([EG \times X]/G) \rightarrow K^*(X, G)$ can fail to be an isomorphism. The simplest example of this is obtained by taking X to be a point and $G = \mathbf{Z}/2\mathbf{Z}$.

When G has torsion, $K_*^\tau([EG \times X]/G)$ appears to be only a first approximation to $K^*(X, G)$ and $K_*[C_0(X) \rtimes G]$. The key point is that when G has torsion, there will be proper G -manifolds on which the G -action is not free.

4. SOLVABLE SIMPLY CONNECTED LIE GROUPS

The conjecture stated in §2 above is verified for (connected) solvable simply connected Lie groups by

PROPOSITION 1. *Let G be a (connected) solvable simply connected Lie group, and let X be a G -manifold. Then there is a commutative diagram*

$$\begin{array}{ccc} K^*(X, G) & \xrightarrow{\mu} & K_*[C_0(X) \rtimes G] \\ \downarrow & & \downarrow \\ K^*(X) & \longrightarrow & K_*[C_0(X)] \end{array}$$

in which each arrow is an isomorphism.

The proof depends on

LEMMA 2. *Let G be a (connected) solvable simply connected Lie group, and let Z be a proper G -manifold. Then there exists a G -map from Z to G .*

Proof of Lemma 2. Since the action of G on Z is proper all isotropy groups are compact. G has no non-trivial compact subgroups, so the action of G on Z is free. Therefore Z is a principal G -bundle with base Z/G . As G is itself a contractible space on which G acts freely, there is a G -map from Z to G . \square

Proof of Proposition 1. In the diagram of the proposition the right vertical arrow is the Thom isomorphism of [13]. The lower horizontal arrow is the standard isomorphism which is valid for any locally compact Hausdorff topological space.

To define the left vertical arrow the first step is to use the lemma to construct an isomorphism

$$(1) \quad K^*(X, G) \rightarrow K_G^*(T^*[X \times G] \oplus \pi_1^* T^* X).$$

Here G acts on $X \times G$ by

$$(x, g_1)g = (xg, g_1g)$$

and $\pi_1: X \times G \rightarrow X$ is the projection.

If (Z, ξ, f) is a K -cocycle for (X, G) then according to the lemma there exists a G -map $\psi: Z \rightarrow G$. Define $h: Z \rightarrow X \times G$ by $h(z) = (fz, \psi z)$ so that there is the evident commutative diagram

$$\begin{array}{ccc} Z & \xrightarrow{h} & X \times G \\ f \searrow & & \swarrow \pi_1 \\ & X & \end{array}$$

The isomorphism (1) is

$$(Z, \xi, f) \rightarrow h_!(\xi).$$

Next, $T^*[X \times G] \oplus \pi_1^* T^* X$ has a G -invariant Spin^c -structure so by the Thom isomorphism theorem of §2, there is an isomorphism

$$(2) \quad K_G^*(T^*[X \times G] \oplus \pi_1^* T^* X) \cong K_G^*(X \times G).$$

Finally, the action of G on $X \times G$ is free and has $[X \times G]/G = X$. This yields an isomorphism

$$(3) \quad K_G^*(X \times G) \cong K^*(X).$$

Composing (1), (2), (3) gives the left vertical arrow of the proposition. \square

REMARK 3. The two vertical arrows in the diagram of the proposition are not quite canonical. First an orientation must be chosen for the Lie algebra of G . There is no dimension shift in the horizontal arrows of the proposition. If $\epsilon = \dim(G)$, then the left vertical arrow maps $K^i(X, G)$ to $K^{i+\epsilon}(X)$, and the right vertical arrow maps $K_i[C_0(X) \rtimes G]$ to $K_{i+\epsilon}[C_0(X)]$.