Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: p-ADIC L-FUNCTION OF TWO VARIABLES
Autor: Fox, Glenn J.
Kapitel: 2.2 Generalized Bernoulli polynomials
DOI: https://doi.org/10.5169/seals-64800

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 07.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Since any Dirichlet character χ is multiplicative, we must have $\chi(-1) = \pm 1$. A character χ is said to be odd if $\chi(-1) = -1$, and even if $\chi(-1) = 1$.

2.2 Generalized Bernoulli polynomials

Let χ be a Dirichlet character with conductor f_χ. Then we define the functions, $B_{n,\chi}(t)$, $n \in \mathbf{Z}$, $n \geq 0$, by the generating function

$$\sum_{a=1}^{f_\chi} \frac{\chi(a) x e^{(a+t)x}}{e^{f_\chi x} - 1} = \sum_{n=0}^{\infty} B_{n,\chi}(t) \frac{x^n}{n!}, \quad |x| < \frac{2\pi}{f_\chi}. \tag{1}$$

We define the generalized Bernoulli numbers associated with χ, $B_{n,\chi}$, $n \in \mathbf{Z}$, $n \geq 0$, by

$$\sum_{a=1}^{f_\chi} \frac{\chi(a) x e^{ax}}{e^{f_\chi x} - 1} = \sum_{n=0}^{\infty} B_{n,\chi} \frac{x^n}{n!}, \quad |x| < \frac{2\pi}{f_\chi},$$

so that $B_{n,\chi}(0) = B_{n,\chi}$. Note that

$$\sum_{a=1}^{f_\chi} \frac{\chi(a) x e^{(a+t)x}}{e^{f_\chi x} - 1} = e^{tx} \sum_{a=1}^{f_\chi} \frac{\chi(a) x e^{ax}}{e^{f_\chi x} - 1},$$

which implies that

$$\sum_{n=0}^{\infty} B_{n,\chi}(t) \frac{x^n}{n!} = e^{tx} \sum_{n=0}^{\infty} B_{n,\chi} \frac{x^n}{n!},$$

and from this we obtain

$$B_{n,\chi}(t) = \sum_{m=0}^{n} \binom{n}{m} B_{n-m,\chi} t^m. \tag{2}$$

Thus the functions $B_{n,\chi}(t)$, defined in (1), are actually polynomials, called the generalized Bernoulli polynomials associated with χ. Let $\mathbf{Z}[\chi]$ denote the ring generated over $\mathbf{Z}$ by all the values $\chi(a)$, $a \in \mathbf{Z}$, and $\mathbf{Q}(\chi)$ the field generated over $\mathbf{Q}$ by all such values. Then it can be shown that $f_\chi B_{n,\chi}$ must be in $\mathbf{Z}[\chi]$ for each $n \geq 0$ whenever $\chi \neq 1$. In general, we have $B_{n,\chi} \in \mathbf{Q}(\chi)$ for each $n \geq 0$, and so $B_{n,\chi}(t) \in \mathbf{Q}(\chi)[t]$. The polynomials $B_{n,\chi}(t)$ exhibit the property that, for all $n \geq 0$,

$$B_{n,\chi}(-t) = (-1)^n \chi(-1) B_{n,\chi}(t), \tag{3}$$

whenever $\chi \neq 1$. Thus $B_{n,\chi}(t)$, for $\chi \neq 1$, is either an even function or an odd function according to whether $(-1)^n \chi(-1)$ is 1 or -1. From (3) we obtain

$$B_{n,\chi} = (-1)^n \chi(-1) B_{n,\chi},$$

and so $B_{n,\chi} = 0$ whenever n is even and χ is odd, or whenever n is odd and χ is even, $\chi \neq 1$. Another property that the polynomials satisfy is that for $m \in \mathbf{Z}$, $m \geq 1$,

$$B_{n,\chi}(mf_\chi + t) - B_{n,\chi}(t) = n \sum_{a=1}^{mf_\chi} \chi(a)(a+t)^{n-1}, \tag{4}$$

for all $n \geq 0$. This can be derived from (1). Note that for $\chi = 1$ and $t = 0$ this becomes

$$\frac{1}{n}\left(B_{n,1}(m) - B_{n,1}\right) = \sum_{a=1}^{m} a^{n-1}.$$

If $\chi \neq 1$, then it can be shown that $\sum_{a=1}^{f_\chi} \chi(a) = 0$, and from the above relations we can derive

$$B_{0,\chi} = \frac{1}{f_\chi} \sum_{a=1}^{f_\chi} \chi(a)$$

for all χ. Therefore

$$B_{0,\chi} = \begin{cases} 0, & \text{if } \chi \neq 1 \\ 1, & \text{if } \chi = 1. \end{cases}$$

The ordinary Bernoulli polynomials, $B_n(t)$, $n \in \mathbf{Z}$, $n \geq 0$, are defined by

$$\frac{xe^{tx}}{e^x - 1} = \sum_{n=0}^{\infty} B_n(t) \frac{x^n}{n!}, \quad |x| < 2\pi, \tag{5}$$

and the Bernoulli numbers, B_n, $n \in \mathbf{Z}$, $n \geq 0$,

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}, \quad |x| < 2\pi.$$

From this we obtain the values $B_0 = 1$, $B_1 = -1/2$, $B_2 = 1/6$, $B_4 = -1/30, \ldots$, with $B_n = 0$ for odd $n \geq 3$. For even $n \geq 2$, we have

$$B_n = -\frac{1}{n+1} \sum_{m=0}^{n-1} \binom{n+1}{m} B_m.$$

Note that we again have the relations $B_n(0) = B_n$ and

$$B_n(t) = \sum_{m=0}^{n} \binom{n}{m} B_{n-m} t^m,$$

as we did for the generalized Bernoulli polynomials.

Some of the more important properties of Bernoulli polynomials are that

$$B_n(t+1) - B_n(t) = nt^{n-1} \tag{6}$$

for all $n \geq 1$, and

$$B_n(1-t) = (-1)^n B_n(t)$$

for $n \geq 0$. Each of these results can be derived from the generating function (5) above.

Similar to (4) for the generalized Bernoulli polynomials, whenever $m, n \in \mathbf{Z}$, $m \geq 1$, $n \geq 1$,

$$\frac{1}{n}(B_n(m) - B_n) = \sum_{a=0}^{m-1} a^{n-1},$$

where we take 0^0 to be 1 in the case of $a = 0$ and $n = 1$. Note that this can be derived from (6) since

$$B_n(m) - B_n = \sum_{a=0}^{m-1} (B_n(a+1) - B_n(a)) .$$

The Bernoulli numbers are rational numbers, and, in fact, the von Staudt-Clausen theorem states that for even $n \geq 2$,

$$B_n + \sum_{\substack{p \text{ prime} \\ (p-1)|n}} \frac{1}{p} \in \mathbf{Z}.$$

Thus the denominator of each B_n must be square-free.

The ordinary Bernoulli numbers are related to the generalized Bernoulli numbers in that for $\chi = 1$ we have

$$\frac{xe^x}{e^x - 1} = \sum_{n=0}^{\infty} B_{n,1} \frac{x^n}{n!}, \quad |x| < 2\pi,$$

and since

$$\frac{xe^x}{e^x - 1} = x + \frac{x}{e^x - 1},$$

we see that $B_{n,1} = B_n$ for all $n \neq 1$, and $B_{1,1} = -B_1$. In fact, this can be written as $B_{n,1} = (-1)^n B_n$, and for the polynomials, $B_{n,1}(t) = (-1)^n B_n(-t)$.