

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: p-ADIC L-FUNCTION OF TWO VARIABLES
Autor: Fox, Glenn J.
Kapitel: 4.1 A SYMMETRY PROPERTY IN t
DOI: <https://doi.org/10.5169/seals-64800>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

where the power series converges in the domain \mathfrak{D} , and

$$a_{-1}(\tau) = \begin{cases} 1 - \frac{1}{p}, & \text{if } \chi = 1 \\ 0, & \text{if } \chi \neq 1. \end{cases} \quad \square$$

Since $L_p(s, \tau; \chi)$ is defined for each $\tau \in \mathbf{C}_p$ such that $|\tau|_p \leq 1$, we now have a p -adic function of two variables, $L_p(s, t; \chi)$, where $s \in \mathfrak{D}$, $s \neq 1$ if $\chi = 1$, and $t \in \mathbf{C}_p$ with $|t|_p \leq 1$.

4. PROPERTIES OF $L_p(s, t; \chi)$

Most of the properties that follow are direct consequences of similar properties that hold for the generalized Bernoulli polynomials. In all of the following we will take p prime and χ a Dirichlet character with conductor f_χ .

4.1 A SYMMETRY PROPERTY IN t

The first property we obtain regarding $L_p(s, t; \chi)$ is a direct consequence of the generalized Bernoulli polynomials being either odd or even functions, except when $\chi = 1$. Recall that $L_p(s, t; \chi)$ interpolates the values

$$(18) \quad L_p(1 - n, t; \chi) = -\frac{1}{n} b_n(t),$$

for $n \in \mathbf{Z}$, $n \geq 1$, and $t \in \mathbf{C}_p$, $|t|_p \leq 1$, where

$$(19) \quad b_n(t) = B_{n, \chi_n}(qt) - \chi_n(p) p^{n-1} B_{n, \chi_n}(p^{-1}qt),$$

and we define

$$(20) \quad c_n(t) = \sum_{m=0}^n \binom{n}{m} (-1)^{n-m} b_m(t).$$

LEMMA 4.1. For all $n \in \mathbf{Z}$, $n \geq 0$, we have

$$B_{n,1}(-t) = (-1)^n B_{n,1}(t) - (-1)^n n t^{n-1}.$$

Proof. This holds for $n = 0$ since $B_{0,1}(t) = 1$. Now assume that $n \geq 1$. Because $B_{n,1} = 0$ for odd $n \geq 3$, we can write (2) in the form

$$B_{n,1}(t) = \sum_{\substack{m=0 \\ n-m \text{ even}}}^n \binom{n}{m} B_{n-m,1} t^m + n B_{1,1} t^{n-1}.$$

Any m such that $n - m$ is even must have the same parity as n . Thus

$$\begin{aligned} B_{n,1}(-t) &= (-1)^n \sum_{\substack{m=0 \\ n-m \text{ even}}}^n \binom{n}{m} B_{n-m,1} t^m + (-1)^{n-1} n B_{1,1} t^{n-1} \\ &= (-1)^n B_{n,1}(t) - 2(-1)^n n B_{1,1} t^{n-1}. \end{aligned}$$

From the value $B_{1,1} = -B_1 = 1/2$, the lemma then follows. \square

LEMMA 4.2. For all $n \in \mathbf{Z}$, $n \geq 0$,

$$b_n(-t) = \chi(-1)b_n(t).$$

Proof. This is obviously true for $n = 0$ since

$$b_0(t) = (1 - \chi(p)p^{-1}) B_{0,\chi},$$

and $B_{0,\chi} = 0$ except when $\chi = 1$, in which case $B_{0,1} = 1$. So we can assume that $n \geq 1$.

First consider the case of $\chi_n = 1$. This implies that $\chi = \omega^n$. By Lemma 4.1,

$$\begin{aligned} b_n(-t) &= B_{n,1}(-qt) - p^{n-1} B_{n,1}(-p^{-1}qt) \\ &= (-1)^n B_{n,1}(qt) - (-1)^n n (qt)^{n-1} \\ &\quad - p^{n-1} \left((-1)^n B_{n,1}(p^{-1}qt) - (-1)^n n (p^{-1}qt)^{n-1} \right) \\ &= (-1)^n \left(B_{n,1}(qt) - p^{n-1} B_{n,1}(p^{-1}qt) \right) \\ &= (-1)^n b_n(t). \end{aligned}$$

Since $\chi = \omega^n$ and $\omega(-1) = -1$, the lemma holds for $\chi_n = 1$.

Now suppose that $\chi_n \neq 1$. Then, from (3),

$$\begin{aligned} b_n(-t) &= B_{n,\chi_n}(-qt) - \chi_n(p)p^{n-1} B_{n,\chi_n}(-p^{-1}qt) \\ &= (-1)^n \chi_n(-1) \left(B_{n,\chi_n}(qt) - \chi_n(p)p^{n-1} B_{n,\chi_n}(p^{-1}qt) \right) \\ &= (-1)^n \chi_n(-1) b_n(t). \end{aligned}$$

Note that $\chi_n = \chi\omega^{-n}$, which implies that $\chi_n(-1) = (-1)^n \chi(-1)$. Thus the lemma also holds for $\chi_n \neq 1$.

Since the lemma holds for both $\chi_n = 1$ and $\chi_n \neq 1$, the proof must be complete. \square

Using this result, we can prove

THEOREM 4.3. *Let $t \in \mathbf{C}_p$, $|t|_p \leq 1$, and $s \in \mathfrak{D}$, except $s \neq 1$ if $\chi = 1$. Then*

$$L_p(s, -t; \chi) = \chi(-1)L_p(s, t; \chi).$$

Proof. From Lemma 4.2 we see that

$$b_n(-t) = \chi(-1)b_n(t).$$

Also, (20) implies that

$$c_n(-t) = \chi(-1)c_n(t).$$

From (16), whenever $n \geq -1$,

$$a_n(-t) = \chi(-1)a_n(t),$$

which implies that

$$L_p(s, -t; \chi) = \chi(-1)L_p(s, t; \chi). \quad \square$$

If $\chi(-1) = -1$ and $t = 0$, then

$$L_p(s, 0; \chi) = -L_p(s, 0; \chi),$$

which implies that

$$L_p(s; \chi) = -L_p(s; \chi),$$

and thus $L_p(s; \chi) = 0$ for all $s \in \mathfrak{D}$, as we would expect.

4.2 $L_p(s, t; \chi)$ AS A POWER SERIES IN $t - \alpha$, $\alpha \in \mathbf{C}_p$, $|\alpha|_p \leq 1$

To develop $L_p(s, t; \chi)$ in terms of a power series in t will enable us to find a derivative of this function with respect to this second variable. All this we shall do, but before doing so we need to specify some notation.

LEMMA 4.4. *Let $t \in \mathbf{C}_p$, $|t|_p \leq 1$. Then for $n \in \mathbf{Z}$, $n \geq 1$,*

$$\lim_{s \rightarrow 1-n} \binom{-s}{n} L_p(s+n, t; \chi) = -\frac{1}{n} (1 - \chi(p)p^{-1}) B_{0, \chi}.$$

Proof. Recall that, from Theorem 3.13, we can write

$$L_p(s, t; \chi) = \frac{a_{-1}(t)}{s-1} + \sum_{m=0}^{\infty} a_m(t)(s-1)^m,$$

where $a_{-1}(t) = (1 - \chi(p)p^{-1})B_{0, \chi}$. Thus