Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: p-ADIC L-FUNCTION OF TWO VARIABLES
Autor: Fox, Glenn J.
Kapitel: 4.4 Generalized Bernoulli power series
DOI: https://doi.org/10.5169/seals-64800

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Now let $\tau \in pq^{-1}F_0\mathbf{Z}_p$, and let $\{\tau_i\}_{i=1}^{\infty}$ be a sequence in $pq^{-1}F_0\mathbf{Z}$, with $\tau_i > 0$ for each i, such that $\tau_i \to \tau$. We are working with polynomials, so that

$$\lim_{i\to\infty}\left(\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau_i) - \binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0)\right) = \binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau) - \binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0),$$

which must be in $\mathbf{Z}_p[\chi]$ since the limit of any sequence in $\mathbf{Z}_p[\chi]$ must also be in $\mathbf{Z}_p[\chi]$. Now let n' be a positive integer, and consider

$$\lim_{i\to\infty}\left(\left(\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau_i)-\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0)\right)-\left(\binom{q^{-1}\Delta_c}{k}\beta_{n',\chi}(\tau_i)-\binom{q^{-1}\Delta_c}{k}\beta_{n',\chi}(0)\right)\right)$$
$$=\left(\left(\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau)-\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0)\right)-\left(\binom{q^{-1}\Delta_c}{k}\beta_{n',\chi}(\tau)-\binom{q^{-1}\Delta_c}{k}\beta_{n',\chi}(0)\right)\right).$$

The quantity on the left must be 0 modulo $q\mathbf{Z}_p[\chi]$, which implies that the value of

$$\binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(\tau) - \binom{q^{-1}\Delta_c}{k}\beta_{n,\chi}(0)$$

modulo $q\mathbf{Z}_p[\chi]$ is independent of n. $\square$

4.4 Generalized Bernoulli power series

In [9] we find a definition of ordinary Bernoulli numbers of negative index, B_{-n}, where $n \in \mathbf{Z}$, $n \geq 1$, in the field $\mathbf{Q}_p$, given by

$$B_{-n} = \lim_{k\to\infty} B_{\phi(p^k)-n}, \tag{26}$$

where the limit is taken in a p-adic sense. Note that $\phi(p^k) \to 0$ in $\mathbf{Z}_p$ as $k \to \infty$. Since $|B_m|_p$ is bounded for all $m \in \mathbf{Z}$, $m \geq 0$, we must have

$$\begin{aligned} B_{-n} &= \lim_{k\to\infty}\left(1 - p^{\phi(p^k)-n-1}\right)B_{\phi(p^k)-n} \\ &= \lim_{k\to\infty} -\left(\phi\left(p^k\right)-n\right)L_p\left(1-\left(\phi\left(p^k\right)-n\right);\omega^{-n}\right) \\ &= nL_p\left(n+1;\omega^{-n}\right). \end{aligned}$$

implying that the limit exists and can be described in familiar terms.

Recall that $B_m = 0$ for any odd $m \in \mathbf{Z}$, $m \geq 3$. Thus (26) implies that $B_{-n} = 0$ for any odd $n \in \mathbf{Z}$, $n \geq 1$. Furthermore, we have the following:

THEOREM 4.13. *Let $n \in \mathbf{Z}$ be even, $n \geq 2$. Then*

$$B_{-n} + \sum_{\substack{r \text{ prime} \\ (r-1)|n}} \frac{1}{r} \in \mathbf{Z}_p ,$$

where each prime r is taken to be a rational prime.

REMARK. Since $1/r \in \mathbf{Z}_p$ for any rational prime $r \neq p$, this implies that $B_{-n} + 1/p \in \mathbf{Z}_p$ whenever $(p-1) \mid n$, and $B_{-n} \in \mathbf{Z}_p$ otherwise.

Proof. By the von Staudt-Clausen theorem, we know that

$$B_m + \sum_{\substack{r \text{ prime} \\ (r-1)|m}} \frac{1}{r} \in \mathbf{Z}$$

for any even $m \in \mathbf{Z}$, $m \geq 2$.

Let $n \in \mathbf{Z}$ be even, $n \geq 2$. For any integer $k \geq 2$, $\phi(p^k)$ is even and $(p-1) \mid \phi(p^k)$. Thus $\phi(p^k) - n$ is even, and $(p-1) \mid n$ if and only if $(p-1) \mid (\phi(p^k) - n)$. Therefore, if k is sufficiently large,

$$B_{\phi(p^k)-n} + \sum_{\substack{r \text{ prime} \\ (r-1)|n}} \frac{1}{r} \in \mathbf{Z}_p ,$$

and the result follows from (26). $\square$

In a similar manner we define generalized Bernoulli numbers of negative index, $B_{-n,\chi}$, where $n \in \mathbf{Z}$, $n \geq 1$, in the field $\mathbf{C}_p$ according to

$$B_{-n,\chi} = \lim_{k \to \infty} B_{\phi(p^k)-n,\chi} , \tag{27}$$

where the limit is once again taken in a p-adic sense. For each $m \in \mathbf{Z}$, $m \geq 0$, the quantity $|B_{m,\chi}|_p$ is bounded. Thus, since $\chi_{\phi(p^k)} = \chi$ for all characters χ and for all $k \in \mathbf{Z}$, $k \geq 1$, we can write

$$\begin{aligned} B_{-n,\chi} &= \lim_{k \to \infty} \left(1 - \chi_{\phi(p^k)}(p) p^{\phi(p^k)-n-1}\right) B_{\phi(p^k)-n,\chi_{\phi(p^k)}} \\ &= \lim_{k \to \infty} -\left(\phi\left(p^k\right) - n\right) L_p\left(1 - \left(\phi\left(p^k\right) - n\right); \chi_n\right) \\ &= n L_p\left(n+1; \chi_n\right), \end{aligned}$$

so that the limit exists. Since $B_{\phi(p^k)-n,1} = B_{\phi(p^k)-n}$ for $n, k \in \mathbf{Z}$, with $n \geq 1$ and k sufficiently large, we obtain $B_{-n,1} = B_{-n}$ for all such n.

If $k \geq 2$, then $\phi(p^k)$ is even. Thus n and $\phi(p^k) - n$ are of the same parity. Recall that

$$\delta_\chi = \begin{cases} 1, & \text{if } \chi \text{ is odd} \\ 0, & \text{if } \chi \text{ is even}. \end{cases}$$

Then $B_{\phi(p^k)-n,\chi} = 0$ whenever $n \not\equiv \delta_\chi \pmod 2$, provided $\phi(p^k) - n > 1$. Because of this, the relation (27) implies that $B_{-n,\chi} = 0$ whenever $n \not\equiv \delta_\chi \pmod 2$ for all $n \in \mathbf{Z}$, $n \geq 1$. Furthermore, we can obtain

THEOREM 4.14. *Let χ be such that $\chi \neq 1$, and let $n \in \mathbf{Z}$, $n \geq 1$. Then $f_\chi B_{-n,\chi} \in \mathbf{Z}_p[\chi]$.*

Proof. Recall that when $\chi \neq 1$, $f_\chi B_{m,\chi} \in \mathbf{Z}[\chi]$ for all $m \in \mathbf{Z}$, $m \geq 0$. Thus

$$f_\chi B_{-n,\chi} = \lim_{k\to\infty} f_\chi B_{\phi(p^k)-n,\chi}$$

must be in the p-adic completion of $\mathbf{Z}[\chi]$ for any $n \in \mathbf{Z}$, $n \geq 1$. Since the p-adic completion of $\mathbf{Z}[\chi]$ is $\mathbf{Z}_p[\chi]$, the theorem must hold. $\square$

We now define what we shall refer to as generalized Bernoulli power series of negative index in $\mathbf{Z}_p[\chi]$. For $n \in \mathbf{Z}$, $n \geq 1$, and for $t \in \mathbf{C}_p$, $|t|_p \leq |q|_p$, let

$$B_{-n,\chi}(t) = \lim_{k\to\infty} B_{\phi(p^k)-n,\chi}(t).$$

Then

$$\begin{aligned} B_{-n,\chi}(qt) &= \lim_{k\to\infty} \left(B_{\phi(p^k)-n,\chi_{\phi(p^k)}}(qt) - \chi_{\phi(p^k)}(p)p^{\phi(p^k)-n-1}B_{\phi(p^k)-n,\chi_{\phi(p^k)}}(p^{-1}qt)\right) \\ &= \lim_{k\to\infty} -(\phi(p^k) - n)L_p(1 - (\phi(p^k) - n), t; \chi_n) \\ &= nL_p(n+1, t; \chi_n). \end{aligned}$$

Since $L_p(n+1, t; \chi_n)$ exists for each $n \in \mathbf{Z}$, $n \geq 1$, and $t \in \mathbf{C}_p$, $|t|_p \leq 1$, we see that $B_{-n,\chi}(qt)$ must also exist for such t. Thus $B_{-n,\chi}(t)$ exists for $t \in \mathbf{C}_p$, $|t|_p \leq |q|_p$. Now, by Theorem 4.5, we can expand this quantity as a power series, obtaining

$$\begin{aligned} B_{-n,\chi}(qt) &= n\sum_{m=0}^{\infty} \binom{-(n+1)}{m} q^m t^m L_p(n+m+1; \chi_{n+m}) \\ &= n\sum_{m=0}^{\infty} \binom{-(n+1)}{m} q^m t^m \frac{1}{n+m} B_{-(n+m),\chi} \\ &= \sum_{m=0}^{\infty} \binom{-n}{m} B_{-(n+m),\chi} q^m t^m. \end{aligned}$$

Since $|B_{-(n+m),\chi}|_p \leq \max\{|p|_p^{-1}, |f_\chi|_p^{-1}\}$ and

$$\binom{-n}{m} = (-1)^m \binom{n+m-1}{m},$$

this sum converges for $|qt|_p < 1$. Thus we have the relation

$$B_{-n,\chi}(t) = \sum_{m=0}^{\infty} \binom{-n}{m} B_{-n-m,\chi} t^m, \tag{28}$$

converging for all $t \in \mathbf{C}_p$, $|t|_p < 1$. Note that this is in the same form as (2) for the generalized Bernoulli polynomials having positive index, which we can rewrite as

$$B_{n,\chi}(t) = \sum_{m=0}^{\infty} \binom{n}{m} B_{n-m,\chi} t^m,$$

since $\binom{n}{m} = 0$ for $m, n \in \mathbf{Z}$, $m > n \geq 0$. By setting $t = 0$ in (28), we see that $B_{-n,\chi}(0) = B_{-n,\chi}$ for all $n \in \mathbf{Z}$, $n \geq 1$.

THEOREM 4.15. *Let $n \in \mathbf{Z}$, $n \geq 1$. Then for any $m \in \mathbf{Z}$, $m \geq 1$, such that $q \mid mf_\chi$,*

$$B_{-n,\chi}(mf_\chi) - B_{-n,\chi}(0) = -n \sum_{\substack{a=1 \\ (a,p)=1}}^{mf_\chi} \chi(a) a^{-n-1}.$$

Proof. By definition, since $|mf_\chi|_p \leq |q|_p$,

$$\begin{aligned} B_{-n,\chi}(mf_\chi) - B_{-n,\chi}(0) &= \lim_{k\to\infty} \left(B_{\phi(p^k)-n,\chi}(mf_\chi) - B_{\phi(p^k)-n,\chi}(0)\right) \\ &= \lim_{k\to\infty} \left(\phi(p^k) - n\right) \sum_{a=1}^{mf_\chi} \chi(a) a^{\phi(p^k)-n-1}, \end{aligned}$$

following from (4). Now, $v_p(\phi(p^k)) = k-1$, and $a^{\phi(p^k)} \equiv 1 \pmod{p^k}$ for $(a,p) = 1$. These imply that

$$\lim_{k\to\infty} \left(\phi(p^k) - n\right) \sum_{a=1}^{mf_\chi} \chi(a) a^{\phi(p^k)-n-1} = -n \sum_{\substack{a=1 \\ (a,p)=1}}^{mf_\chi} \chi(a) a^{-n-1},$$

completing the proof. $\square$

THEOREM 4.16. *Let $n \in \mathbf{Z}$, $n \geq 1$. Then for all χ and for all $t \in \mathbf{C}_p$, $|t|_p < 1$,*

$$B_{-n,\chi}(-t) = (-1)^n \chi(-1) B_{-n,\chi}(t) .$$

Proof. Since

$$B_{-n,\chi}(t) = \sum_{m=0}^{\infty} \binom{-n}{m} B_{-n-m,\chi} t^m ,$$

and $B_{-n-m,\chi} = 0$ whenever $n+m \not\equiv \delta_\chi \pmod 2$ for each $m \in \mathbf{Z}$, $m \geq 1$, we see that $B_{-n,\chi}(t)$ is either an odd or an even function according to whether $n + \delta_\chi$ is odd or even, respectively. Thus

$$\begin{aligned} B_{-n,\chi}(-t) &= (-1)^{n+\delta_\chi} B_{-n,\chi}(t) \\ &= (-1)^n \chi(-1) B_{-n,\chi}(t) , \end{aligned}$$

and the proof is complete. □

REFERENCES

[1] ANKENY, N., E. ARTIN and S. CHOWLA. The class number of real quadratic number fields. *Ann. of Math. (2) 56* (1952), 479–493.

[2] BARSKY, D. Sur la norme de certaines séries d'Iwasawa (une démonstration analytique p-adique du théorème de Ferrero-Washington). *Study group on ultrametric analysis, 10th year: 1982/83, No. 1.* Inst. Henri Poincaré, Paris, 1984.

[3] BERGER, A. Recherches sur les nombres et les fonctions de Bernoulli. *Acta Math. 14* (1890/1891), 249–304.

[4] BERNOULLI, J. *Ars Conjectandi.* Basel, 1713. Reprinted in *Die Werke von Jakob Bernoulli.* Vol. 3. Birkhäuser, Basel, 1975.

[5] CARLITZ, L. Arithmetic properties of generalized Bernoulli numbers. *J. reine angew. Math. 202* (1959), 174–182.

[6] COMTET, L. *Advanced Combinatorics. The Art of Finite and Infinite Expansions.* Revised and enlarged edition. D. Reidel Publishing Co., Dordrecht, 1974.

[7] FRESNEL, J. Nombres de Bernoulli et fonctions L p-adiques. *Ann. Inst. Fourier (Grenoble) 17* (1967), fasc. 2, 281–333 (1968).

[8] GOUVÊA, F. Q. *p-adic Numbers. An Introduction.* Universitext. Springer, Berlin, 1993.

[9] GRANVILLE, A. Arithmetic properties of binomial coefficients. I. Binomial coefficients modulo prime powers. *Organic Mathematics (Burnaby, BC, 1995).* CMS Conf. Proc. 20, Amer. Math. Soc., Providence, RI, 1997, 253–276.