

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: IDEAL SOLUTIONS OF THE TARRY-ESCOTT PROBLEM OF DEGREE FOUR AND A RELATED DIOPHANTINE SYSTEM
Autor: Choudhry, Ajai
Kapitel: 4. The diophantine system $\sum_{i=1}^5 a_i^r = \sum_{i=1}^5 b_i^5$,
 $\quad r=1,2,3,4,6$
DOI: <https://doi.org/10.5169/seals-64802>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We may apply Frolov's theorem to the above solution to obtain other non-symmetric solutions. For instance, an arbitrary constant K can be added to all the terms $a_i, b_i, i = 1, 2, 3, 4, 5$.

As a numerical example, taking $m = 3, n = 1$, we get, on suitable re-arrangement, the following solution:

$$\begin{aligned} (-1659)^r + 1406^r + 2784^r + 4025^r + 5915^r \\ = (-1675)^r + 1659^r + 2366^r + 4256^r + 5865^r, \end{aligned}$$

where $r = 1, 2, 3, 4$. Adding the constant 1676 to all the terms, we get the following solution in positive integers:

$$17^r + 3082^r + 4460^r + 5701^r + 7591^r = 1^r + 3335^r + 4042^r + 5932^r + 7541^r,$$

where $r = 1, 2, 3, 4$.

4. THE DIOPHANTINE SYSTEM $\sum_{i=1}^5 a_i^r = \sum_{i=1}^5 b_i^r, \quad r = 1, 2, 3, 4, 6$

We will now state the theorem given by Gloden [7, p.24] to which a reference has already been made in the introduction and then apply it to obtain a parametric solution of this diophantine system.

THEOREM 4.1. *If*

$$\sum_{i=1}^{k+1} a_i^r = \sum_{i=1}^{k+1} b_i^r, \quad r = 1, 2, \dots, k$$

then

$$\sum_{i=1}^{k+1} (a_i + t)^r = \sum_{i=1}^{k+1} (b_i + t)^r, \quad r = 1, 2, \dots, k, k + 2,$$

where

$$t = -\left(\sum_{i=1}^{k+1} a_i\right)/(k + 1).$$

As we have already obtained, in the preceding section, a parametric solution of $\sum_{i=1}^5 a_i^r = \sum_{i=1}^5 b_i^r, r = 1, 2, 3, 4$, a direct application of the above theorem gives a parametric solution of $\sum_{i=1}^5 a_i^r = \sum_{i=1}^5 b_i^r, r = 1, 2, 3, 4$ and 6. We multiply the $(a_i + t), (b_i + t), i = 1, 2, 3, 4, 5$ by 5 to cancel out denominators,

and we rename the resulting expressions as a_i , b_i , $i = 1, 2, 3, 4, 5$, so that the parametric solution of the diophantine system

$$\sum_{i=1}^5 a_i^r = \sum_{i=1}^5 b_i^r, \quad r = 1, 2, 3, 4, 6$$

may be written as

$$\begin{aligned} a_1 &= 4m^8 - 30m^7n + 98m^6n^2 - 34m^5n^3 - 9m^4n^4 \\ &\quad - 80m^3n^5 + 97m^2n^6 - 6mn^7 - 16n^8, \\ a_2 &= 4m^8 - 20m^7n + 63m^6n^2 - 184m^5n^3 + 116m^4n^4 \\ &\quad + 60m^3n^5 - 53m^2n^6 - 26mn^7 + 24n^8, \\ a_3 &= -16m^8 + 53m^6n^2 + 46m^5n^3 - 279m^4n^4 \\ &\quad + 460m^3n^5 - 388m^2n^6 + 144mn^7 - 16n^8, \\ a_4 &= -16m^8 + 90m^7n - 262m^6n^2 + 361m^5n^3 - 279m^4n^4 \\ &\quad + 145m^3n^5 - 73m^2n^6 + 54mn^7 - 16n^8, \\ a_5 &= 24m^8 - 40m^7n + 48m^6n^2 - 189m^5n^3 + 451m^4n^4 \\ &\quad - 585m^3n^5 + 417m^2n^6 - 166mn^7 + 24n^8, \\ b_1 &= 24m^8 - 40m^7n + 23m^6n^2 + 36m^5n^3 - 284m^4n^4 \\ &\quad + 520m^3n^5 - 393m^2n^6 + 134mn^7 - 16n^8, \\ b_2 &= -16m^8 - 10m^7n + 138m^6n^2 - 254m^5n^3 + 391m^4n^4 \\ &\quad - 540m^3n^5 + 437m^2n^6 - 166mn^7 + 24n^8, \\ b_3 &= 4m^8 - 70m^7n + 188m^6n^2 - 229m^5n^3 + 121m^4n^4 \\ &\quad + 55m^3n^5 - 143m^2n^6 + 74mn^7 - 16n^8, \\ b_4 &= 4m^8 + 20m^7n - 127m^6n^2 + 86m^5n^3 + 121m^4n^4 \\ &\quad - 260m^3n^5 + 172m^2n^6 - 16mn^7 - 16n^8, \\ b_5 &= -16m^8 + 100m^7n - 222m^6n^2 + 361m^5n^3 - 349m^4n^4 \\ &\quad + 225m^3n^5 - 73m^2n^6 - 26mn^7 + 24n^8. \end{aligned}$$

As a numerical example, when $m = 3$ and $n = 1$, we get, on removal of common factors and suitable re-arrangement, the following solution:

$$\begin{aligned} &1449^r + 7654^r + 17104^r + (-5441)^r + (-20766)^r \\ &= 8809^r + 16854^r + (-641)^r + (-4176)^r + (-20846)^r, \end{aligned}$$

where $r = 1, 2, 3, 4$ and 6 .