

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE SPECTRAL MAPPING THEOREM, NORMS ON RINGS, AND RESULTANTS
Autor: Laksov, D. / Svensson, L. / Thorup, A.
Kapitel: 3. A REDUCTION
DOI: <https://doi.org/10.5169/seals-64805>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. A REDUCTION

To prove the Spectral Mapping Theorem it suffices to verify that it holds for the polynomial ring $k[t_0, \dots, t_m]$ in variables t_0, \dots, t_m over k , and for the polynomial $F(x) = t_0 + t_1x + \dots + t_mx^m$. This is because any polynomial $G(x) = b_0 + b_1x + \dots + b_mx^m$ with coefficients in a ring R containing k as a subring is the image of $F(x)$ by the map $g: k[t_0, \dots, t_m][x] \rightarrow R[x]$ defined by $g(a) = a$ for $a \in k$, by $g(t_i) = b_i$ for $i = 0, \dots, m$, and by $g(x) = x$. If we can prove the equality $\det F(M) = \prod_{i=1}^n F(\lambda_i)$ in $k[t_0, \dots, t_m]$ we obtain that $\det G(M) = g(\det F(M)) = \prod_{i=1}^n g(F(\lambda_i)) = \prod_{i=1}^n G(\lambda_i)$ in R .

4. THE PROOF

Clearly (2.1) holds when F is a constant a where it simply states that $\det(aI_n) = a^n$. We shall prove (2.1) for polynomials F of degree $m > 0$ by induction on m .

We first note that if $F(x)$ has a root λ in R , so that $F(x) = (x - \lambda)G(x)$ in $R[x]$, then (2.1) holds for $F(x)$. Indeed, $G(x)$ is of degree $m - 1$ so it follows from the induction hypothesis that $\det G(M) = \prod_{i=1}^n G(\lambda_i)$. Since $F(M) = (M - \lambda I_n)G(M)$ we obtain:

$$\begin{aligned} \det F(M) &= \det(M - \lambda I_n) \det G(M) \\ &= \prod_{i=1}^n (\lambda_i - \lambda) \prod_{i=1}^n G(\lambda_i) = \prod_{i=1}^n (\lambda_i - \lambda) G(\lambda_i) = \prod_{i=1}^n F(\lambda_i). \end{aligned}$$

As we saw in Section 3 it suffices to prove the result for the ring $Q = k[t_0, \dots, t_m]$ and the polynomial $F(x) = t_0 + t_1x + \dots + t_mx^m$. Let x and y be independent variables over the ring Q . The polynomial $F(x) - F(y)$ in x with coefficients in $Q[y]$ has the root $x = y$. Hence, as we just observed, (2.1) holds for the polynomial $F(x) - F(y)$. We obtain the equation:

$$(4.1) \quad \det(F(M) - F(y)I_n) = \prod_{i=1}^n (F(\lambda_i) - F(y))$$

in $Q[y]$.

The equation (2.1) is a consequence of (4.1). To see this we observe that $F(y)$ in $Q[y]$ is transcendental over Q , that is the element $F(y)$ in $Q[y]$ does not satisfy a polynomial relation $a_0 + a_1F(y) + \dots + a_lF(y)^l = 0$ with coefficients a_i in Q and $a_l \neq 0$, because the coefficient $a_l t_m^l$ of the highest