

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	46 (2000)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE SPECTRAL MAPPING THEOREM, NORMS ON RINGS, AND RESULTANTS
Autor:	Laksov, D. / Svensson, L. / Thorup, A.
Kapitel:	6. NORMS AND RESULTANTS
DOI:	https://doi.org/10.5169/seals-64805

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5.1. LEMMA. *Let N be a norm of degree n on a k -algebra A . For each element α of A the characteristic polynomial $P_\alpha^N(t) = N_{k[t]}(t - \alpha)$ is monic of degree n .*

Moreover, the trace Tr^N is a k -linear map $A \rightarrow k$.

Proof. Let s, t, u, v be independent variables over the ring k . For each element β in A the norm $N_{k[s,t,u]}(t - \alpha s - \beta u)$ is a polynomial in $k[s, t, u]$. Since N is of degree n we have that $N_{k[s,t,u,v]}(vt - \alpha vs - \beta vu) = v^n N_{k[s,t,u]}(t - \alpha s - \beta u)$. It follows that $N_{k[s,t,u]}(t - \alpha s - \beta u)$ is homogeneous of degree n in $k[s, t, u]$. In particular the coefficient of t^{n-1} is of the form $as + bu$ with a and b in k . By evaluating the polynomial $N_{k[s,t,u]}(t - \alpha s - \beta u)$ at $s = 0, u = 0$, it follows that the coefficient to t^n is equal to 1. Hence $N_{k[t]}(t - \alpha)$ is a monic polynomial of degree n , and $a = -\text{Tr}^N(\alpha)$. Similarly, $b = -\text{Tr}^N(\beta)$. Hence we have that $\text{Tr}^N(\alpha s + \beta u) = -(as + bu) = \text{Tr}^N(\alpha)s + \text{Tr}^N(\beta)t$. Specializing s and t to any pair of elements of k the second part of the Lemma follows. \square

5.2. EXAMPLE. Let M be a free module of rank n over k , or more generally a projective k -module of constant rank n . Then the determinant defines a norm of degree n on $\text{End}_k(M)$.

Let A be a k -algebra which is free of rank n as a k -module. Left multiplication by elements of A define an injection $A \rightarrow \text{End}_k(A)$ of k -algebras. By restriction we obtain a norm of degree n on A .

6. NORMS AND RESULTANTS

Let $F(x) = f_0 + \cdots + f_m x^m$ and $P(x) = p_0 + \cdots + p_n x^n$ be polynomials of degree m , respectively n in the k -algebra $k[x]$ of polynomials in the variable x with coefficients in k . The *resultant* $\text{Res}(F, P)$ of F and P is the determinant of the $(m+n) \times (m+n)$ -matrix $D(F, P)$ whose columns are the coefficients of the polynomials $F, xF, \dots, x^{n-1}F, P, xP, \dots, x^{m-1}P$. Note that the definition is asymmetric in F and P in the sense that $\text{Res}(F, P) = (-1)^{mn} \text{Res}(P, F)$.

When P is monic the resultant is equal to the determinant of the endomorphism induced by multiplication by F on the free k -module $k[x]/(P(x))$ of rank n . To see this we note that for $i = 0, \dots, n-1$ we can write $x^i F = Q_i P + R_i$ in $k[x]$, where $Q_i(x)$ and $R_i(x)$ are of degrees at most $m-1$, respectively $n-1$. It follows that the determinant of $D(F, P)$ is equal to the determinant of the $(m+n) \times (m+n)$ -matrix $B(F, P)$ whose columns are the

coefficients of the polynomials $R_0, \dots, R_{n-1}, P, xP, \dots, x^{m-1}P$. We see that the $n \times n$ -block $C(F, P)$ in the upper left corner of $B(F, P)$ is the matrix $C(F, P)$ of the map induced by multiplication by F on $k[x]/(P(x))$, and the $m \times m$ -block in the lower right corner is upper triangular with 1's on the diagonal. Moreover, the entries of $C(F, P)$ are the only non-zero entries in the first n columns of $B(F, P)$. It follows that $\text{Res}(F, P) = \det C(F, P)$, as we claimed.

6.1. EXAMPLE. When P is a monic polynomial we saw in Example 5.2 that the k -algebra $k[x]/(P(x))$ which is free of rank n as a k -module has a canonical norm. Via the canonical map $k[x] \rightarrow k[x]/(P(x))$ we obtain a canonical norm N'_P on $k[x]$. The above interpretation of the resultant can then be written as

$$(6.1.1) \quad (N'_P)_R(F) = \text{Res}(F, P)$$

for all commutative k -algebras R and all polynomials $F(x)$ in $R[x] = R \otimes_k k[x]$. By an easy computation of the determinant defining $\text{Res}(t - x, P)$, we obtain that the characteristic polynomial of x with respect to N'_P is

$$P_x^{N'_P}(t) = P(t).$$

6.2. EXAMPLE. We shall introduce a second important norm on $k[x]$. Let $P(x)$ be a monic polynomial of degree n in the k -algebra $k[x]$. There is a canonical ring extension $k \subseteq k' = k[\lambda_1, \dots, \lambda_n]$ such that $P(x)$ splits as $P(x) = \prod_{i=1}^n (x - \lambda_i)$ in $k'[x]$. The extension is obtained by induction starting with $k = k_0$ and $P_0(x) = P(x)$, and constructing $k_i = k[\lambda_1, \dots, \lambda_i]$ and $P_i(x) \in k[\lambda_1, \dots, \lambda_i][x]$ from k_{i-1} and P_{i-1} , for $i = 1, 2, \dots, n$, by $k_i = k_{i-1}[x]/(P_{i-1}(x)) = k_{i-1}[\lambda_i]$, where λ_i is the class of x , and by $P_i(x) = P_{i-1}(x)/(x - \lambda_i)$. We note that k' is a free k -module of rank $n!$. The algebra k' is sometimes called the *universal decomposition algebra* for P (see [B1], §6, p. 68).

For every commutative k -algebra R and every polynomial G in $R[x] = R \otimes_k k[x]$ we have that $\prod_{i=1}^n G(\lambda_i)$ is symmetric in $\lambda_1, \dots, \lambda_n$, and consequently lies in the image of the inclusion $R \subseteq k' \otimes_k R$. We obtain a map $(N''_P)_R: R \otimes_k k[x] \rightarrow R$ defined by $(N''_P)_R(G) = \prod_{i=1}^n G(\lambda_i)$. In this way we obtain a norm N''_P of degree n on $k[x]$ and the characteristic polynomial of x with respect to the norm N''_P is

$$P_x^{N''_P}(t) = \prod_{i=1}^n (t - \lambda_i) = P(t).$$