Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 46 (2000)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NOTE OF THE EDITORS

Autor: Laksov, D. / Svensson, L. / Thorup, A. DOI: https://doi.org/10.5169/seals-64806

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 09.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

NOTE OF THE EDITORS

concerning

"The Spectral Mapping Theorem, norms on rings, and resultants" by D. LAKSOV, L. SVENSSON and A. THORUP

The Editors of L'Enseignement Mathématique have decided to publish here the referee's report, as it throws some interesting light on the authors' paper. At the referee's request, it appears without any editing as an anonymous contribution.

The main result, and many other related results, can be deduced from the fact that if one takes a generic square matrix of size n of indeterminates over the integers (call A_n the ring generated by the entries) and forces the characteristic polynomial to factor "generically", i.e., one adjoins n new indeterminates and equates the elementary symmetric functions of these with the coefficients of the characteristic polynomial (with suitable signs), then the resulting ring R_n is an integral domain. This takes a little proving and I do not know a formal reference, but it has been known to me for many years, and I strongly suspect that it is in the literature.

If one knows this (and it does not follow from trivial generalities about integral extensions — the authors are quite right about that), then, because R_n maps to every commutative ring such that a size n matrix has a split characteristic polynomial, all questions about polynomial identities in the entries of the matrix and its eigenvalues immediately reduce to the question of whether the identities hold in the generic case, i.e. for R_n . But since the generic ring is an integral domain, one may pass to its fraction field, which has characteristic 0, and even to the complex numbers. (The argument works even if the generic ring is merely reduced, i.e., has no nonzero nilpotents, since then it injects into a product of fields.)

Here is a sketch of a proof that R_n is a domain. First off, if one starts with any ring A and elements $a_1, \ldots, a_n \in A$, and forms the quotient $A[Y_1, \ldots, Y_n]/(e_i - a_i)$, where the e_i are the elementary symmetric functions

 $Y_1 + \cdots + Y_n, \ldots, Y_1 \cdot \ldots \cdot Y_n$ of the Y_i , the quotient ring is a finitely generated free module over A, of rank n!. Killing $e_1 - a_1$ gives a monic equation of degree 1 for Y_1 in terms of the other variables. Then killing $e_2 - a_2$ gives a monic quadratic equation for Y_2 in terms of the remaining variables. Etc.

From this it follows that R_n is free of rank n! over A_n . Thus, the issue of whether it is a domain is unaffected by inverting the nonzero elements of A_n , and one may pass to (i.e., tensor with) the fraction field B of A_n . Let L be the splitting field of the characteristic polynomial over B. It will suffice to see that the B-algebra homomorphism from $B \otimes R_n \to L$ that sends the images of the Y_i to the eigenvalues of the matrix in L is an isomorphism, and it is clearly onto.

Thus, it will be an isomorphism provided that L has degree n! over B. We have reduced to showing that the splitting field of the characteristic polynomial of a generic matrix has degree n! over the field generated by the entries. But this follows because the indeterminates can be specialized so as to give the companion matrix of any polynomial. (E.g., specialize the last column to elementary symmetric functions of n variables, the diagonal just below the main diagonal to all 1's, and the other entries to 0.)

So the short argument above (which would be shorter in a research paper version) shows that *every* polynomial identity with integer coefficients relating eigenvalues of a matrix and entries of that matrix which holds for matrices over the complex numbers holds in every commutative ring. There is one extra step needed for the Spectral Mapping Theorem: the coefficients of the polynomial F also need to be introduced as indeterminates, or instead one can point out that the identity really comes down to a finite number of identities saying that coefficients of polynomials in the extra variables are equal.

A simpler example is the Cayley-Hamilton theorem. It suffices to prove the result for a matrix of indeterminates over the integers, and so one is working over a polynomial ring over the integers. Here, one does not need to prove that any auxiliary ring is a domain. Since such a matrix is diagonalizable over an algebraically closed field containing the fraction field of the polynomial ring, one reduces the result for commutative rings to the case of diagonal matrices over a field, which is clear.