

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE WITT GROUP OF LAURENT POLYNOMIALS
Autor: Ojanguren, Manuel / Panin, Ivan
Kapitel: 2.K-theoretic PRELIMINARIES
DOI: <https://doi.org/10.5169/seals-64807>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We now recall three elementary, well-known facts about hermitian spaces.

PROPOSITION 1.5. *Let (P, α) be any space. Then:*

1. *The space $(P, \alpha) \perp (P, -\alpha)$ is hyperbolic.*
2. *If L is a lagrangian of (P, α) , then (P, α) is isometric to $H(L)$.*
3. *If M is a sublagrangian of (P, α) , then the map α induces on M^\perp/M a natural structure of hermitian space that makes it Witt equivalent to (P, α) .*

2. K -THEORETIC PRELIMINARIES

We recall a few results proved in the twelfth chapter of Bass' book [1]. For any ring A we denote by $K_0(A)$ the Grothendieck group of finitely generated projective right A -modules and by $K_1(A)$ the abelianized general linear group of A : $K_1(A) = GL(A)/[GL(A), GL(A)]$. By Whitehead's lemma $K_1(A)$ is also the quotient of $GL(A)$ by the subgroup $E(A)$ generated by all elementary matrices over A .

For any functor F from rings to abelian groups we denote by $N_+F(A)$ the kernel of the map $F(A[t]) \rightarrow F(A)$ obtained by putting $t = 0$. Similarly, we denote by $N_-F(A)$ the kernel of $F(A[t^{-1}]) \rightarrow F(A)$ obtained by putting $t^{-1} = 0$. The inclusions of $A[t]$ and $A[t^{-1}]$ into $A[t, t^{-1}]$ define a map

$$N_+F(A) \oplus N_-F(A) \longrightarrow F(A[t, t^{-1}])$$

whose cokernel will be denoted by $LF(A)$. The functor LK_1 turns out to be naturally isomorphic to K_0 , hence we will denote LK_i by K_{i-1} for $i = 1$ and also for $i = 0$.

THEOREM 2.1. *Let A be any associative ring.*

(a) *For $i = 0$ or 1 there exists a natural embedding*

$$\lambda_i: K_{i-1}(A) \longrightarrow K_i(A[t, t^{-1}])$$

such that the composite

$$K_{i-1}(A) \xrightarrow{\lambda_i} K_i(A[t, t^{-1}]) \rightarrow LK_i(A) = K_{i-1}(A)$$

is the identity.

(b) *The embedding λ_i and the canonical homomorphism*

$$N_{\pm}K_i(A) \rightarrow K_i(A[t, t^{-1}])$$

yield canonical decompositions

$$K_1(A[t, t^{-1}]) = K_1(A) \oplus N_+K_1(A) \oplus N_-K_1(A) \oplus K_0(A)$$

and

$$K_0(A[t, t^{-1}]) = K_0(A) \oplus N_+K_0(A) \oplus N_-K_0(A) \oplus K_{-1}(A).$$

Proof. See [1], Theorem 7.4 of chapter XII. \square

We will also use the following well-known result.

PROPOSITION 2.2. *If 2 is invertible in A, the groups $N_{\pm}K_1(A)$ are uniquely divisible by 2.*

Proof. By [1], XII, 5.3, every element of $N_+K_1(A)$ can be represented by a matrix $\alpha = 1 + \nu t$, with ν a nilpotent matrix of $M_n(A)$. Let

$$P(X) = \sum_0^{\infty} \binom{1/2}{n} X^n \in \mathbf{Z}[1/2][X].$$

Then $P(\nu t) \in M_n(A[t])$ and $(P(\nu t))^2 = 1 + \nu t$. This shows that $N_+K_1(A)$ is divisible by 2. To show uniqueness it suffices to show that $N_+K_1(A)$ has no 2-torsion. Take $\alpha = 1 + \nu t$ as before and suppose that $\alpha^2 \in E(A[t])$. Put $s = t(2 + \nu t)$, so that $\alpha^2 = 1 + \nu s$. Since

$$t = \sum_1^{\infty} \binom{1/2}{n} \nu^{n-1} s^n$$

we have $M_n(A)[t] = M_n(A)[s]$. If $\alpha^2 = 1 + \nu s \in E(A[s]) = E(M_n(A)[s])$ we clearly also have $\alpha = 1 + \nu t \in E(M_n(A)[t])$. \square

COROLLARY 2.3. *If 2 is invertible in A, the groups $N_{\pm}K_0(A)$ are uniquely divisible by 2.*

Proof. $K_0(A)$ is a direct factor of $K_1(A[X, X^{-1}])$, hence $N_{\pm}K_0(A)$ is a direct factor of $N_{\pm}K_1(A[X, X^{-1}])$. \square

Assume now that A has an involution. Associating to any projective module its dual and to any matrix its conjugate transpose yields actions of $\mathbf{Z}/2$ on K_0 and K_1 which are compatible with the decompositions of Theorem 2.1. From Corollary 2.3 we immediately deduce

COROLLARY 2.4. *Suppose that A is a ring with involution, in which 2 is invertible. Then*

$$H^2(\mathbf{Z}/2, K_0(A[t, t^{-1}])/K_0(A)) = H^2(\mathbf{Z}/2, K_{-1}(A)).$$

3. THE WITT GROUP OF POLYNOMIAL RINGS

THEOREM 3.1. *Let A be an associative ring with involution, in which 2 is invertible. Let ϵ be 1 or -1 and let W be the Witt group functor of ϵ -hermitian spaces. The natural homomorphism*

$$W(A) \longrightarrow W(A[t])$$

is an isomorphism.

Proof. It suffices to show that the homomorphism $W(A[t]) \rightarrow W(A)$ given by the evaluation at $t = 0$ is an isomorphism. Surjectivity is obvious. To prove injectivity let (P, α) be a space over $A[t]$ and $(P(0), \alpha(0))$ its reduction modulo t . Suppose that $(P(0), \alpha(0))$ is isometric to some hyperbolic space $H(Q)$. Choosing a projective module Q' such that $Q \oplus Q'$ is free and adding to (P, α) the space $H(Q'[t])$ we may assume that $P(0)$ is the hyperbolic space over a free module. The class of P in $K_0(A[t])/K_0(A) = N_+(A)$ is a symmetric element. By Corollary 2.4 it can be written as $a + a^*$, hence, adding to (P, α) a suitable free hyperbolic space, we may assume that (P, α) is of the form

$$H(A^n[t]) \perp (R \oplus R^*, \beta).$$

Let R' be an $A[t]$ -module such that $R \oplus R'$ is free. Adding to (P, α) the hyperbolic space $H(R')$ we are reduced to the case in which P is free and α is an invertible ϵ -hermitian matrix with entries in $A[t]$.

LEMMA 3.2. *Let $\alpha = \epsilon\alpha^* \in M_n(A[t])$ be any ϵ -hermitian matrix. There exist an integer m and a matrix $\tau \in \text{GL}_{n+2m}(A[t])$ (actually in $E_{n+2m}(A[t])$) such that*

$$\tau^* \begin{pmatrix} \alpha & 0 \\ 0 & \chi \end{pmatrix} \tau = \alpha_0 + t\alpha_1,$$

where α_0 and α_1 are constant matrices and χ is a sum of hyperbolic blocks $\begin{pmatrix} 0 & 1 \\ \epsilon 1 & 0 \end{pmatrix}$ of various sizes.