

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE WITT GROUP OF LAURENT POLYNOMIALS
Autor: Ojanguren, Manuel / Panin, Ivan
Kapitel: 3. The Witt group of polynomial rings
DOI: <https://doi.org/10.5169/seals-64807>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

COROLLARY 2.4. *Suppose that A is a ring with involution, in which 2 is invertible. Then*

$$H^2(\mathbf{Z}/2, K_0(A[t, t^{-1}])/K_0(A)) = H^2(\mathbf{Z}/2, K_{-1}(A)).$$

3. THE WITT GROUP OF POLYNOMIAL RINGS

THEOREM 3.1. *Let A be an associative ring with involution, in which 2 is invertible. Let ϵ be 1 or -1 and let W be the Witt group functor of ϵ -hermitian spaces. The natural homomorphism*

$$W(A) \longrightarrow W(A[t])$$

is an isomorphism.

Proof. It suffices to show that the homomorphism $W(A[t]) \rightarrow W(A)$ given by the evaluation at $t = 0$ is an isomorphism. Surjectivity is obvious. To prove injectivity let (P, α) be a space over $A[t]$ and $(P(0), \alpha(0))$ its reduction modulo t . Suppose that $(P(0), \alpha(0))$ is isometric to some hyperbolic space $H(Q)$. Choosing a projective module Q' such that $Q \oplus Q'$ is free and adding to (P, α) the space $H(Q'[t])$ we may assume that $P(0)$ is the hyperbolic space over a free module. The class of P in $K_0(A[t])/K_0(A) = N_+(A)$ is a symmetric element. By Corollary 2.4 it can be written as $a + a^*$, hence, adding to (P, α) a suitable free hyperbolic space, we may assume that (P, α) is of the form

$$H(A^n[t]) \perp (R \oplus R^*, \beta).$$

Let R' be an $A[t]$ -module such that $R \oplus R'$ is free. Adding to (P, α) the hyperbolic space $H(R')$ we are reduced to the case in which P is free and α is an invertible ϵ -hermitian matrix with entries in $A[t]$.

LEMMA 3.2. *Let $\alpha = \epsilon\alpha^* \in M_n(A[t])$ be any ϵ -hermitian matrix. There exist an integer m and a matrix $\tau \in \text{GL}_{n+2m}(A[t])$ (actually in $E_{n+2m}(A[t])$) such that*

$$\tau^* \begin{pmatrix} \alpha & 0 \\ 0 & \chi \end{pmatrix} \tau = \alpha_0 + t\alpha_1,$$

where α_0 and α_1 are constant matrices and χ is a sum of hyperbolic blocks $\begin{pmatrix} 0 & 1 \\ \epsilon 1 & 0 \end{pmatrix}$ of various sizes.

Proof of the lemma. Write $\alpha = \gamma + \delta t^N$, where δ is constant and γ of degree less than N . Assume that N is at least 2. Since δ is ϵ -hermitian and 2 is invertible in A we can write $\delta = \sigma + \epsilon\sigma^*$. Then

$$\begin{pmatrix} 1 & t & -\sigma^*t^{N-1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \gamma + \sigma t^N + \epsilon\sigma^*t^N & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & \epsilon & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ t & 1 & 0 \\ -\sigma t^{N-1} & 0 & 1 \end{pmatrix}$$

is of degree $\leq N - 1$ and after $N - 1$ such transformations we get a linear matrix. \square

Writing $\alpha = \alpha_0 + t\alpha_1$ as $\alpha_0(1 + \nu t)$ we see immediately that, α being invertible, ν is nilpotent. The formal power series

$$\tau = (1 + \nu t)^{-1/2} = \sum \binom{-1/2}{k} (\nu t)^k$$

is a polynomial. From $\alpha = \epsilon\alpha^*$ we get $\alpha_0^* = \epsilon\alpha_0$ and $\nu^*\alpha_0^* = \epsilon\alpha_0\nu$. This implies that $\tau^*\alpha_0^* = \epsilon\alpha_0\tau$ and therefore

$$\tau^*\alpha\tau = \tau^*\alpha_0(1 + \nu t)\tau = \alpha_0\tau(1 + \nu t)\tau = \alpha_0.$$

This proves that (P, α) is Witt equivalent to $(P(0), \alpha(0))$ and is, therefore, hyperbolic. \square

4. THE WITT GROUP OF TORSION MODULES

Let M be a finitely generated right $A[t]$ -module and suppose that it is a t -torsion module and that it is projective as an A -module. Obviously, it will be finitely generated over A . We denote by M^\sharp the left $A[t]$ -module $\text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t])$ and we consider it as a right module through the involution on $A[t]$.

Recall that, as an A -module, the quotient $A[t, t^{-1}]/A[t]$ can be written as a direct sum

$$A[t, t^{-1}]/A[t] = At^{-1} \oplus At^{-2} \oplus \dots$$

Thus, to any $f \in \text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t])$ we can associate an A -linear map $f_{-1}: M \rightarrow A$, which is defined as the composite of f with the projection onto At^{-1} .