

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE WITT GROUP OF LAURENT POLYNOMIALS
Autor: Ojanguren, Manuel / Panin, Ivan
Kapitel: 4. The Witt group of torsion modules
DOI: <https://doi.org/10.5169/seals-64807>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proof of the lemma. Write $\alpha = \gamma + \delta t^N$, where δ is constant and γ of degree less than N . Assume that N is at least 2. Since δ is ϵ -hermitian and 2 is invertible in A we can write $\delta = \sigma + \epsilon\sigma^*$. Then

$$\begin{pmatrix} 1 & t & -\sigma^* t^{N-1} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \gamma + \sigma t^N + \epsilon\sigma^* t^N & 0 & 0 \\ 0 & 0 & 1 \\ 0 & \epsilon & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ t & 1 & 0 \\ -\sigma t^{N-1} & 0 & 1 \end{pmatrix}$$

is of degree $\leq N-1$ and after $N-1$ such transformations we get a linear matrix. \square

Writing $\alpha = \alpha_0 + t\alpha_1$ as $\alpha_0(1 + \nu t)$ we see immediately that, α being invertible, ν is nilpotent. The formal power series

$$\tau = (1 + \nu t)^{-1/2} = \sum \binom{-1/2}{k} (\nu t)^k$$

is a polynomial. From $\alpha = \epsilon\alpha^*$ we get $\alpha_0^* = \epsilon\alpha_0$ and $\nu^*\alpha_0^* = \epsilon\alpha_0\nu$. This implies that $\tau^*\alpha_0^* = \epsilon\alpha_0\tau$ and therefore

$$\tau^*\alpha\tau = \tau^*\alpha_0(1 + \nu t)\tau = \alpha_0\tau(1 + \nu t)\tau = \alpha_0.$$

This proves that (P, α) is Witt equivalent to $(P(0), \alpha(0))$ and is, therefore, hyperbolic. \square

4. THE WITT GROUP OF TORSION MODULES

Let M be a finitely generated right $A[t]$ -module and suppose that it is a t -torsion module and that it is projective as an A -module. Obviously, it will be finitely generated over A . We denote by $M^\#$ the left $A[t]$ -module $\text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t])$ and we consider it as a right module through the involution on $A[t]$.

Recall that, as an A -module, the quotient $A[t, t^{-1}]/A[t]$ can be written as a direct sum

$$A[t, t^{-1}]/A[t] = At^{-1} \oplus At^{-2} \oplus \dots.$$

Thus, to any $f \in \text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t])$ we can associate an A -linear map $f_{-1}: M \rightarrow A$, which is defined as the composite of f with the projection onto At^{-1} .

PROPOSITION 4.1. *The map*

$$\partial = \partial_M: M^\# = \text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t]) \longrightarrow \text{Hom}_A(M, A) = M^*$$

obtained by associating f_{-1} to f is a functorial A -linear isomorphism.

Proof. It is clear that ∂ is A -linear. To show that it is bijective we construct its inverse. Given any $g \in M^*$ define \tilde{g} by the (finite !) sum

$$\tilde{g}(x) = t^{-1}g(x) + t^{-2}g(tx) + t^{-3}g(t^2x) + \dots$$

It is easy to check that $\tilde{g} \in M^\#$, $(\tilde{g})_{-1} = g$ and $\tilde{f_{-1}} = f$. Functoriality is clear. \square

COROLLARY 4.2. *For any finitely generated t -torsion module M which is projective as an A -module the canonical homomorphism $M \rightarrow M^{\#\#}$ is an isomorphism.*

Proof. It suffices to remark that the diagram

$$\begin{array}{ccc} M & & \\ \swarrow \text{can} & & \searrow \text{can} \\ M^{\#\#} & \xrightarrow{(\partial_M^*)^{-1} \circ \partial_{M^\#}} & M^{**} \end{array}$$

commutes and that $M \xrightarrow{\text{can}} M^{**}$ is an isomorphism. \square

An ϵ -hermitian t -torsion space (or, briefly, a t -torsion space) is a pair $(M, \langle \cdot, \cdot \rangle)$ consisting of a finitely generated t -torsion right $A[t]$ -module M which is projective as an A -module, and a perfect ϵ -hermitian pairing $\langle \cdot, \cdot \rangle: M \times M \rightarrow A[t, t^{-1}]/A[t]$. Giving $\langle \cdot, \cdot \rangle$ is the same, of course, as giving its adjoint $\varphi: M \rightarrow M^\#$ defined by $\varphi(a)(b) = \langle a, b \rangle$.

Isometries and orthogonal sums are defined in the obvious way. For any subset $X \subset M$ we define its orthogonal as

$$X^\perp = \{y \in M \mid \langle x, y \rangle = 0 \quad \forall x \in X\}.$$

A sublagrangian of (M, φ) is an $A[t]$ -submodule L of M which satisfies the following two conditions :

- (1) It is contained in its own orthogonal: $L \subseteq L^\perp$.
- (2) The quotient M/L is projective over A (which is the same as saying that L , as an A -module, is a direct factor of M).

A sublagrangian L is a *lagrangian* if $L = L^\perp$. A t -torsion space is *metabolic* if it has a lagrangian. The Witt group of t -torsion spaces is the quotient of the Grothendieck group of t -torsion spaces with respect to orthogonal sums, modulo the subgroup generated by the metabolic spaces. We will denote it by $W_{tors}(A[t])$. Lemma 4.6 below will show that the opposite of the class of (M, φ) is the class of $(M, -\varphi)$.

LEMMA 4.3. *Let M and N be finitely generated t -torsion modules and $i: N \rightarrow M$ an $A[t]$ -linear homomorphism. Assume that as A -modules M and N are projective. Then the map $i^\sharp: M^\sharp \rightarrow N^\sharp$ is surjective (respectively injective) if and only if $i^*: M^* \rightarrow N^*$ is surjective (respectively injective).*

Proof. Look:

$$\begin{array}{ccc} M^\sharp & \xrightarrow{i^\sharp} & N^\sharp \\ \partial_M \downarrow & & \downarrow \partial_N \\ M^* & \xrightarrow{i^*} & N^* \end{array}$$

□

PROPOSITION 4.4. *Let (M, φ) be a t -torsion space and L an $A[t]$ -submodule of M . If M/L is projective over A , then $L = L^{\perp\perp}$ and L^\perp is a direct factor of M as an A -module.*

Proof. First observe that as an A -module L is finitely generated and projective. Let $i: L \rightarrow M$ be the natural injection. By Lemma 4.3 the map $i^\sharp \circ \varphi$ is surjective, thus the sequence

$$0 \longrightarrow L^\perp \xrightarrow{j} M \xrightarrow{i^\sharp \circ \varphi} L^\sharp \longrightarrow 0$$

is exact. Hence L^\perp is a direct factor of M as an A -module; in particular it is A -projective. Identifying L with $L^{\sharp\sharp}$ we can write the dual sequence as

$$0 \longrightarrow L \xrightarrow{i} M \xrightarrow{j^\sharp \circ \varphi^\sharp} (L^\perp)^\sharp \longrightarrow 0.$$

Notice that it is exact by Lemma 4.3. Again by Lemma 4.3 the sequence

$$0 \longrightarrow L^{\perp\perp} \longrightarrow M \xrightarrow{j^\sharp \circ \varphi} (L^\perp)^\sharp \longrightarrow 0$$

is exact because L^\perp is a direct factor of M as an A -module. Since $\varphi^\sharp = \pm \varphi$, comparing the last two sequences we get the result. □

We now prove a fundamental result on the equivalence of t -torsion spaces.

THEOREM 4.5. *Let (M, φ) be an ϵ -hermitian t -torsion space and L a sublagrangian of (M, φ) . The quotient L^\perp/L carries a natural structure of t -torsion ϵ -hermitian space and its class in $W_{tors}(A[t])$ is the same as that of (M, φ) .*

Proof. We first prove the following lemma.

LEMMA 4.6. *Let (M, φ) be any ϵ -hermitian t -torsion space. The space $(M, \varphi) \perp (M, -\varphi)$ is metabolic.*

Proof of Lemma 4.6. We show that the image $L = \Delta(M)$ of the diagonal map $M \xrightarrow{\Delta} M \oplus M$ is a lagrangian. The condition $L \subseteq L^\perp$ is immediately verified. The quotient $(M \oplus M)/L$ is isomorphic to M , hence it is projective over A . It remains to see that $L^\perp \subseteq L$. If $(a, b) \in L^\perp$ we have $0 = \langle (a, b), (x, x) \rangle = \langle a - b, x \rangle$ for any $x \in M$. Since the pairing \langle , \rangle is perfect, this implies $a = b$, i.e. $(a, b) \in L$. \square

We now prove the theorem. By Proposition 4.4, L^\perp is a direct factor of M as an A -module. Since $L \subseteq L^\perp$ is also a direct factor of M , the quotient L^\perp/L is projective. Denoting by \bar{a}, \bar{b} the classes modulo L of two elements $a, b \in L$, we define the hermitian structure of L^\perp/L by $\langle \bar{a}, \bar{b} \rangle = \langle a, b \rangle$. It is clear that $\langle a, b \rangle$ only depends on \bar{a} and \bar{b} . We first check that this pairing defines a t -torsion space. It is clearly ϵ -hermitian. The injectivity of the adjoint map $L^\perp/L \rightarrow (L^\perp/L)^\sharp$ follows immediately from Proposition 4.4. To show surjectivity consider any $A[t]$ -linear map $f: L^\perp \rightarrow A[t, t^{-1}]/A[t]$. Since L^\perp is a direct factor of M as an A -module, f , by Lemma 4.3, extends to an $A[t]$ -linear map $\tilde{f}: M \rightarrow A[t, t^{-1}]/A[t]$. Choose an $m \in M$ for which $\tilde{f} = \langle m, \cdot \rangle$. If \tilde{f} vanishes on L , then m is in L^\perp . This proves that L^\perp/L is a t -torsion space.

To show that L^\perp/L is equivalent to (M, φ) we check that the image of the diagonal map $\Delta: L^\perp \rightarrow M \oplus L^\perp/L$ is a lagrangian of $(M, -\varphi) \perp L^\perp/L$ which is, therefore, metabolic. It is easy to check that $\Delta(L^\perp)$ is contained in its own orthogonal. Conversely, if $(a, \bar{b}) \in M \oplus L^\perp/L$ is orthogonal to every (x, \bar{x}) , then $\langle a - b, x \rangle = 0$ for every $x \in L^\perp$. This means that $a - b$ is in $L^{\perp\perp}$, which by Proposition 4.4 coincides with L . We thus have $(a, \bar{b}) = (a, \bar{a}) \in \Delta(L^\perp)$. \square

The next proposition connects the Witt group of t -torsion spaces with the Witt group of A .

PROPOSITION 4.7. *The isomorphisms*

$$\partial_M: \text{Hom}_{A[t]}(M, A[t, t^{-1}]/A[t]) \rightarrow \text{Hom}_A(M, A)$$

induce a surjective homomorphism

$$\partial^W: W_{tors}(A[t]) \rightarrow W(A).$$

Proof. Associating to any t -torsion space (M, φ) the hermitian space $(M, \partial_M \circ \varphi)$ preserves isometries and orthogonal sums and, by Lemma 4.3, transforms metabolic t -torsion spaces into hyperbolic spaces (with the same lagrangian). Therefore it induces a homomorphism

$$\partial^W: W_{tors}(A[t]) \rightarrow W(A).$$

To find a preimage (M, φ) of a space (M, α) over A consider M as an $A[t]$ -module annihilated by t and replace $\alpha: M \rightarrow M^*$ by $\varphi = \partial_M^{-1} \circ \alpha$. \square

5. THE WITT GROUP OF EXTENDED SPACES

Let $W'(A[t, t^{-1}])$ be the group defined in the introduction.

THEOREM 5.1. *Let A be an associative ring with involution, in which 2 is invertible. The homomorphism*

$$\psi: W(A) \oplus W(A) \rightarrow W'(A[t, t^{-1}])$$

mapping (ξ, η) to $\xi + t\eta$ is an isomorphism.

Proof. The injectivity of ψ is based on the following result, whose proof will be given in §6.

PROPOSITION 5.2. *There exists a homomorphism*

$$\text{Res}: W'(A[t, t^{-1}]) \rightarrow W(A)$$

with the following properties:

R_1 : For any constant space $\xi \in W(A) \subset W'(A[t, t^{-1}])$, $\text{Res}(\xi) = 0$.

R_2 : For any constant space $\xi \in W(A) \subset W'(A[t, t^{-1}])$, $\text{Res}(t \cdot \xi) = \xi$.

Proof. See Theorem 6.7. \square