

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 46 (2000)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE SIXTH FERMAT NUMBER AND PALINDROMIC CONTINUED FRACTIONS  
**Autor:** Dyson, Freeman  
**DOI:** <https://doi.org/10.5169/seals-64808>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## THE SIXTH FERMAT NUMBER AND PALINDROMIC CONTINUED FRACTIONS

by Freeman DYSON

ABSTRACT. An elementary argument is presented, verifying the known factorization of the sixth Fermat number  $2^{64} + 1$ . The verification is made more elegant by using an old theorem of Serret concerning palindromic continued fractions.

It is easy to explain why the fifth Fermat number  $2^{32} + 1$  is divisible by 641 [Hardy and Wright, 1938]. The prime 641 has the two representations

$$(1) \quad 641 = 5 \cdot 2^7 + 1 = 5^4 + 2^4.$$

The congruence

$$(2) \quad 2^{32} = 2^4 \cdot 2^{28} \equiv -5^4 \cdot 2^{28} = -(5 \cdot 2^7)^4 \equiv -1 \pmod{641}$$

follows immediately from (1).

The purpose of this note is to explain in a similarly elementary way why the sixth Fermat number  $2^{64} + 1$  is divisible by 274177. The divisor has the representation

$$(3) \quad q = 274177 = 1 + 2^8 f, \quad f = (2^6 - 1)(2^4 + 1),$$

and it is easily verified that

$$(4) \quad 2^{24} - 1 = fg, \quad g = (2^6 + 1)(2^8 - 2^4 + 1).$$

We look for a factorization of the form

$$(5) \quad 2^{64} + 1 = (x^2 + y^2)(z^2 + w^2), \quad 2^{32} - i = (x + iy)(z - iw),$$

so that we require integers  $x, y, z, w$  satisfying

$$(6) \quad xz + yw = 2^{32}, \quad xw - yz = 1, \quad x^2 + y^2 = q.$$

When  $z = gx$ ,  $w = gy$ , the right side of (5),  $(x + iy)g(x - iy)$ , becomes  $gq$ ; and  $gq = 2^{32} + a$ , very close to  $2^{32}$ , with the difference  $a = g - 2^8 = 15409$

by (3) and (4). This implies that the ratios  $(z/x)$  and  $(w/y)$  must both be very close to  $g$ . We therefore try to set

$$(7) \quad z = gx - s, \quad w = gy - t,$$

with integers  $s, t$  expected to be small. Then (5) will hold, provided that

$$(8) \quad x^2 + y^2 = q, \quad xs + yt = a, \quad ys - xt = 1.$$

From (8) it follows that

$$(9) \quad a^2 + 1 = qu, \quad u = s^2 + t^2.$$

Since the integers  $q$  and  $a$  are known and comparatively small, it is not difficult to find an integer solution of (8) and (9) by trial and error, namely  $x = 516$ ,  $y = 89$ ,  $s = 29$ ,  $t = 5$ ,  $u = 866$ . The solution of (8) and (9), with  $z$  and  $w$  given by (7), provides a solution of (5). This completes the verification of the factorization of the sixth Fermat number. The factorization (4) of  $2^{24} - 1$  leads directly to the factorization (5) of  $2^{64} + 1$ .

Instead of using trial and error to solve (8) and (9), we can obtain a solution more elegantly by using an old theorem of Serret [Serret, 1848; Perron, 1954] on continued fractions. In the statement of the theorem,  $q$  and  $a$  are any pair of coprime integers with  $0 < a < q$ . The ratio  $q/a$  may be expressed as a continued fraction with partial quotients  $(a_j, j = 1, \dots, n)$  in precisely two ways, one with  $n$  even and one with  $n$  odd. In one way, the last two quotients are  $[j, 1]$ , where  $j$  may be any integer. In the other way these two quotients are replaced by the single quotient  $[j + 1]$ , all other quotients remaining the same. We say that the continued fraction  $q/a$  with  $n$  quotients is a palindrome if

$$(10) \quad a_j = a_{n+1-j}, \quad j = 1, \dots, n.$$

**SERRET'S THEOREM.** *Suppose that the continued fraction  $q/a$  with  $n$  partial quotients is given. It will be a palindrome if and only if an integer  $u$  exists with*

$$(11) \quad qu = a^2 + (-1)^n.$$

*Proof of theorem.* Let  $A$  be any finite ordered set of positive integers, and let  $S(A)$  be the numerator of the continued fraction whose partial quotients are the members of  $A$ . Euler discovered the explicit formula for  $S(A)$  [Euler, 1764; Perron, 1954],

$$(12) \quad S(A) = \sum_B P(B),$$

where the sum extends over certain ordered subsets  $B$  of  $A$ , and  $P(B)$  is the product of the members of  $B$ . The subsets included in the sum (12) are  $A$  itself, and those  $B$  for which the complement  $A - B$  is a union of non-overlapping pairs, each pair consisting of two consecutive members of  $A$ . The empty set  $E$  will appear in the sum (12) if and only if the number of members of  $A$  is even. Thus, for example,  $S(E) = 1$ ,  $S(a) = a$ ,  $S(a, b) = 1 + ab$ ,  $S(a, b, c) = a + c + abc$ ,  $S(a, b, c, d) = 1 + ab + ad + cd + abcd$ . For a lucid and accessible explanation of Euler's formula, see [Davenport, 1982], chapter 4, pp. 82–84.

Let now  $A$  be the set  $(a_j, j = 1, \dots, n)$  of partial quotients of one of the two continued fractions  $q/a$ . We write

$$(13) \quad N(k, m) = S(a_k, \dots, a_m),$$

so that

$$(14) \quad q = N(1, n), \quad a = N(2, n).$$

Euler's formula (12) implies the symmetry relation

$$(15) \quad S(a_1, \dots, a_n) = S(a_n, \dots, a_1),$$

and the recurrence relations

$$(16) \quad N(1, n) = N(1, k)N(k+1, n) + N(1, k-1)N(k+2, n),$$

$$(17) \quad N(1, n)N(2, n-1) - N(1, n-1)N(2, n) = (-1)^n.$$

Suppose that (11) holds. Define

$$(18) \quad b = N(1, n-1), \quad v = N(2, n-1).$$

The symmetry (15) implies

$$(19) \quad q/b = S(a_n, \dots, a_1)/S(a_{n-1}, \dots, a_1),$$

so that one of the two continued fractions for  $q/b$  has the same partial quotients as the continued fraction for  $q/a$ , in reversed order. Then (11), (17) and (18) imply

$$(20) \quad qv = ab + (-1)^n, \quad q(u-v) = a(a-b).$$

Now  $q > a > 0$  by hypothesis, and  $q > b > 0$  by (19), so that  $q > |a-b|$ . If  $a \neq b$ , the fraction  $(u-v)/(a-b)$  is equal to  $a/q$ , which is impossible because  $a$  and  $q$  are coprime and  $|a-b| < q$ . Therefore  $a = b$ ,  $u = v$ , and the continued fractions for  $q/a$  and  $q/b$  are identical, so that (10) holds and the continued fraction is a palindrome. Conversely, if (10) holds, (13) and (18) imply  $a = b$ , and then (17) implies (11) with  $u = v$  defined by (18).  $\square$

Applying the theorem to  $q$  and  $a$  given by (3) and (9), we deduce that the continued fraction  $q/a$  with  $n$  even must be a palindrome. The partial quotients are easily calculated, with the result

$$(21) \quad q = S(17, 1, 3, 1, 5, 5, 1, 3, 1, 17), \quad a = S(1, 3, 1, 5, 5, 1, 3, 1, 17).$$

We now apply the recurrence relation (16) to (21) with  $k = n/2 = 5$ . As a result of the symmetry (15) and the palindrome property of  $q$ , (16) gives a solution of (8) with

$$(22) \quad \begin{aligned} x &= S(17, 1, 3, 1, 5) = 516, & y &= S(17, 1, 3, 1) = 89, \\ s &= S(1, 3, 1, 5) = 29, & t &= S(1, 3, 1) = 5. \end{aligned}$$

In this way Serret's theorem leads to a solution of (8), following a deductive route rather than trial and error.

The same method leads to the factorization of the fifth Fermat number, starting from

$$(23) \quad q = 1 + 2^7 f, \quad 2(2^8 - 1) = fg, \quad f = 2^2 + 1, \quad g = 2(2^2 - 1)(2^4 + 1).$$

The palindromic continued fraction  $q/a$  has numerator  $q = S(4, 6, 6, 4)$ , and the factorization of  $2^{32} + 1$  results with  $x = S(4, 6) = 25$ ,  $y = S(4) = 4$ ,  $s = S(6) = 6$ ,  $t = S(E) = 1$ ,  $x^2 + y^2 = 641$ . But in this case the argument using (1) and (2) gives the same result more quickly. It would be more interesting if we could understand in a similar way the factorizations of the seventh and higher Fermat numbers. For a summary of the known factorizations, see [Brent, 1999]. The seventh Fermat number has two prime factors with 17 and 22 decimal digits.

According to (21), the partial quotients of  $q/a$  in the case of the sixth Fermat number are all odd, while in the case of the fifth Fermat number the partial quotients are all even. To decide whether this unexpected behavior of the partial quotients is a numerical accident or a general rule, we need to find more examples.

The factorization of the sixth Fermat number was originally published in [Landry, 1880], without any explanation of how it was found. For a conjectured reconstruction of Landry's method see [Williams, 1993]. Landry worked for several months to find the factorization. The argument presented in this note only verifies the factorization after the factors are known.

ACKNOWLEDGMENT. The author is grateful to a referee for informing him of the existence of Serret's theorem and providing the references to Perron, Serret and Davenport.

## REFERENCES

- BRENT, R. P., 1999. Factorization of the tenth Fermat number. *Math. Comp.* 68 (1999), 429–451.
- DAVENPORT, H. *The Higher Arithmetic*. 5<sup>th</sup> edition, Cambridge University Press, Cambridge, 1982.
- EULER, L. Specimen algorithmi singularis. (1764; commentatio 281 indicis Ene-strœmiani.) *Novi Commentarii Academiæ Scientiarum Imperialis Petropolitanae* 9 (1762–1763), 53–69. Reprinted in L. Euler, *Opera Omnia*, series I, vol. 15. (Ed. Georg Faber.) B.G. Teubner, Leipzig and Berlin, 1927. (In the *Opera Omnia* edition, our equations (12) and (15) appear on page 36, (17) on page 41, (16) on page 47.)
- HARDY, G. H. and E. M. WRIGHT. *An Introduction to the Theory of Numbers*. Oxford University Press, Oxford, 1938. (For the fifth and sixth Fermat numbers, see pp. 14–15.)
- LANDRY, F. Sur la décomposition du nombre  $2^{64} + 1$ . *C. R. Acad. Sci. Paris* 91 (1880), 138.
- PERRON, O. *Die Lehre von den Kettenbrüchen*. Volume 1, 3<sup>rd</sup> edition, pp. 6–7, 28–29. B.G. Teubner, Stuttgart, 1954.
- SERRET, J. A. Sur un théorème relatif aux nombres entiers. *J. de mathématiques pures et appliquées* 13 (1848), 12–14.
- WILLIAMS, H. C. How was  $F_6$  factored? *Math. Comp.* 61 (1993), 463–474.

(Reçu le 10 décembre 1999; version révisée reçue le 19 septembre 2000)

Freeman Dyson

The Institute for Advanced Study  
 School of Natural Sciences  
 Princeton, N.J. 08540  
 U. S. A.  
*e-mail*: dyson@ias.edu

**vide-leer-empty**