

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ALMOST COMPLEX STRUCTURES ON 8-MANIFOLDS
Autor: MÜLLER, Stefan / GEIGES, Hansjörg
Kapitel: 2. Stable almost complex structures
DOI: <https://doi.org/10.5169/seals-64796>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 04.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Theorem 4 in the general case is also due to the first author. The present joint note grew from the second author's observation that the original proof can be simplified and be made more geometric by appealing to considerations in [5].

This work was completed during a visit by the second author to the Forschungsinstitut für Mathematik (FIM) of the ETH Zürich. He thanks the FIM for its hospitality and support.

2. STABLE ALMOST COMPLEX STRUCTURES

To derive Theorem 4(a) from Theorem 3(a) we merely have to show that the condition $w_8(M) \in \text{Sq}^2 H^6(M; \mathbf{Z})$ is void in case I and equivalent to $\chi(M) \equiv 0 \pmod{2}$ (i.e. $w_8(M) = 0$) in case II.

Indeed, in case I we find an integral lift u of $w_2(M)$ whose free part is indivisible. Then there is a dual element $u' \in H^6(M; \mathbf{Z})$ such that $uu' = 1 \in H^8(M; \mathbf{Z})$. By the Wu formula it follows that

$$\text{Sq}^2 H^6(M; \mathbf{Z}) = w_2(M) H^6(M; \mathbf{Z}) = \rho_2(u H^6(M; \mathbf{Z})) = H^8(M; \mathbf{Z}_2).$$

In case II, on the other hand, we can lift $w_2(M)$ to a torsion class $u \in H^2(M; \mathbf{Z})$, thus

$$\text{Sq}^2 H^6(M; \mathbf{Z}) = \rho_2(u H^6(M; \mathbf{Z})) = 0.$$

3. THE TOP-DIMENSIONAL OBSTRUCTION

Assume we have an almost complex structure J_0 over M with a disc D^8 removed (which is homotopy equivalent to the 7-skeleton of M). Thinking of an almost complex structure J as a section of the SO_8/U_4 -bundle associated to TM , we may interpret J_0 as such a section defined only over the 7-skeleton of M . The obstruction $\sigma(M, J_0)$ to extending J_0 to an almost complex structure on M then lives in

$$H^8(M; \pi_7(\text{SO}_8/U_4)) \cong \pi_7(\text{SO}_8/U_4) \cong \mathbf{Z} \oplus \mathbf{Z}_2.$$

See [13] for references to the computations of the homotopy group above and others used below. The homotopy group involved here is in fact the