

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 46 (2000)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** HARTREE'S THEOREM ON EXISTENCE OF THE QUANTUM DEFECT

**Autor:** KOOSIS, Paul

#### Bibliographie

**DOI:** <https://doi.org/10.5169/seals-64799>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$\operatorname{Re}(1/\kappa) + m > -1$  and bounded away from  $-1$  as  $\kappa \rightarrow -1/p$ . For the  $\gamma_\kappa$  we can take *circular arcs* lying in the fourth quadrant, *orthogonal to the real axis* and running from  $0$  to  $(1-\kappa)/(1+\kappa)$ ; these  $\gamma_\kappa$  will stay away from  $1$  while  $(1-\kappa)/(1+\kappa) \rightarrow (p+1)/(p-1)$ , i.e., while  $\kappa \rightarrow -1/p$ . Then  $|1-s|$  will be uniformly bounded away from  $0$  for  $s$  on these  $\gamma_\kappa$ , and the *integral* in (102) thus *remain bounded* as  $\kappa \rightarrow -1/p$ . At the same time, however, the *sum* in (102) will tend to  $\infty$  like  $1/((1/\kappa)+p)$ . Hence  $A(\kappa)$  will tend to  $\infty$  as  $\kappa \rightarrow -1/p$  in the manner described.

This being so for *each* of the points  $-1/p$ ,  $p = 2, 3, \dots$ ,  $A(\kappa)$  *can have no analytic continuation from  $\mathcal{D}_+$  into any neighbourhood of  $0$* , since any such continuation would have to coincide with the one just constructed on the intersection of the open second quadrant with the neighbourhood in question. That is what we needed to prove.

Before concluding, I must again thank my friend Victor Havin for having, during a conversation, expressed a thought which, indirectly, got me onto a path leading to the above argument.

## BIBLIOGRAPHY

- [1] SCHIFF, L. I. *Quantum Mechanics*. McGraw-Hill, New York, 1955.
- [2] MARGENAU, H. and G. M. MURPHY. *The Mathematics of Physics and Chemistry*. Van Nostrand, New York, 1943.
- [3] HARTREE, D. R. The wave mechanics of an atom with a non-Coulomb central field. Part III. Term values and intensities in series in optical spectra. *Proc. Camb. Phil. Soc.* 24 (1928), 426–437.
- [4] SEATON, M. J. Quantum defect theory. *Reports Prog. Phys.* 46 (1983), 167–257.
- [5] GALLAGHER, T. F. *Rydberg Atoms*. Cambridge Univ. Press, 1994.
- [6] TITCHMARSH, E. C. *Eigenfunction Expansions*. Oxford, Clarendon Press, 1946.
- [7] HAM, F. S. The quantum defect method. In: *Solid State Physics*, edited by Seitz and Turnbull. Vol. 1 (1955), 127–192. Academic Press, New York.
- [8] MOTT, N. F. and H. S. W. MASSEY. *The Theory of Atomic Collisions*. Oxford, Clarendon Press; 2nd edition, 1949; 3rd edition, 1965.
- [9] FANO, U. and A. R. P. RAU. *Atomic Collisions and Spectra*. Academic Press, Orlando, 1986.
- [10] RAU, A. R. P. and M INOKUTI. The quantum defect: Early history and recent developments. *Amer. J. Phys.* 65:3 (1997), 221–225.
- [11] INCE, E. L. *Ordinary Differential Equations*. Dover, New York, 1956.

- [12] TITCHMARSH, E. C. *Eigenfunction Expansions*. Part II. Oxford, Clarendon Press, 1958.
- [13] SHUBOV, M. A. Asymptotic behavior of the quantum defect as a function of an angular momentum and almost normalization of eigenfunctions for a three-dimensional Schrödinger operator with nearly Coulomb potential. *Differential Integral Equations* 8: 8 (1995), 1885–1910.

(Reçu le 25 octobre 1999)

Paul Koosis

Department of Mathematics & Statistics  
McGill University  
Montreal  
Canada

**vide-leer-emptv**