

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PROJECTIVE GEOMETRY OF POLYGONS AND DISCRETE 4-
VERTEX AND 6-VERTEX THEOREMS
Autor: Ovsienko, V / Tabachnikov, S.
Kapitel: 2.1 DISCRETE 4-VERTEX THEOREM
DOI: <https://doi.org/10.5169/seals-65426>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. THEOREMS ON PLANE POLYGONS

In this section we formulate our results for plane polygonal curves. The proofs will be given in Section 4.1.

2.1 DISCRETE 4-VERTEX THEOREM

The *osculating circle* of a smooth plane curve at a point is the circle (or straight line) that has 3rd order of contact with the curve at the given point. One may say that the osculating circle goes through 3 infinitely close points; at a vertex the osculating circle passes through 4 infinitely close points. Moreover, a generic curve crosses the osculating circle at a generic point and stays on one side of it at a vertex. This well-known fact motivates the following definition.

Let P be a plane convex n -gon; throughout this section we assume that $n \geq 4$. Denote the consecutive vertices by V_1, \dots, V_n ; the subscripts are understood cyclically, that is, $V_{n+1} = V_1$, etc.

DEFINITION 2.1. A triple of vertices (V_i, V_{i+1}, V_{i+2}) is said to be *extremal*¹⁾ if V_{i-1} and V_{i+3} lie on the same side of the circle through V_i, V_{i+1}, V_{i+2} (this does not exclude the case where V_{i-1} or V_{i+3} belongs to the circle).




FIGURE 1

The next result follows from a somewhat more general theorem due to O. Musin and V. Sedykh [12] (see also [13]).

¹⁾ We have a terminological difficulty here: as we are dealing with polygons, we cannot use the term “vertex” in the same sense as in the smooth case; hence the term “extremal”.

THEOREM 2.2. *Every plane convex polygon P has at least 4 extremal triples of vertices.*

EXAMPLE 2.3. If P is a quadrilateral then the theorem holds tautologically since the $(i - 1)^{\text{st}}$ vertex coincides with the $(i + 3)^{\text{rd}}$ for every i .

REMARK 2.4. An alternative approach to discretization of the 4-vertex theorem consists in inscribing circles in consecutive triples of sides of a polygon (the centre of such a circle is the intersection point of the bisectors of consecutive angles of the polygon). Then a triple of sides (l_i, l_{i+1}, l_{i+2}) is said to be *extremal* if the lines l_{i-1}, l_{i+3} either both intersect the corresponding circle or both fail to intersect it. With this definition an analogue of Theorem 2.2 holds true [19, 16], and this, in the limit, also provides the smooth 4-vertex theorem.

Both formulations, concerning circumscribed or inscribed circles, make sense on the sphere. Moreover, they are equivalent via projective duality.

2.2 DISCRETE THEOREM ON 6 AFFINE VERTICES

Five generic points in the plane determine a conic. Considering the plane as an affine part of the projective plane, the complement of the conic has two connected components. Let P be a plane convex n -gon; throughout this section we assume that $n \geq 6$. As in the previous section, we introduce the following definition.

DEFINITION 2.5. Five consecutive vertices V_i, \dots, V_{i+4} are said to be *extremal* if V_{i-1} and V_{i+5} lie on the same side of the conic through these 5 points (this does not exclude the case where V_{i-1} or V_{i+5} belongs to the conic).

If P is replaced by a smooth convex curve, and V_i, \dots, V_{i+4} are infinitely close points, we recover the definition of an affine vertex. Hence the following theorem is a discrete version of the smooth theorem on 6 affine vertices.

THEOREM 2.6. *Every plane convex polygon P has at least 6 extremal quintuples of vertices.*

EXAMPLE 2.7. If P is a hexagon then the theorem holds tautologically for the same reason as in Example 2.3.