Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 47 (2001)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PROJECTIVE GEOMETRY OF POLYGONS AND DISCRETE 4-

VERTEX AND 6-VERTEX THEOREMS

Autor: Ovsienko, V / Tabachnikov, S. Kapitel: 3.1 Intersection multiplicities

DOI: https://doi.org/10.5169/seals-65426

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 30.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. MAIN THEOREM

All theorems from Section 2 are consequences of one theorem on the least number of flattenings of a closed polygon in real projective space.

In his remarkable work [3], M. Barner introduced the notion of a *strictly convex* curve in real projective space: this is a smooth closed curve $\gamma \subset \mathbf{RP}^d$ such that for every (d-1)-tuple of points on γ there exists a hyperplane through these points that does not intersect γ at any other points. Barner discovered the following theorem:

A strictly convex curve has at least d+1 distinct flattening points.

Recall that a flattening point of a projective space curve is a point at which the osculating hyperplane is stationary; in other words, this is a singularity of the projectively dual curve. In fact, Barner's result is considerably stronger but we shall not dwell on it here – see [15] for an exposition.

Our goal in this section is to provide a discrete version of Barner's theorem. First we need to develop an elementary intersection formalism for polygonal lines.

3.1 Intersection multiplicities

Throughout this section we shall look at closed polygons $P \subset \mathbf{RP}^d$ with vertices V_1, \ldots, V_n $(n \ge d+1)$ in general position. In other words, for every set of vertices V_{i_1}, \ldots, V_{i_k} , where $k \le d+1$, the span of V_{i_1}, \ldots, V_{i_k} is (k-1)-dimensional.


DEFINITION 3.1. A polygon P is said to be *transverse* to a hyperplane H at a point $X \in P \cap H$ if

- (a) X is an interior point of an edge and this edge is transverse to H, or
- (b) X is a vertex, the two edges incident to X are transverse to H and are locally separated by H.

Clearly, transversality is an open condition.

DEFINITION 3.2. A polygon P is said to intersect a hyperplane H with multiplicity k if for every hyperplane H' sufficiently close to H and transverse to P, the number of points $P \cap H'$ does not exceed k and, moreover, k is attained for some H'.

This definition does not exclude the case where a number of vertices of P lie in H.



LEMMA 3.3. Let V_{i_1}, \ldots, V_{i_k} with $k \leq d$ be vertices of P. Then any hyperplane H passing through V_{i_1}, \ldots, V_{i_k} meets P with multiplicity at least k.

Proof. Move each V_{i_j} (j = 1, ..., k) slightly along the edge (V_{i_j}, V_{i_j+1}) to obtain a new point V'_{i_j} . Let us show that a generic hyperplane H' through $V'_{i_1}, ..., V'_{i_k}$ is transverse to P. This will imply the lemma because H' has at least k intersections with P.

It suffices to show that H' does not contain any vertex of P. First we note that, since P is in general position, a generic hyperplane H through V_{i_1}, \ldots, V_{i_k} does not contain any other vertex. The same holds true for every hyperplane which is sufficiently close to H. It remains to show that the chosen H' does not contain any of V_{i_1}, \ldots, V_{i_k} .

Suppose H' contains V_{i_j} . Then H' contains the edge (V_{i_j}, V_{i_j+1}) and therefore also V_{i_j+1} . If $i_j+1 \notin \{i_1,\ldots,i_k\}$ we obtain a contradiction with the previous paragraph. If, on the other hand, $i_j+1 \in \{i_1,\ldots,i_k\}$ then we can proceed in the same way with V_{i_j+1} . However, we cannot go on indefinitely since k < n. \square

The next definition is topological in nature.

DEFINITION 3.4. Consider a continuous curve in \mathbb{RP}^d with endpoints A and Z. Let H be a hyperplane not containing A or Z. We say that A and Z are on one side of H if one can connect A and Z by a curve not intersecting H in such a way that the resulting closed curve is contractible. Otherwise we say that A and Z are separated by H.

Clearly, if one has only two points A and Z (and no curve connecting

them), then one cannot say that the points are on one side of, or separated by, a hyperplane.

LEMMA 3.5. Let $\Gamma = (A, ..., Z)$ be a broken line in general position in \mathbf{RP}^d , and let H be a hyperplane not containing A or Z. Denote by k the intersection multiplicity of Γ with H. Then A and Z are separated by H if k is odd and not separated otherwise.

Proof. Connect Z and A by a segment so as to obtain a closed polygon $\overline{\Gamma}$ and consider a hyperplane H' close to H, transverse to $\overline{\Gamma}$ and intersecting Γ in k points. Since $\overline{\Gamma}$ is contractible, H' intersects $\overline{\Gamma}$ in an even number of points. Therefore, H' intersects the segment (Z,A) for odd k and does not intersect it for even k. \square


The next definition introduces a significant class of polygons which is our main object of study.

DEFINITION 3.6. A polygon P is called *strictly convex* if through every d-1 vertices there passes a hyperplane H whose intersection multiplicity with P is equal to d-1.

This definition becomes, in the smooth limit, that of strict convexity for smooth curves, due to Barner.

DEFINITION 3.7. A *d*-tuple of consecutive vertices (V_i, \ldots, V_{i+d-1}) of a polygon P in \mathbb{RP}^d is called a *flattening* if the endpoints V_{i-1} and V_{i+d} of the broken line $(V_{i-1}, \ldots, V_{i+d})$ are:

- (a) separated by the hyperplane through (V_i, \ldots, V_{i+d-1}) if d is even,
- (b) not separated if d is odd.



a) d = 2

b) d = 3

REMARK 3.8. A curve in \mathbf{RP}^d can be lifted to $\mathbf{R}^{d+1} \setminus \{0\}$; the lifting is not unique. Given a polygon $P \subset \mathbf{RP}^d$ with vertices V_1, \ldots, V_n , we lift it to \mathbf{R}^{d+1} as a polygon \widetilde{P} and denote its vertices by $\widetilde{V}_1, \ldots, \widetilde{V}_n$. Then a d-tuple (V_i, \ldots, V_{i+d-1}) is a flattening if and only if the determinant

$$\Delta_j = \left| \widetilde{V}_j \dots \widetilde{V}_{j+d} \right|$$

changes sign as j varies from i-1 to i.

This property is independent of the lifting.

3.2 A SIMPLEX IS STRICTLY CONVEX

Define a simplex $S_d \subset \mathbf{RP}^d$ with vertices V_1, \dots, V_{d+1} as the projection from the punctured \mathbf{R}^{d+1} of the polygonal line:

(3.2)
$$\widetilde{V}_1 = (1, 0, \dots, 0), \quad \widetilde{V}_2 = (0, 1, 0, \dots, 0), \quad \dots, \widetilde{V}_{d+1} = (0, \dots, 0, 1)$$
 and

$$(3.3) \widetilde{V}_{d+2} = (-1)^{d+1} \widetilde{V}_1.$$

The last vertex has the same projection as the first one; S_d is contractible for odd d, and non-contractible for even d.




FIGURE 5

PROPOSITION 3.9. The polygon S_d is strictly convex.

Proof. We need to prove that through every (d-1)-tuple

$$(V_1,\ldots,\widehat{V}_i,\ldots,\widehat{V}_i,\ldots,V_{d+1})$$

there passes a hyperplane H intersecting P with multiplicity d-1. Select a point W on the line $(\widetilde{V}_i,\widetilde{V}_j)$ in such a manner that W lies on the segment $(\widetilde{V}_i,\widetilde{V}_j)$ if j-i is even, and does not lie on it if j-i is odd. Define \widetilde{H} as the linear span of $\widetilde{V}_1,\ldots,\widehat{\widetilde{V}}_i,\ldots,\widehat{\widetilde{V}}_j,\ldots,\widetilde{V}_{d+1},W$. We claim that its projection $H\subset\mathbf{RP}^d$ meets S_d with multiplicity $\leq d-1$.