

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVERTING RADON TRANSFORMS : THE GROUP-THEORETIC APPROACH
Autor: Rouvière, François
Kapitel: 2.1 Double fibrations of homogeneous spaces
DOI: <https://doi.org/10.5169/seals-65436>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We briefly recall some classical semisimple notations, as used for instance in Helgason's books. Let \mathfrak{a} be a maximal abelian subspace of \mathfrak{p} . Related to the restricted root system of the pair $(\mathfrak{g}, \mathfrak{a})$ are the eigenspaces \mathfrak{g}_α , the *Iwasawa decomposition* $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}$ of the Lie algebra and $G = KAN$ for the group (unique decomposition of each element of G into a product of factors in the respective subgroups); the subgroups A , resp. N , of G are abelian, resp. nilpotent. The half sum of positive roots (counted with multiplicities) is a linear form ρ on \mathfrak{a} ; we write $a^\rho = e^{\rho(\log a)}$ for $a \in A$. Let M , resp. M' , denote the centralizer, resp. normalizer, of A in K . Then $W = M'/M$ is a finite group called the Weyl group.

Let y_o denote the orbit $N \cdot x_o \subset X$. The *horocycles* of X are the submanifolds $g \cdot y_o$, for $g \in G$. Since $g \cdot y_o = y_o$ (globally) if and only if $g \in MN$, the space of all horocycles is $Y = G/MN$.

e. ISOTROPIC RIEMANNIAN SYMMETRIC SPACES. A Riemannian manifold X is called *isotropic* if, for every $x \in X$ and every pair of unit tangent vectors V, W to X at x , there exists an isometry of X leaving x fixed and mapping V to W . The connected isotropic Riemannian manifolds are the Euclidean spaces \mathbf{R}^n , the *hyperbolic spaces* i.e. the Riemannian symmetric spaces of the noncompact type and of rank one ($\dim \mathfrak{a} = 1$), and their compact analogues, spheres and projective spaces. The compact spaces will not be considered in this paper, so that most of our examples will be taken from the list

$$\mathbf{R}^n, H^n(\mathbf{R}), H^n(\mathbf{C}), H^n(\mathbf{H}), H^{16}(\mathbf{O}).$$

Among them we shall often restrict ourselves to the *classical hyperbolic spaces* $H^n(\mathbf{F})$, with $\mathbf{F} = \mathbf{R}, \mathbf{C}$ or \mathbf{H} .

2. GEOMETRIC SETTING

2.1 DOUBLE FIBRATIONS OF HOMOGENEOUS SPACES

The general group-theoretic setting for Radon transforms, introduced by Helgason in the sixties, is motivated by the well-known example of points and hyperplanes in the Euclidean space \mathbf{R}^n . The set of points and the set of hyperplanes are both homogeneous spaces of the isometry group of \mathbf{R}^n , and it turns out that the fundamental "incidence" relation (a point x belongs to a hyperplane y), as well as the defining integral of the Radon transform, have simple expressions in terms of Lie groups and invariant measures. This observation suggests considering the following general situation.

Let X and Y be two manifolds, with given origins $x_o \in X$ and $y_o \in Y$, and assume a real Lie group G acts transitively on both manifolds X and Y . Two elements $x \in X$ and $y \in Y$ are said to be *incident* if there exists some $g \in G$ such that $x = g \cdot x_o$ and $y = g \cdot y_o$. Roughly speaking, if we think of g as a motion, this means that x and y have the same relative position as the origins x_o and y_o .

A more convenient formulation is obtained in terms of the isotropy subgroups K , resp. H , of x_o , resp. y_o , in G . They are closed Lie subgroups of G , and the manifolds X, Y can be identified with the homogeneous spaces of left cosets $G/K, G/H$ respectively; in particular we may write $x_o = K, y_o = H, g \cdot x_o = gK$, etc. The points $x = g'K \in X$ and $y = g''H \in Y$ are then incident if and only if there exists $g \in G$ such that $g'K = g \cdot x_o = gK$ and $g''H = g \cdot y_o = gH$, in other words if the left cosets $g'K$ and $g''H$, as subsets of G , are not disjoint (they meet at g).

Given $y = g''H$, we see that x is incident to y if and only if $x = g'hK$ for some $h \in H$. Given $x = g'K$, the point y is incident to x if and only if $y = g'kH$ for some $k \in K$.

In the above example X , resp. Y , is the set of points, resp. hyperplanes, of \mathbf{R}^n and G is the group of all isometries. But hyperplanes can also be viewed as subsets of $X = \mathbf{R}^n$, and the incidence relation boils down to the familiar "the point x belongs to the hyperplane y " if and only if the chosen origin x_o belongs to the chosen origin y_o . Lemma 1 below extends this fact to Riemannian manifolds. More general incidence relations can be considered, however, and will be helpful in Section 6.

Clearly, the group G acts transitively on the subset Z of $X \times Y$ consisting of all incident couples $(x, y) = (g \cdot x_o, g \cdot y_o)$, with $K \cap H$ as the isotropy subgroup of the origin $(x_o, y_o) \in Z$. Thus $Z = G/(K \cap H)$ can be endowed with a structure of manifold, and the present setting can be summarized by the following *double fibration of homogeneous spaces*

$$\begin{array}{ccc} Z = G/(K \cap H) & \subset & X \times Y \\ \downarrow & & \searrow \\ X = G/K & & Y = G/H, \end{array}$$

where the arrows denote the natural projections.

Radon transforms can be studied with more general double fibrations of manifolds X, Y, Z (without groups), as introduced by Gel'fand et al. [4]. We refer to Guillemin and Sternberg ([6], p.340, 370) for their basic properties; this theory has been developed in several papers by Boman, Quinto, and others.