

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	47 (2001)
<b>Heft:</b>	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
<b>Artikel:</b>	INVERTING RADON TRANSFORMS : THE GROUP-THEORETIC APPROACH
<b>Autor:</b>	Rouvière, François
<b>Kapitel:</b>	2.2 Group-theoretic Radon transforms
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-65436">https://doi.org/10.5169/seals-65436</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 2.2 GROUP-THEORETIC RADON TRANSFORMS

Let  $G$  be a real Lie group and  $K$  a (closed) Lie subgroup, equipped with left-invariant Haar measures  $dg$ ,  $dk$  respectively. If the homogeneous space  $G/K$  admits a  $G$ -invariant measure  $d(gK)$ , the measures can then be normalized so that

$$\int_G f(g) dg = \int_{G/K} d(gK) \int_K f(gk) dk,$$

for any  $f \in C_c(G)$ . This applies in particular if  $K$  is compact (on invariant measures, see [9], Chap. I, § 1).

Throughout the paper  $G$  will be a Lie group,  $K$  a *compact* subgroup, and  $H$  a (closed) Lie subgroup of  $G$ . The Haar measure  $dk$  of  $K$  will be normalized by  $\int_K dk = 1$ .

Let  $u$  be a (complex-valued) function on  $X = G/K$ . Its *Radon transform* is the function  $Ru$  on  $Y = G/H$  defined by

$$Ru(gH) = \int_H u(ghK) dh,$$

for  $g \in G$ , whenever this makes sense (e.g. if  $u \in C_c(X)$ ). The left invariance of  $dh$  implies that the integral only depends on the left coset  $gH$  of  $g$ . Given  $y = gH$  in  $Y = G/H$ , the value  $Ru(y)$  is an integral of  $u$  over all  $x$  incident to  $y$ . A more precise statement can be given in the following important example.

**EXAMPLE.** Let  $X$  be a connected Riemannian manifold,  $G$  a transitive Lie group of *isometries* of  $X$  and  $K$  the isotropy subgroup of some origin  $x_o \in X$ ; then  $K$  is compact ([8], p. 204) and  $X = G/K$ . Let  $y_o$  be a given *closed submanifold* of  $X$ , containing  $x_o$ , and let  $Y$  be the set of all submanifolds  $y = g \cdot y_o$  of  $X$ , with  $g \in G$ .

The set  $H$  of all  $h \in G$  such that  $h \cdot y_o = y_o$  (i.e. the submanifold  $y_o$  is globally invariant under  $h$ ) is a closed Lie subgroup of  $G$ . Indeed if  $h_n \in H$  converges to  $h$  in  $G$ , for any  $x \in y_o$  the point  $\lim h_n \cdot x = h \cdot x$  belongs to  $y_o$ ; similarly  $h^{-1} \cdot x \in y_o$ , so that  $h \cdot y_o = y_o$ . Thus  $Y = G/H$  can be endowed with a structure of manifold and we obtain a double fibration of homogeneous spaces.

The following lemma allows one to compute the Radon transform without knowing  $H$  explicitly.

LEMMA 1. *Keeping the notation of this example, assume furthermore that  $y_o = G' \cdot x_o$  is a closed orbit of the origin  $x_o = K$  under some Lie subgroup  $G'$  of  $G$ .*

*Then  $G' \subset H \subset G'K$  and  $y_o = H \cdot x_o$ . The incidence relation between  $X = G/K$  and  $Y = G/H$  is simply  $x \in y$ . Besides, the left-invariant Haar measures  $dh, dg'$  of the groups  $H, G'$  can be normalized so that*

$$\begin{aligned} Ru(y) &= \int_H u(gh \cdot x_o) dh = \int_{G'} u(gg' \cdot x_o) dg' \\ &= \int_y u(x) dm_y(x), \end{aligned}$$

where  $dm_y$  is the Riemannian measure induced by  $X$  on its submanifold  $y = g \cdot y_o$ .

REMARK. The subgroup  $H$  can of course be strictly bigger than  $G'$ . This occurs for instance if  $y_o$  is a line in  $X = \mathbf{R}^n$  and  $G'$  is the group of translations along this line, or a horocycle in a Riemannian symmetric space  $X$  of the noncompact type (for which  $G' = N$  and  $H = MN = NM$  in the usual semisimple notations).

*Proof of Lemma 1.* If  $y_o = G' \cdot x_o$ , then  $H$  obviously contains  $G'$  and it follows that

$$y_o = G' \cdot x_o \subset H \cdot x_o \subset y_o,$$

whence  $H \cdot x_o = G' \cdot x_o$  and  $H \subset G'K$ .

A point  $x \in X$  is incident to  $y = g \cdot y_o \in Y$  if and only if there exists  $h \in H$  such that  $x = gh \cdot x_o$ , i.e.  $x \in gH \cdot x_o = g \cdot y_o = y$ .

An isometry  $g$  transforms the Riemannian measure of  $y_o$  into the Riemannian measure of  $y = g \cdot y_o$ , and it suffices to prove the integral formula for  $g = e$ . Now  $y_o = H \cdot x_o$  can be identified to the homogeneous space  $H/(H \cap K)$ , and  $dm_{y_o}$  (which is invariant under all isometries of  $y_o$ ) to an  $H$ -invariant measure on this space. The Haar measure  $dh$  can therefore be normalized so that the corresponding measure on  $H/(H \cap K)$  satisfies

$$\begin{aligned} \int_{y_o} u(x) dm_{y_o}(x) &= \int_{H/(H \cap K)} u(h \cdot x_o) d(h(H \cap K)) \\ &= \int_H u(h \cdot x_o) dh = Ru(y_o). \end{aligned}$$

The proof is similar for  $\int_{G'}$ , whence the lemma.  $\square$

Going back to general double fibrations, the *Radon dual transform* of a (continuous, say) function  $v$  on  $Y = G/H$  is the function on  $X = G/K$  defined by

$$R^*v(gK) = \int_K v(gkH) dk,$$

for  $g \in G$ , an integral of  $v$  over all  $y$  incident to  $x = gK$ . The word “dual” is of course motivated by the classical projective duality between points and hyperplanes in the basic example, but it also stems from the following proposition.

**PROPOSITION 2.** *Let  $X = G/K$  with  $K$  compact, and assume that  $Y = G/H$  has a  $G$ -invariant measure. Let  $u \in C_c(X)$ ,  $v \in C(Y)$ . Then  $Ru \in C_c(Y)$ ,  $R^*v \in C(X)$  and*

$$\int_X u(x) R^*v(x) dx = \int_Y Ru(y) v(y) dy = \int_Z u(x) v(y) dz,$$

where  $dx, dy, dz$  are the respective  $G$ -invariant measures on  $X, Y$  and  $Z = G/(K \cap H)$ .

In the latter integral  $u(x)v(y)$  is considered as a function of  $z = (x, y)$  on  $Z$  (Section 2.1). We omit the proof, a classical exercise on invariant integrals (cf. [9], p. 144 and [11], p. 41); all groups are assumed unimodular there, but the proof only uses the invariant measures on the homogeneous spaces, thus extends to the present situation.

Proposition 2 allows a natural extension of the transforms  $R$  and  $R^*$  to distributions. Given  $u \in \mathcal{E}'(X)$ , the distribution  $Ru \in \mathcal{E}'(Y)$  is defined by

$$\langle Ru, v \rangle = \langle u, R^*v \rangle,$$

for all test functions  $v \in C^\infty(Y)$ . Similarly, given  $v \in \mathcal{D}'(Y)$ , the distribution  $R^*v \in \mathcal{D}'(X)$  is defined by

$$\langle R^*v, u \rangle = \langle v, Ru \rangle,$$

for all  $u \in \mathcal{D}(X)$ . Again we refer to Helgason ([11], p. 42) for details, based on the compactness of  $K$ . These definitions do extend the Radon integrals for functions, as Proposition 2 shows, when identifying a function  $u$  with the distribution  $u(x) dx$ , and similarly for  $v$ .