Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 47 (2001)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVERTING RADON TRANSFORMS : THE GROUP-THEORETIC APPROACH

Autor: Rouvière, François

Kapitel: 3.2 Radon inversion by convolution

DOI: https://doi.org/10.5169/seals-65436

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 17.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$\int_N u(n\cdot x_o)\,dn = 2^{p-2+(q/2)}\omega_p\omega_q \int_0^\infty \int_0^\infty u(\exp(rH)\cdot x_o)\,x^{(p/2)-1}y^{(q/2)-1}dx\,dy$$
$$= \int_0^\infty u(\exp(rH)\cdot x_o)f(r)\,dr\,.$$

The latter expression follows from the change of variables $(x,r)\mapsto(x,y)$, with Jacobian $\sinh 2r$; here

$$f(r) = 2^{p-2+(q/2)}\omega_p\omega_q \sinh 2r \int_0^{\cosh r-1} x^{(p/2)-1}\left(\cosh^2 r-(1+x)^2\right)^{(q/2)-1}dx\,.$$

Setting $x = t(\cosh r-1)$ we find

$$f(r) = 2^{(3p+q)/2}\omega_{n-1}\Big(\sinh r\Big)^{q-1}\Big(\sinh\frac{r}{2}\Big)^p\cosh r$$
$$\times\frac{\Gamma((p+q)/2)}{\Gamma(p/2)\Gamma(q/2)}\int_0^1 t^{(p/2)-1}(1-t)^{(q/2)-1}\Big(1+t\tanh^2\frac{r}{2}\Big)^{(q/2)-1}dt$$
$$= 2^{(3p+q)/2}\omega_{n-1}\Big(\sinh r\Big)^{q-1}\Big(\sinh\frac{r}{2}\Big)^p\cosh r$$
$$\times\,{}_2F_1\Big(\frac{p}{2},1-\frac{q}{2};\frac{p+q}{2};-\tanh^2\frac{r}{2}\Big)\,,$$

by Euler's integral formula for the hypergeometric function. From a quadratic transformation formula for ${}_2F_1$ ([3], p. 113, formula (35)) we finally obtain

$$f(r) = 2^{(n-1)/2}\omega_{n-1}(\sinh r)^{n-2}(\cosh r)^q\,{}_2F_1\left(\frac{\rho-1}{2},\frac{\rho}{2};\frac{n-1}{2};-\sinh^2 r\right).$$

Thus, for K-invariant u,

$$\int_N u(n\cdot x_o)\,dn = \int_0^\infty u\,(\exp(rH)\cdot x_o)\;S(r)A(r)\,dr = \int_X u(x)\,S(x)\,dx\,,$$

where $A(r)=\omega_n(\sinh r)^{n-1}(\cosh r)^q$ and $S(r)=f(r)/A(r)$. □

3.2 Radon inversion by convolution

Radon inversion formulas will follow from Section 3.1 if we can solve for u the convolution equation $u*S=R^*Ru$, in the form

$$u = DR^*Ru\,. \tag{2}$$

To recover $u(x)$ from Ru the recipe will be to integrate $Ru(y)$ over all y incident to x, and to apply the operator D on the x variable.

As noted in the proof of Proposition 3, R^*R commutes with the action of G on X, and it is natural to look for a D with the same property, i.e. a convolution operator: if T is a distribution on X such that $S*T=\delta$, then

$$u = (R^* Ru) * T .$$

Though the question can be tackled by harmonic analysis on X (cf. Section 5), a G-invariant linear differential operator D can sometimes be found directly, such that $DS = \delta$. Then (2) follows from the equality $u = u * DS = D(u * S)$. Indeed, for any test function φ,

$$\begin{aligned} \langle D(u * S), \varphi \rangle &= \langle u * S, {}^t D\varphi \rangle \\ &= \big\langle u(g \cdot x_o), \langle S, ({}^t D\varphi) \circ \tau(g) \rangle \big\rangle \qquad \text{by (1)} \\ &= \big\langle u(g \cdot x_o), \langle S, {}^t D(\varphi \circ \tau(g)) \rangle \big\rangle , \end{aligned}$$

since the transpose operator ${}^t D$ is G-invariant too, as follows from the existence of a G-invariant measure on X. Finally,

$$\begin{aligned} \langle D(u * S), \varphi \rangle &= \big\langle u(g \cdot x_o), \langle DS, \varphi \circ \tau(g) \rangle \big\rangle \\ &= \langle u * DS, \varphi \rangle , \end{aligned}$$

as claimed; assuming G unimodular (as in [9], p. 291) is thus unnecessary here.

The method applies whenever we can *find a G-invariant differential operator D on X with given fundamental solution S*. We shall now investigate this question on the basis of Propositions 4 and 5.

4. Radon transforms on isotropic spaces

Throughout this section X will be an *isotropic* connected *noncompact* Riemannian manifold, that is a Euclidean space or a Riemannian globally symmetric space of rank one:

$$X = \mathbf{R}^n \text{ or } H^m(\mathbf{R}),\ H^{2m}(\mathbf{C}),\ H^{4m}(\mathbf{H}),\ H^{16}(\mathbf{O}) ,$$

where all superscripts denote the real dimension of these real, complex, quaternionic or Cayley hyperbolic spaces (cf. Wolf [18], §8.12). We first try to invert the d-geodesic Radon transform on X, defined by integrating over a family of d-dimensional totally geodesic submanifolds of X. At the end of this section we shall see that the same tools provide an inversion formula for the horocycle Radon transform on $H^{2k+1}(\mathbf{R})$.

4.1 Totally geodesic submanifolds

Our first goal is to describe these submanifolds and the corresponding functions S in Proposition 4.