

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FINITE TYPE LINK-HOMOTOPY INVARIANTS
Autor: Lin, Xiao-Song
Kapitel: 3. Construction of the invariant
DOI: <https://doi.org/10.5169/seals-65440>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

This bijection is an anti-homomorphism: $\overline{\sigma_1\sigma_2} = \overline{\sigma_2}\overline{\sigma_1}$. This bijection induces an operation on $\mathbf{Z}^{(k)}_{(3)}$.

THEOREM 2.9. *The operation on $\mathbf{Z}^{(k)}_{(3)}$ induced by reversing the orientation of each component of a string link is to change each $\mu(rst)$ to $-\mu(rst)$ followed by the translation operation*

$$\mu(rst) \longrightarrow \mu(rst) - l_{rs}l_{rt} + l_{rs}l_{st} - l_{rt}l_{st}.$$

Proof. Consider the normal form (2) of $\sigma \in \mathcal{H}(k)/\mathcal{H}(k)_3$ in the r, s, t -th components. The normal form for $\overline{\sigma}$ is obtained as follows:

$$\begin{aligned} \overline{\sigma} &= [\tau_{rt}, \tau_{st}]^{-\delta} \tau_{st}^{\gamma} \tau_{rt}^{\beta} \tau_{rs}^{\alpha} \\ &= \tau_{rs}^{\alpha} \tau_{rt}^{\beta} \tau_{st}^{\gamma} [\tau_{rt}, \tau_{st}]^{-\delta - \alpha\beta + \alpha\gamma - \beta\gamma}. \end{aligned}$$

Thus the operation on $\mathbf{Z}^{(k)}_{(3)}$ induced by $\sigma \mapsto \overline{\sigma}$ is given by

$$\mu(rst) \longrightarrow -\mu(rst) - l_{rs}l_{rt} + l_{rs}l_{st} - l_{rt}l_{st}. \quad \square$$

3. CONSTRUCTION OF THE INVARIANT

By Theorems 2.2 and 2.7, we shall look for polynomials in l_{ij} and $\mu(rst)$ invariant under the translation operations on $\{\mu(rst)\} \in \mathbf{Z}^{(k)}_{(3)}$ induced by partial conjugations. There are $k(k-1)$ partial conjugations altogether and their induced translations subject to $2k$ linear equations given in Theorem 2.8. If these equations are linearly independent for generic values of $\{l_{ij}\}$, the sublattice of $\mathbf{Z}^{(k)}_{(3)}$ generated by the translation vectors of the partial conjugations will be of dimension no larger than $k(k-1) - 2k = k^2 - 3k$.

LEMMA 3.1. *For $k > 3$, the $2k$ equations in Theorem 2.8 are linearly independent for generic values of $\{l_{ij}\}$.*

Proof. We write the two sets of equations in Theorem 2.8 as follows:

$$\begin{aligned} \mathbf{1}^i + \mathbf{2}^i + \cdots + \mathbf{j}^i + \cdots + \mathbf{k}^i &= 0, \quad j \neq i; \\ l_{i1}\mathbf{i}^1 + l_{i2}\mathbf{i}^2 + \cdots + l_{ij}\mathbf{i}^j + \cdots + l_{ik}\mathbf{i}^k &= 0, \quad j \neq i, \end{aligned}$$

for each $i = 1, 2, \dots, k$.

For generic values of $\{l_{ij}\}$, using the first $k-1$ equations from the first set of k equations, we can solve for $\mathbf{k}^1, \mathbf{k}^2, \dots, \mathbf{k}^{k-1}$. Similarly, we can solve

for $\mathbf{1}^k, \mathbf{2}^k, \dots, (\mathbf{k} - \mathbf{1})^k$ from the first $k - 1$ equations of the second set of k equations. The remaining vectors \mathbf{i}^j , $i, j \neq k$, have to satisfy another two equations obtained from the last equations in those two sets of k equations, respectively, by substituting \mathbf{k}^i and \mathbf{i}^k with their solutions in terms of \mathbf{i}^j for $i, j \neq k$. It is then easy to check that these two equations are linearly independent when $k > 3$. \square

LEMMA 3.2. For $k = 4, 5$, we have $\binom{k}{3} = k^2 - 3k$. For $k \geq 6$, we have $\binom{k}{3} > k^2 - 3k$.

Proof. We have

$$\binom{k}{3} - (k^2 - 3k) = \frac{k}{6}(k^2 - 9k + 20) = \frac{k}{6}(k - 4)(k - 5). \quad \square$$

THEOREM 3.3. For $k \geq 6$, there exists a polynomial in l_{ij} and $\mu(rst)$ which is a link-homotopy invariant of ordered, oriented links with k components. This link-homotopy invariant is of finite type.

Proof. In $\mathbf{Z}^{\binom{k}{3}}$, let \mathcal{P} be the sublattice generated by the translation vectors of partial conjugations. Then we have

$$\dim(\mathcal{P}) \leq k^2 - 3k < \binom{k}{3}.$$

Let $\Omega \in \mathbf{Z}^{\binom{k}{3}}$ be a non-zero vector perpendicular to \mathcal{P} . We can choose such an Ω so that its coordinates are polynomials in $\{l_{ij}\}$ and the inner product $\mathbf{i}^j \cdot \Omega$ is identically zero. This can be achieved by considering generic values of $\{l_{ij}\}$ first and solving a system of homogeneous equations (with more equations than unknowns) whose coefficients are polynomials in l_{ij} ³). Then since $\mathbf{i}^j \cdot \Omega = 0$ for generic values of $\{l_{ij}\}$, it has to be zero identically. Let $\mu = \{\mu(rst)\} \in \mathbf{Z}^{\binom{k}{3}}$. The inner product $\mu \cdot \Omega$ is invariant under the translations by vectors in \mathcal{P} . This is a desired link-homotopy invariant of ordered, oriented links since

$$(\mu + \mathbf{i}^j) \cdot \Omega = \mu \cdot \Omega$$

for all $i, j = 1, 2, \dots, k$.

The fact that the invariant $\mu \cdot \Omega$ is of finite type is a direct consequence of the fact that the linking numbers and the triple linking numbers are all finite

³) This will be made explicit in the example following this proof.

type invariants of string links ([7], [2]). If we have a singular link, we may put it into the form of the closure of a single string link. Since polynomials of finite type invariants are still of finite type, $\mu \cdot \Omega$ vanishes on singular string links with a sufficiently large number of double points. This implies that it is a finite type link invariant. \square

We now consider in some detail the case $k = 6$. Let us order $\mu(rst)$, $1 \leq r < s < t \leq 6$ in lexicographic order. So

$$\mu = (\mu(123), \mu(124), \mu(125), \mu(126), \mu(134), \mu(135), \mu(136), \mu(145), \mu(146), \mu(156), \\ \mu(234), \mu(235), \mu(236), \mu(245), \mu(246), \mu(256), \mu(345), \mu(346), \mu(356), \mu(456)).$$

Then the vectors of the translation operations $\mathbf{1}^2, \mathbf{1}^3, \mathbf{1}^4, \mathbf{1}^5, \mathbf{1}^6, \mathbf{2}^1, \mathbf{2}^3, \mathbf{2}^4, \mathbf{2}^5, \mathbf{2}^6, \mathbf{3}^1, \mathbf{3}^2, \mathbf{3}^4, \mathbf{3}^5, \mathbf{3}^6, \mathbf{4}^1, \mathbf{4}^2, \mathbf{4}^3, \mathbf{4}^5, \mathbf{4}^6, \mathbf{5}^1, \mathbf{5}^2, \mathbf{5}^3, \mathbf{5}^4, \mathbf{5}^6, \mathbf{6}^1, \mathbf{6}^2, \mathbf{6}^3, \mathbf{6}^4, \mathbf{6}^5$ are the row vectors of the following 30×20 matrix, from top to bottom respectively:

l_{13}	l_{14}	l_{15}	l_{16}	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$-l_{12}$	0	0	0	l_{14}	l_{15}	l_{16}	0	0	0	0	0	0	0	0	0	0	0	0	0
0	$-l_{12}$	0	0	$-l_{13}$	0	0	l_{15}	l_{16}	0	0	0	0	0	0	0	0	0	0	0
0	0	$-l_{12}$	0	0	$-l_{13}$	0	$-l_{14}$	0	l_{16}	0	0	0	0	0	0	0	0	0	0
0	0	0	$-l_{12}$	0	0	$-l_{13}$	0	$-l_{14}$	$-l_{15}$	0	0	0	0	0	0	0	0	0	0
$-l_{23}$	$-l_{24}$	$-l_{25}$	$-l_{26}$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
l_{12}	0	0	0	0	0	0	0	0	0	0	l_{24}	l_{25}	l_{26}	0	0	0	0	0	0
0	l_{12}	0	0	0	0	0	0	0	0	0	$-l_{23}$	0	0	l_{25}	l_{26}	0	0	0	0
0	0	l_{12}	0	0	0	0	0	0	0	0	0	$-l_{23}$	0	$-l_{24}$	0	l_{26}	0	0	0
0	0	0	l_{12}	0	0	0	0	0	0	0	0	0	$-l_{23}$	0	$-l_{24}$	$-l_{25}$	0	0	0
l_{23}	0	0	0	$-l_{34}$	$-l_{35}$	$-l_{36}$	0	0	0	0	0	0	0	0	0	0	0	0	0
$-l_{13}$	0	0	0	0	0	0	0	0	0	0	$-l_{34}$	$-l_{35}$	$-l_{36}$	0	0	0	0	0	0
0	0	0	0	l_{13}	0	0	0	0	0	0	l_{23}	0	0	0	0	0	l_{35}	l_{36}	0
0	0	0	0	0	l_{13}	0	0	0	0	0	l_{23}	0	0	0	0	0	$-l_{34}$	0	l_{36}
0	0	0	0	0	0	l_{13}	0	0	0	0	l_{23}	0	0	0	0	0	$-l_{34}$	$-l_{35}$	0
0	l_{24}	0	0	l_{34}	0	0	$-l_{45}$	$-l_{46}$	0	0	0	0	0	0	0	0	0	0	0
0	$-l_{14}$	0	0	0	0	0	0	0	0	0	l_{34}	0	0	$-l_{45}$	$-l_{46}$	0	0	0	0
0	0	0	0	$-l_{14}$	0	0	0	0	0	0	$-l_{24}$	0	0	0	0	0	$-l_{45}$	$-l_{46}$	0
0	0	0	0	0	0	0	l_{14}	0	0	0	0	0	0	l_{24}	0	0	l_{34}	0	l_{46}
0	0	0	0	0	0	0	0	l_{14}	0	0	0	0	0	l_{24}	0	0	l_{34}	0	$-l_{45}$
0	0	l_{25}	0	0	l_{35}	0	l_{45}	0	$-l_{56}$	0	0	0	0	0	0	0	0	0	0
0	0	$-l_{15}$	0	0	0	0	0	0	0	0	l_{35}	0	l_{45}	0	$-l_{56}$	0	0	0	0
0	0	0	0	0	$-l_{15}$	0	0	0	0	0	$-l_{25}$	0	0	0	0	l_{45}	0	$-l_{56}$	0
0	0	0	0	0	0	0	$-l_{15}$	0	0	0	0	0	$-l_{25}$	0	0	$-l_{35}$	0	0	$-l_{56}$
0	0	0	0	0	0	0	0	0	0	l_{15}	0	0	0	0	0	l_{25}	0	0	l_{35}
0	0	0	l_{26}	0	0	l_{36}	0	l_{46}	l_{56}	0	0	0	0	0	0	0	0	0	0
0	0	0	$-l_{16}$	0	0	0	0	0	0	0	0	l_{36}	0	l_{46}	l_{56}	0	0	0	0
0	0	0	0	0	0	$-l_{16}$	0	0	0	0	0	$-l_{26}$	0	0	0	0	l_{46}	l_{56}	0
0	0	0	0	0	0	0	0	$-l_{16}$	0	0	0	0	0	$-l_{26}$	0	0	$-l_{36}$	0	l_{56}
0	0	0	0	0	0	0	0	0	$-l_{16}$	0	0	0	0	0	0	l_{26}	0	0	$-l_{36}$

We shall pick out the 18 rows of this matrix corresponding to the translation operations of $1^2, 1^3, 1^4, 1^5, 2^1, 2^3, 2^4, 2^5, 3^1, 3^2, 3^4, 3^5, 4^1, 4^2, 4^3, 4^5, 5^1, 5^2$, respectively. Calculation using *Mathematica*[®] shows that these 18 vectors are linearly independent generically.

Consider now the operation of reversing the orientation. The vector $R = \{R(rst)\} \in \mathbf{Z}^{20}$ of the translation operation in Theorem 2.9 is given by

$$R(rst) = -l_{rs} l_{rt} + l_{rs} l_{st} - l_{rt} l_{st}.$$

One can verify that the vector R and the previous 18 vectors are linearly independent. Let \mathcal{M} be the 19×20 matrix formed by these 19 vectors. Let $\mathcal{M}^{(i)}$ be the 19×19 matrix obtained from \mathcal{M} by deleting the i^{th} column from \mathcal{M} , $i = 1, 2, \dots, 20$. Let

$$\Omega_i = (-1)^{i-1} \det(\mathcal{M}^{(i)})$$

and $\Omega = (\Omega_1, \Omega_2, \dots, \Omega_{20})$.

THEOREM 3.4. $\mu \cdot \Omega$ is a finite type link-homotopy invariant of ordered, oriented links with 6 components. When the orientation of every component is reversed, this invariant is changed only by a sign.

Proof. Using the fact that the rows of the cofactor matrix A^* of a given matrix A are perpendicular to different rows of A , we see that Ω is perpendicular to all the vectors of translation operation induced by partial conjugations as well as the vector R . Certainly, $\Omega \neq 0$. So $\mu \cdot \Omega$ is a non-trivial link-homotopy invariant of ordered, oriented links with 6 components. It is of finite type since it is a polynomial in l_{ij} and $\mu(rst)$. Under the reversion of orientation, μ changes to $-\mu + R$. Since $R \cdot \Omega = 0$, the invariant $\mu \cdot \Omega$ is only changed by a sign under the reversion of orientation. \square

To finish, let us furnish some data obtained using *Mathematica*. Let $\deg(l_{ij}) = 1$, then Ω_i is a homogeneous polynomial of degree 20 in l_{ij} . Let L_i be the number of monomials in Ω_i , the sequence $\{L_1, L_2, \dots, L_{20}\}$ is given as follows:

$$\{5531, 5555, 5555, 5531, 5424, 5769, 5802, 5734, 5753, 5432, \\ 5432, 5753, 5802, 5734, 5769, 5424, 5928, 5922, 5922, 5928\}.$$

Thus $\mu \cdot \Omega$ is linear and homogeneous in $\mu(rst)$ and has 113,700 monomials.