

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 47 (2001)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** GROUPS ACTING ON THE CIRCLE  
**Autor:** GHYS, Étienne  
**Kapitel:** 2. Some classical definitions  
**DOI:** <https://doi.org/10.5169/seals-65441>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## CONTENTS

1. Introduction .....	329
2. Some classical definitions .....	330
3. Two basic examples .....	332
3.1. The projective group .....	333
3.2. Piecewise linear groups .....	336
4. The group of homeomorphisms of the circle .....	338
4.1. Locally compact groups acting on the circle .....	345
5. Rotation numbers .....	349
5.1. Dynamics of a single homomorphism .....	349
5.2. Tits' alternative .....	359
6. Bounded Euler class .....	363
6.1. Group cohomology .....	363
6.2. The Euler class of a group acting on the circle .....	366
6.3. Bounded cohomology and the Milnor-Wood inequality ..	367
6.4. Explicit bounds on the Euler class .....	372
6.5. Actions of the real line and orderings .....	373
6.6. Some examples .....	382
7. Higher rank lattices .....	386
7.1. Witte's theorem .....	387
7.2. Actions of higher rank lattices .....	390
7.3. Lattices in linear groups .....	392
7.4. Some groups that do act... .....	401
References .....	404

## 2. SOME CLASSICAL DEFINITIONS

We begin with some very general definitions concerning group actions. For an introduction to this subject, we refer to [42].

Let  $\Gamma$  be any group and  $X$  be any topological space. An *action* of  $\Gamma$  on  $X$  is a homomorphism  $\phi$  from  $\Gamma$  to the group  $\text{Homeo}(X)$  of homeomorphisms of  $X$ . An element  $\gamma \in \Gamma$  and a point  $x \in X$  produce the point  $\gamma \cdot x = \phi(\gamma)(x)$ . Conversely a map

$$(\gamma, x) \in \Gamma \times X \mapsto \gamma \cdot x \in X$$

comes from an action if for every  $\gamma$ , the point  $\gamma \cdot x$  depends continuously on  $x$  and if for every  $\gamma_1, \gamma_2$  we have  $\gamma_1 \cdot (\gamma_2 \cdot x) = (\gamma_1 \gamma_2) \cdot x$  and  $e \cdot x = x$  ( $e$  denotes the identity element in  $\Gamma$ ).

Two actions  $\phi_1$  and  $\phi_2$  of  $\Gamma$  on  $X_1$  and  $X_2$  are *conjugate* if there exists a homeomorphism  $h$  from  $X_1$  to  $X_2$  such that for every  $\gamma \in \Gamma$ , one has  $\phi_2(\gamma) = h\phi_1(\gamma)h^{-1}$ .

An action  $\phi$  is *faithful* if it is injective, *i.e.* if non trivial elements in the group act non trivially on the space. This is a minor assumption since we can always consider the associated faithful action of the quotient group  $\Gamma / \ker(\phi)$ .

The *orbit* of a point  $x$  is the set  $\mathcal{O}(x) = \{\phi(\gamma)(x) \mid \gamma \in \Gamma\} \subset X$ . The main object of topological dynamics is to study the topological properties of the partition of  $X$  into orbits. An action is *transitive* if there is only one orbit. We say in this case that  $X$  is *homogeneous* under the action of  $\Gamma$ . Of course, these transitive actions are quite trivial from the topological dynamics point of view but this does not mean that the geometrical study of homogeneous spaces is not interesting !

The *stabilizer* of the point  $x$  is the subgroup

$$Stab(x) = \{\gamma \in \Gamma \mid \phi(\gamma)(x) = x\} \subset \Gamma.$$

There is a natural bijection between the quotient  $\Gamma / Stab(x)$  and the orbit  $\mathcal{O}(x)$ . Note that the stabilizers of two points in the same orbit are conjugate subgroups in  $\Gamma$ . An action is *free* if the stabilizer of every point is trivial, *i.e.* if the action of a non trivial element of  $\Gamma$  has no fixed point.

In some cases,  $\Gamma$  might be a topological group. In these cases, we frequently consider *continuous actions* such that  $\gamma \cdot x$  is a continuous function on  $\Gamma \times X$ . The orbit map bijection from  $\Gamma / Stab(x)$  to  $\mathcal{O}(x)$  is continuous but is usually not a homeomorphism when  $\mathcal{O}(x)$  is equipped with the induced topology from  $X$ . The easiest non trivial example is the case where  $\Gamma = \mathbf{R}$ , *i.e.* of a topological flow: if the stabilizer of a point  $x$  is trivial, the orbit  $\mathcal{O}(x)$  is the image of a continuous bijection  $\mathbf{R} \rightarrow \mathcal{O}(x) \subset X$  but in many cases this orbit might be recurrent (for instance dense in  $X$ ) and this bijection is not a homeomorphism. There is however a special case in which this map is indeed a homeomorphism and we use this fact constantly (and sometimes implicitly) in these notes. Consider a Lie group  $G$  acting continuously and transitively on a manifold  $M$  and denote by  $H$  the stabilizer of a point. Then  $H$  is a closed subgroup of  $G$ , hence a closed Lie subgroup, and the quotient space  $G/H$  is naturally a smooth manifold. In this case, the orbit map from  $G/H$  to  $M$  is a homeomorphism.