

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON AN ASSERTION IN RIEMANN'S HABILITATIONSVORTRAG
Autor: Di SCALA, Antonio J.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-65428>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.11.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON AN ASSERTION IN RIEMANN'S HABILITATIONSVORTRAG

by Antonio J. DI SCALA^{*})

ABSTRACT. We study an assertion in Riemann's Habilitation Lecture of 1854. Namely, the determination of the metric given $n^{\frac{n-1}{2}}$ sectional curvatures.

1. INTRODUCTION

Modern differential geometry was born with Riemann's Habilitation Lecture "Ueber die Hypothesen, welche der Geometrie zu Grunde liegen" (*On the Hypotheses which lie at the Foundations of Geometry*) of 1854 at Göttingen [R], [We]. In this lecture Riemann defines the curvature tensor R . One says that M is *flat* if M is locally isometric to \mathbf{R}^n with the usual metric; the tensor R vanishes if and only if the metric is flat. M. Spivak [Sp1] translates Riemann's Lecture and explains it in modern terms. Let

$$Q(X, Y) := \frac{\langle R(X, Y) Y, X \rangle}{|X \wedge Y|^2}$$

be the sectional curvature. Spivak [Sp1, p.4B-25], [Sp2, p.176] makes the following

^{*}) Supported by a CONICET fellowship, partially supported by CONICOR and Secyt-UNC.

ASSERTION 1.1. *If M is n -dimensional and if $Q = 0$ for $n^{\frac{n-1}{2}}$ independent 2-dimensional subspaces of each M_q , then M is flat.*

It is well known that the metric is flat if and only if the sectional curvature Q vanishes identically. The number $n^{\frac{n-1}{2}}$ of Assertion 1.1 is “deduced” from the following “counting argument” given by Riemann: the metric $ds^2 = \sum g_{ij} dx_i dx_j$ contains $\frac{n(n+1)}{2}$ functions while a new coordinate system involves only n functions, so that we can change only n of the g_{ij} , leaving $\frac{n(n-1)}{2}$ other functions which depend on the metric; thus there should be some set of $\frac{n(n-1)}{2}$ functions which will determine the metric completely (see [Di, p.198], [Sp1, p.4B-4]). We quote from the original text as follows [We], [R]:

“...wenn also das Krümmungsmaß in jedem Punkte in $n^{\frac{n-1}{2}}$ Flächenrichtungen gegeben wird, so werden daraus die Maßverhältnisse der Mannigfaltigkeit sich bestimmen lassen, wofern nur zwischen diesen Werthen keine identischen Relationen stattfinden, was in der That, allgemein zu reden, nicht der Fall ist.”

“... es reicht aber nach der frühern Untersuchung, um die Maßverhältnisse zu bestimmen, hin zu wissen, daß es in jedem Punkte in $n^{\frac{n-1}{2}}$ Flächenrichtungen, deren Krümmungsmaße von einander unabhängig sind, Null sei.”

We remark that this text is omitted by Hermann Weyl in his discussion of Riemann’s ideas. Relating the curvature tensor to the metric is a very classical subject and we refer to [Ku, Ya, B] for further details.

In this note we construct several families of counter-examples to Assertion 1.1. In §2 we discuss the space of algebraic curvature tensors and construct an algebraic curvature tensor in dimension 3 which has vanishing sectional curvature on three independent 2-planes: this shows that Assertion 1.1 is not an algebraic consequence of the curvature tensor identities. Let H^2 , S^2 and T^k denote the hyperbolic plane, the sphere and the torus with the metrics of constant curvature -1 , 1 , and 0 . Give $M = S^2 \times H^2 \times T^k$ the product metric; this manifold is not flat. In §3 we construct local orthonormal frames $\{e_i\}$ and local coordinate frames ∂_i for the tangent bundle such that the sectional curvatures $Q(e_i, e_j)$ and $Q(\partial_i, \partial_j)$ vanish for $i \neq j$. Again this shows Assertion 1.1 is false. Finally, in §4, we use warped products to construct still other examples of non-flat metrics which are counter-examples to Assertion 1.1. It is a pleasant task to thank Professors V. Cortez and P. Gilkey for helpful discussions concerning these matters.