

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON AN ASSERTION IN RIEMANN'S HABILITATIONSVORTRAG
Autor: Di SCALA, Antonio J.
Kapitel: 5. CONCLUDING COMMENTS
DOI: <https://doi.org/10.5169/seals-65428>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. CONCLUDING COMMENTS

In order to solve the local equivalence problem (i.e. when two metrics $\mathbf{g}_1, \mathbf{g}_2$ on a differentiable manifold M^n differ (locally) by a diffeomorphism), Riemann tried to compute $n\frac{n-1}{2}$ $\text{Diff}(M^n)$ -equivariant functions (i.e. $K(\mathbf{g}_2)(p) = K(\mathbf{g}_1)(f(p))$ for all $f \in \text{Diff}(M^n)$, $p \in M^n$, $\mathbf{g}_2 = f^*\mathbf{g}_1$). The Gaussian curvature K is such a function when $n = 2$. To do this, Riemann expanded the metric in normal coordinates and defined a map Q from \mathcal{M}_n , the space of Riemannian metrics on M^n , to $C^\infty(G_2(M^n))$, where $G_2(M^n)$ is the two-Grassmannian bundle over M^n . In other words, $Q(\mathbf{g})(\pi_p)$ is the sectional curvature of the 2-plane π_p at $p \in M^n$ with respect to the metric \mathbf{g} . Then he said that "... if the curvature is given in $n\frac{n-1}{2}$ surface directions at every point, then the metric relations of the manifold may be determined ..." [Sp2, p. 144]. More precisely, Riemann took $n\frac{n-1}{2}$ independent sections π_{ij} of the bundle $G_2(M^n)$ and he defined the $n\frac{n-1}{2}$ functions by composing with Q (i.e. a map from \mathcal{M}_n to $\{C^\infty(M^n)\}^{n\frac{n-1}{2}}$). Perhaps the expression of Q in coordinates, the two-dimensional flat case and the counting argument led Riemann to the wrong conclusion that Q can be recovered from evaluation in $n\frac{n-1}{2}$ independent 2-planes. It is hard to believe that he did not observe that this map is not actually a $\text{Diff}(M^n)$ -equivariant morphism, as follows from the fact that a generic diffeomorphism does not preserve the π_{ij} (i.e. $f^*\pi_{ij} \neq \pi_{ij}$) when $n > 2$.

REMARK 5.1. A way of defining $n\frac{n-1}{2}$ $\text{Diff}(M^n)$ -equivariant functions from \mathcal{M}_n to $C^\infty(M^n)$ such that:

- (i) if $n = 2$ then the function is the Gauss curvature K ;
 - (ii) if the $n\frac{n-1}{2}$ functions vanish identically then the metric \mathbf{g} is flat;
- is as follows. Regarding the curvature tensor R as a symmetric endomorphism of the second exterior product bundle $\bigwedge^2(M^n)$ one can take the characteristic polynomial $\chi_R(X)$ of R . Then the coefficients of $\chi_R(X)$ are the required $n\frac{n-1}{2}$ functions.

REFERENCES

- [B] BERGER, M. Riemannian geometry during the second half of the twentieth century. *Jahresber. Deutsch. Math.-Verein.* 100 (1998), 45–208.
- [Di] DIEUDONNÉ, J. *Abrégé d'histoire des mathématiques 1700–1900*. Vol. II. Hermann, 1978.