L'Enseignement Mathématique
Commission Internationale de l'Enseignement Mathématique
47 (2001)
1-2: L'ENSEIGNEMENT MATHÉMATIQUE
THE POSITIVE CONE OF SPHERES AND SOME PRODUCTS OF SPHERES
MATTHEY, Michel / SUTER, Ulrich
3. The -cone and the c-cone
https://doi.org/10.5169/seals-65432

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

computation (based on Proposition 2.4) for the sphere  $S^{2n}$  shows that one has  $q_n = (-1)^{n-1}(n-1)!$ , as claimed.

# 3. The $\gamma$ -cone and the *c*-cone

In general, the problem of computing the geometric dimension of vector bundles is very complicated, as is any general lifting problem in homotopy theory. So, the same is true for the positive cone. That is why we now introduce what we call the  $\gamma$ -cone and the *c*-cone. They are supposed to be easier to compute and might be good approximations to the positive cone. As we will see, these two cones coincide for torsion-free spaces.

DEFINITION 3.1.

i) The  $\gamma$ -cone of X is defined by

$$K_{\gamma}(X) := \{ (n, x) \in \mathbb{Z} \oplus \widetilde{K}(X) \mid \gamma^{k}(x) = 0 \text{ for all } k > n \}.$$

The  $\gamma$ -dimension of a class  $x \in \widetilde{K}(X)$ , denoted by  $\gamma$ -dim(x), is the least integer n such that  $\gamma^k(x) = 0$  for all k > n, in other words, it is the degree (in the variable t) of the polynomial  $\gamma_t(x)$ .

ii) The *c*-cone of X is defined by

 $K_c(X) := \{(n, x) \in \mathbb{Z} \oplus \widetilde{K}(X) \mid c_k(x) = 0 \text{ for all } k > n \}.$ 

The *c*-dimension of a class  $x \in \widetilde{K}(X)$ , denoted by c-dim(x), is the least integer n such that  $c_k(x) = 0$  for all k > n, in other words, it is the degree (in the variable t) of the polynomial  $c_x(t)$ .

Let us point out that the "lower boundary" of the positive cone  $K_+(X)$ , as a subset of  $\widetilde{K}(X) \oplus \mathbb{Z}$ , coincides with the graph of the geometric dimension function g-dim:  $\widetilde{K}(X) \longrightarrow \mathbb{Z}$  (the positive elements consisting exactly of the boundary and the points located above it). The analogous statements hold for the  $\gamma$ -cone and the *c*-cone with respect to the corresponding dimension function.

The following results on these objects follow readily from our preliminaries on K-theory.

**PROPOSITION 3.2.** Let X be a connected finite CW-complex. Then

- i)  $g-\dim(x) \le \dim(X)/2$ , for any  $x \in \widetilde{K}(X)$ ;
- ii)  $\gamma$ -dim(x)  $\leq$  g-dim(x), for any  $x \in \widetilde{K}(X)$ ;
- iii)  $K_+(X) \subseteq K_{\gamma}(X)$ ;
- iv)  $\operatorname{c-dim}(x) \leq \operatorname{g-dim}(x)$ , for any  $x \in \widetilde{K}(X)$ ;
- v)  $K_+(X) \subseteq K_c(X)$ .

This proposition shows that the  $\gamma$ -cone and the *c*-cone are approximations of the positive cone, more precisely, that they constitute upper bounds of the latter.

It turns out that the  $\gamma$ -cone and the *c*-cone coincide for torsion-free spaces, i.e. those spaces having no torsion in their integral cohomology.

PROPOSITION 3.3. Let X be a connected finite CW-complex. If X is torsion-free, then

$$K_{\gamma}(X) = K_{c}(X) \, .$$

*Proof.* The result follows immediately from Proposition 2.2 and injectivity of the Chern character for a torsion-free space.  $\Box$ 

It is worth mentioning that there is no general comparison statement for the  $\gamma$ -cone and the *c*-cone, i.e. there are spaces with torsion for which the  $\gamma$ -cone is not contained in the *c*-cone, and spaces with torsion for which the *c*-cone is not contained in the  $\gamma$ -cone. Moreover, there exist spaces for which the  $\gamma$ -cone and the *c*-cone strictly contain the positive cone (the product  $S^4 \times S^4$  is such an example as we will later see). We now illustrate the situation by three examples.

# EXAMPLES.

i) Let  $j: BSU(3) \longrightarrow BU(3)$  be the map induced by the inclusion of the special unitary group SU(3) in U(3). Then the composition map

$$BSU(3) \xrightarrow{J} BU(3) \xrightarrow{\gamma_3} BU$$

lifts to a map  $f: BSU(3) \longrightarrow BSU$ . Consider W the homotopy fibre of f. It

enters in a pull-back diagram



where  $SU \simeq \Omega BSU \hookrightarrow PBSU \to BSU$  is the path-loop fibration of BSU. The Leray-Serre spectral sequence in cohomology for this fibration is well-known and maps via  $f^*$  to the corresponding spectral sequence for the fibration  $\pi$ . By Lemma 2.5, one has

$$f^*(\widetilde{c}_3) = j^* \circ \widetilde{\gamma}_3^*(c_3(\widetilde{\rho}_3)) = c_3(\widetilde{\gamma}_3) = 2\widetilde{c}_3.$$

Similarly, one has  $f^*(\tilde{c}_2) = c_2(\tilde{\gamma}_3)$ , which is easily seen to vanish. For the cohomology of W in degree  $\leq 6$ , letting  $a_4 := \pi^*(\tilde{c}_2)$  and  $b_6 := \pi^*(\tilde{c}_3)$ , we have computed that  $x_3^2 = 0$  and

$$H^{\leq 6}(W; \mathbf{Z}) = \mathbf{Z} \cdot 1 \oplus \mathbf{Z} \cdot x_3 \oplus \mathbf{Z} \cdot a_4 \oplus \mathbf{Z} \cdot x_5 \oplus \underbrace{\mathbf{Z} \cdot b_6}_{\cong \mathbf{Z}/2} \cong \mathbf{Z}^4 \oplus \mathbf{Z}/2$$

where deg $(x_{2j+1}) = 2j+1$ . The inclusion  $i: Y := W^{[7]} \hookrightarrow W$  of the 7-skeleton of W induces an isomorphism in cohomology up to degree 6. If we let  $x := i^* \circ \pi^* \circ j^*(\tilde{\rho}_3) \in \tilde{K}(Y)$ , we find  $c_3(x) = b_6 \neq 0$ , whereas  $\gamma^k(x) = 0$ , for all  $k \geq 3$ . Indeed, this is clear for  $k \geq 4$  since then  $\gamma^k(\tilde{\rho}_3) = 0$ , and  $\gamma^3(x) = 0$  because its classifying map is the composition  $f \circ \pi \circ i$ , which is homotopically trivial. Thus c-dim(x) = 3 and  $\gamma$ -dim $(x) \leq 2$ . Consequently, Yis a connected finite CW-complex with a strict inclusion

$$K_c(Y) \subsetneq K_{\gamma}(Y)$$
.

ii) Consider the Moore space  $M = M(\mathbb{Z}/2, 5)$ , i.e. the mapping cone of a continuous map  $f: S^5 \xrightarrow{2} S^5$  of degree two, or more explicitly,  $M = C_f = S^5 \cup_2 e^6$ . The exact sequences in cohomology and in *K*-theory of the cofibration  $S^5 \hookrightarrow M \xrightarrow{q} M/S^5 \simeq S^6$  give epimorphisms

$$q^* \colon \mathbf{Z} \cong H^6(S^6; \, \mathbf{Z}) \twoheadrightarrow H^6(M; \, \mathbf{Z}) \cong \mathbf{Z}/2$$
$$q^* \colon \mathbf{Z} \cong \widetilde{K}(S^6) \twoheadrightarrow \widetilde{K}(M) \cong \mathbf{Z}/2 \,.$$

Let x and a be suitable generators of  $\widetilde{K}(S^6)$  and of  $H^6(S^6; \mathbb{Z})$  respectively, and define  $\overline{x} := q^*(x)$  and  $\overline{a} := q^*(a)$ . For obvious dimensional reasons, the Chern classes  $c_1(\overline{x})$  and  $c_2(\overline{x})$  vanish. Moreover, one has  $c_3(\overline{x}) = q^*(c_3(x)) =$  $q^*(2a) = 0$  (see Proposition 2.4), hence c-dim( $\overline{x}$ ) = 0. On the other hand, we have  $\gamma^1(\overline{x}) = \overline{x} \neq 0$ , so  $\gamma$ -dim( $\overline{x}$ )  $\geq 1$ ; more precisely,  $\gamma^2(\overline{x})$  is  $q^*(-S(3, 2) \cdot x) = q^*(-3x) = \overline{x} \neq 0$  and  $\gamma^3(\overline{x}) = q^*(2S(3, 3) \cdot x) = 0$ , so  $\gamma$ -dim( $\overline{x}$ ) = 2. Consequently, M is a connected finite CW-complex with a strict inclusion

$$K_{\gamma}(M) \subsetneq K_c(M)$$
.

iii) Let  $Z = Y \lor M$  be the wedge of the preceding two examples. It is a 7-dimensional finite connected CW-complex for which none of  $K_{\gamma}(Z)$  and  $K_c(Z)$  contains the other one. (The product  $Y \times M$  would also do.)

To end the present section, we prove that the cones are semigroups and homotopy invariants.

PROPOSITION 3.5. The positive cone, the  $\gamma$ -cone and the c-cone of a connected finite CW-complex X are sub-semigroups of K(X) and homotopy invariants of X. Moreover, the positive cone is a sub- $\lambda$ -semiring of K(X).

*Proof.* The homotopy invariance is obvious for the three cones. We have already mentioned in the preliminaries that the positive cone is a sub-semiring of K(X). It is also clear that it is a sub- $\lambda$ -semiring. The "exponentiality" of  $\gamma_t$  and of c (the total Chern class) immediately show that the  $\gamma$ -cone and the c-cone are sub-semigroups of K(X).

We do not know if in general the  $\gamma$ -cone and the *c*-cone are sub- $\lambda$ -semirings of K(X).

## 4. The positive cone of the spheres

We now intend to compute the positive cone of the spheres. For odddimensional spheres, there is nothing to do since  $\widetilde{K}(S^{2n+1}) = 0$ . Whereas for even-dimensional spheres, one has  $\widetilde{K}(S^{2n}) = \mathbb{Z} \cdot x \cong \mathbb{Z}$ , so we only have to compute g-dim(lx) for all integers l.

By Proposition 2.4, we have

$$c(lx) = c(x)^{l} = (1 + (-1)^{n-1}(n-1)! \cdot a)^{l} = 1 + (-1)^{n-1}l(n-1)! \cdot a,$$