Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 47 (2001)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE POSITIVE CONE OF SPHERES AND SOME PRODUCTS OF

SPHERES

Autor: MATTHEY, Michel / SUTER, Ulrich

Kapitel: 5. FURTHER PROPERTIES OF THE CONES

DOI: https://doi.org/10.5169/seals-65432

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 27.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

LEMMA 4.2. For $m \ge 2k + 1$, the group $\pi_m(BU(k))$ is finite.

Proof. We fix $m \ge 3$. The fibration $BU(k-1) \longrightarrow BU(k)$, with fibre S^{2k-1} , yields the following long exact sequence in homotopy:

$$\dots \to \pi_m(S^{2k-1}) \to \pi_m(BU(k-1)) \to \pi_m(BU(k)) \to \pi_{m-1}(S^{2k-1}) \to \dots$$

By Serre [Serre], $\pi_j(S^{2k-1})$ is finite for $j \neq 2k-1$, and we can conclude by induction over k (with $k \geq 1$ and $2k+1 \leq m$), since when k=1, one has $\pi_m(BU(1)) = \pi_{m-1}(U(1)) = 0$ for $m \geq 3$.

From this, we now infer that the image of $(i_k)_*$ is zero for k < n. This implies that $g\text{-}\dim(lx) = n$ when $l \neq 0$, and concludes the second proof.

REMARK 4.3.

i) Since we were motivated by Elliott's classification of unital C^* -algebras of type AF by means of their K-theory, their positive cone and the K-theory class [1] of the unit (see [Bla1]), it is important to single out the fact that the positive cone of S^{2n} and that of S^{2m} are non-isomorphic as monoids if n is different from m. (There is no need here to distinguish the K-theory class 1 of the trivial one-dimensional bundle.) Let us provide a short proof of this claim. For $n \ge 1$, let M_n denote the positive cone of S^{2n} (identified as above with a sub-monoid of \mathbb{Z}^2 , in order to designate its elements). The abelian monoid M_n has a minimal set A_n of generators, in other words a generating set (as a monoid) that is contained in any other generating set, namely

$$A_n = \{(0, 1)\} \cup \{(k, n) \mid k \in \mathbf{Z} \setminus \{0\}\}.$$

Now, consider the function $\sigma: A_n \longrightarrow \{2, 3, ...\}$ defined, for $x \in A_n$, by $\sigma(x) := \min\{l \ge 2 \mid lx \text{ decomposes as a sum of elements of } A_n \setminus \{x\}\}$.

Clearly, such an l exists for any $x \in A_n$ and $\sigma(A_n) = \{2, 2n\}$. Since A_n and σ are isomorphism invariants of M_n , this proves our claim.

ii) For odd-dimensional spheres the positive cone is "trivial"; in other words, $K(S^{2n-1}) = \mathbb{Z}$ and $K_+(S^{2n-1}) = \mathbb{N}$.

5. FURTHER PROPERTIES OF THE CONES

We now investigate naturality properties and behaviour under products of the positive cone, the γ -cone and the c-cone.

The following result is obvious.

PROPOSITION 5.1. Let $f: X \longrightarrow Y$ be a map between connected finite CW-complexes. Let $f^*: K(Y) \longrightarrow K(X)$ be the λ -homomorphism induced by f. Then, for any $y \in \widetilde{K}(Y)$, one has

$$\begin{aligned} & \text{g-dim}(f^*(y)) \leq \text{g-dim}(y) \\ & \gamma\text{-dim}(f^*(y)) \leq \gamma\text{-dim}(y) \\ & \text{c-dim}(f^*(y)) \leq \text{c-dim}(y) \,, \end{aligned}$$

and in particular,

$$f^*(K_+(Y)) \subseteq K_+(X)$$
$$f^*(K_\gamma(Y)) \subseteq K_\gamma(X)$$
$$f^*(K_c(Y)) \subseteq K_c(X).$$

Furthermore, if f^* is an isomorphism, then

$$f^*(K_{\gamma}(Y)) = K_{\gamma}(X).$$

For the next corollary we need a new definition.

DEFINITION 5.2. Let X and Y be two connected finite CW-complexes. A map $f: X \longrightarrow Y$ is called a K^0 -equivalence (or K-equivalence for short) if there exists a map $g: Y \longrightarrow X$ such that, on the level of the K^0 -groups,

$$f^* \circ g^* = Id_{K^0(X)}$$
 and $g^* \circ f^* = Id_{K^0(Y)}$.

Note that a K-equivalence is *not* necessarily a homotopy equivalence: there are homotopically non-trivial (i.e. non-contractible) finite CW-complexes X for which $\widetilde{K}(X) = 0 = \widetilde{K}(pt)$; see example iii) below.

PROPOSITION 5.3. If $f: X \longrightarrow Y$ is a K-equivalence, then f induces the following isomorphisms of semigroups:

$$K_{+}(Y) \stackrel{f^*}{\cong} K_{+}(X)$$
 and $K_{\gamma}(Y) \stackrel{f^*}{\cong} K_{\gamma}(X)$.

Proof. Applying Proposition 5.1 twice, we get (in the notations of Definition 5.2)

$$K_{+}(X) = f^{*} \circ q^{*}(K_{+}(X)) \subseteq f^{*}(K_{+}(Y)) \subseteq K_{+}(X)$$
.

This establishes the first isomorphism, whereas the second one is obvious. \Box

The following result is more technical to state.

COROLLARY 5.4. Let X and Y be two connected finite CW-complexes. Assume that $K^1(X) = 0$ and that $\widetilde{K}^0(Y) = 0$. Then the projection $p: X \times Y \longrightarrow X$ induces isomorphisms

$$K_{+}(X) \stackrel{p^*}{\cong} K_{+}(X \times Y)$$
 and $K_{\gamma}(X) \stackrel{p^*}{\cong} K_{\gamma}(X \times Y)$.

Proof. Invoking the Künneth theorem for K-theory, our hypotheses imply that $p^*: K^0(X) \longrightarrow K^0(X \times Y)$ is an isomorphism with inverse i^* , where i is the inclusion of X in $X \times Y$. Consequently, p^* is a K-equivalence. \square

The following is a useful result.

PROPOSITION 5.5. Let X and Y be connected finite CW-complexes. Assume that the positive cone and the γ -cone of Y coincide, and let $f: X \longrightarrow Y$ be a map inducing an isomorphism $f^*: K(Y) \longrightarrow K(X)$. Then f induces an isomorphism of positive cones, and the γ -cone of X coincides with the positive cone:

$$K_+(Y) \stackrel{f^*}{\cong} K_+(X) = K_\gamma(X)$$
.

Proof. By Proposition 5.1 we have $f^*(K_+(Y)) = f^*(K_\gamma(Y)) = K_\gamma(X)$ and $f^*(K_+(Y)) \subseteq K_+(X)$, hence $K_\gamma(X) \subseteq K_+(X)$. We conclude with iii) of Proposition 3.2. \square

EXAMPLES.

i) Let X be a connected finite CW-complex of dimension ≤ 3 . Since for suitable CW-decompositions, one has $BU(1)^{[3]} = BU^{[3]}$ and since $BU(1) = \mathbf{C}P^{\infty} = K(\mathbf{Z}, 2)$, any $x \in \widetilde{K}(X) = [X, BU]$ lifts to a class in [X, BU(1)], giving an isomorphism $\widetilde{K}(X) \cong H^2(X; \mathbf{Z})$ mapping x to $c_1(x)$. It follows that the positive cone coincides with the c-cone and is given by

$$K_+(X) = \mathbf{N} \times \{0\} \cup \mathbf{N}^* \times \widetilde{K}(X) \subset \mathbf{Z} \times \widetilde{K}(X).$$

ii) Example i) applies to a closed oriented surface Σ_g of genus g. Since it is torsion-free, its positive cone coincides with its c-cone and with its γ -cone. Moreover, let $f \colon \Sigma_g \longrightarrow S^2$ be a map of degree 1 (it exists, since both the 2-sphere and Σ_g are quotients of the square $[0,1]^2$). Then f not only induces an isomorphism in K-theory, but also an isomorphism of positive cones, as follows from Proposition 5.1.

- iii) Let X and Y be the Moore spaces $M(\mathbb{Z}/3, 2q+11) = S^{2q+11} \cup_3 e^{2q+12}$ and $M(\mathbb{Z}/3, 2q-1) = S^{2q-1} \cup_3 e^{2q}$ respectively. In [Adams], Adams shows that for q large enough, there exists a map $A: X = \Sigma^{12}Y \longrightarrow Y$ such that the induced map $A^*: \widetilde{K}(Y) \longrightarrow \widetilde{K}(X)$ is an isomorphism (take p = m = 3, f = 1 and r = 6 in Theorem 1.7 and in Lemmas 12.4 and 12.5 of [Adams]). Therefore, A is a K-isomorphism between simply connected finite CW-complexes, but it is *not* a homotopy equivalence. The mapping cone C_A is a non-contractible finite CW-complex with $\widetilde{K}(C_A) = 0$. (It is non-contractible because its homology is non-trivial.)
- iv) In [GrMo], pp. 203-206, a CW-complex $X = (S^1 \vee S^2) \cup e^3$ is defined, with the property that the inclusion $i: S^1 = X^{[1]} \hookrightarrow X$ of the 1-skeleton induces an isomorphism in integral homology (and on the level on fundamental groups); however, i is not a homotopy equivalence since $\pi_2(X) \neq 0$. Consequently, by the universal coefficient theorem (see Corollary V.7.2 in [Bred]), i induces an isomorphism in integral cohomology, and, by a direct application of the Atiyah-Hirzebruch spectral sequence, also in K-theory. In particular, i is a K-equivalence, but not an equivalence. (As C_A in the preceding example, the quotient space $X/X^{[1]}$ has vanishing \widetilde{K} , however it is the closed 3-ball and is therefore contractible.)

Let us finally mention that in [Matt], the positive cone, the c-cone and the γ -cone are also studied from the rational point of view, and rational K-theory is considered.

6. The cones of the products $S^n \times S^{2m-1}$

In this section, we will compute the cones for the products $S^{2n} \times S^{2m-1}$ and $S^{2n-1} \times S^{2m-1}$.

We begin with $S^{2n} \times S^{2m-1}$. Since $\widetilde{K}(S^{2m-1}) = 0$ and $K^1(S^{2n}) = 0$, the answer immediately follows from Proposition 5.5.

Theorem 6.1. The projection $p: S^{2n} \times S^{2m-1} \longrightarrow S^{2n}$ induces an isomorphism of positive cones, and, for $S^{2n} \times S^{2m-1}$, the γ -cone and the c-cone coincide with the positive cone:

$$K_{+}(S^{2n}) \stackrel{p^*}{\cong} K_{+}(S^{2n} \times S^{2m-1}) = K_{\gamma}(S^{2n} \times S^{2m-1}).$$