

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	47 (2001)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE POSITIVE CONE OF SPHERES AND SOME PRODUCTS OF SPHERES
Autor:	MATTHEY, Michel / SUTER, Ulrich
Kapitel:	9. The positive cone of some products of even-dimensional spheres
DOI:	https://doi.org/10.5169/seals-65432

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

9. THE POSITIVE CONE OF SOME PRODUCTS OF EVEN-DIMENSIONAL SPHERES

In this section, using known results from the theory of homotopy groups of spheres, we compute the positive cone of $S^4 \times S^4$, $S^4 \times S^6$, $S^6 \times S^6$ and $S^6 \times S^8$. This computation will in particular show that the positive cone and the γ -cone do not coincide for $S^4 \times S^4$. Keeping notations as in Section 7, we describe the positive cone in terms of the geometric dimension function.

A) We start with the case of $S^4 \times S^4$.

THEOREM 9.1. *The geometric dimension on $\tilde{K}(S^4 \times S^4)$ is given as follows: for $x = ax_1 + bx_2 + lx_1x_2 \in \tilde{K}(S^4 \times S^4)$, one has*

$$g\text{-dim}(x) = \begin{cases} 0 & \text{if } a = b = l = 0 \\ 2 & \text{if } a \neq 0, b = l = 0 \\ 2 & \text{if } b \neq 0, l = ab/6, l \text{ even} \\ 3 & \text{if } b \neq 0, l = ab/6, l \text{ odd} \\ 4 & \text{if } l \neq ab/6 \end{cases}$$

Proof. Theorem 8.2 reduces the problem to the computation of the function $s = s(ab)$, i.e. to calculating $g\text{-dim}(x)$ for the particular stable classes $x = ax_1 + bx_2 + (ab/6)x_1x_2$ (where ab is a multiple of 6), or equivalently the order of $[x_1, x_2]$ in both groups $\pi_7(BU(3))$ and $\pi_7(BU(2))$ (with a little abuse of notation, we denote both Whitehead products by the same symbol). By Samelson [Sam], one has

$$\pi_7(BU(2)) \cong \pi_6(U(2)) \cong \pi_6(SU(2)) \cong \pi_6(S^3) \cong \mathbf{Z}/12,$$

precisely generated by $[x_1, x_2]$. This shows that for these particular values of x , $g\text{-dim}(x) = 2$ if and only if ab is a multiple of 12. This completes the proof. \square

REMARK 9.2.

i) Borel and Hirzebruch in [BoHi] (p.355), applying Bott's results of [Bott1], have proved that

$$\pi_{2n+1}(BU(n)) \cong \pi_{2n}(SU(n)) \cong \mathbf{Z}/n! \quad (n \geq 2),$$

hence $\pi_7(BU(3)) \cong \mathbf{Z}/6$. Moreover, Corollary 8.3 shows that the order of $[x_1, x_2]$ in $\pi_7(BU(3))$ is 6; it is consequently a generator.

ii) As already alluded to, we have just proved that $S^4 \times S^4$ has its positive cone strictly contained in its γ -cone, although it is a torsion-free space.

B) As for $S^4 \times S^4$, classical results from the theory of homotopy groups of the unitary groups allow one to compute the positive cone of $S^4 \times S^6$. In this case, it coincides with the γ -cone.

THEOREM 9.3. *For the product $S^4 \times S^6$, one has*

$$K_+(S^4 \times S^6) = K_c(S^4 \times S^6) = K_\gamma(S^4 \times S^6).$$

The latter is described in Theorem 7.1.

Proof. By Lundell's tables [Lun] (see also [Mim]) and by Remark i) above, one has

$$\pi_9(BU(3)) \cong \mathbf{Z}/12 \quad \text{and} \quad \pi_9(BU(4)) \cong \mathbf{Z}/24.$$

Corollary 8.3 shows that $[x_1, x_2]$ is of order 12 in $\pi_9(BU(4))$. By naturality of the Whitehead product, the homomorphism $j_* = \pi_9(j)$, induced by the map $j: BU(3) \rightarrow BU(4)$, takes the product $[x_1, x_2] \in \pi_9(BU(3))$ to $[x_1, x_2] \in \pi_9(BU(4))$. This implies that $[x_1, x_2]$ is of order 12 in $\pi_9(BU(3))$ too, and that $[ax_1, bx_2]$ vanishes in $\pi_9(BU(3))$ precisely when it is zero in $\pi_9(BU(4))$. Together with Theorem 8.2, this completes the proof. \square

REMARK 9.4. This proof shows in particular that $[x_1, x_2]$ is a generator of $\pi_9(BU(3)) \cong \mathbf{Z}/12$ and that the map $j_*: \pi_9(BU(3)) \rightarrow \pi_9(BU(4))$ is injective.

C) By similar methods, we now show that the positive cone and the γ -cone coincide for $S^6 \times S^6$ and then for $S^6 \times S^8$.

THEOREM 9.5. *For the product $S^6 \times S^6$, one has*

$$K_+(S^6 \times S^6) = K_c(S^6 \times S^6) = K_\gamma(S^6 \times S^6).$$

The latter is given by Theorem 7.1.

Proof. By Lundell's tables [Lun] (see also [Mim]), one has

$$\pi_{11}(BU(3)) \cong \mathbf{Z}/30 \quad \text{and} \quad \pi_{11}(BU(5)) \cong \mathbf{Z}/120.$$

Corollary 8.3 shows that $[x_1, x_2]$ is of order 30 in $\pi_{11}(BU(5))$. By naturality, the map $j_* = \pi_{11}(j)$, induced by $j: BU(3) \rightarrow BU(5)$, takes the Whitehead product $[x_1, x_2] \in \pi_{11}(BU(3))$ to $[x_1, x_2] \in \pi_{11}(BU(5))$. This implies that $[x_1, x_2]$ is of order 30 in $\pi_{11}(BU(3))$ too, and that $[ax_1, bx_2]$ vanishes in $\pi_{11}(BU(3))$ precisely when it is zero in $\pi_{11}(BU(5))$. Together with Theorem 8.2, this completes the proof. \square

REMARK 9.6.

i) This shows that $[x_1, x_2]$ generates $\pi_{11}(BU(3)) \cong \mathbf{Z}/30$ and that the map $j_*: \pi_{11}(BU(3)) \longrightarrow \pi_{11}(BU(5))$ is injective.

ii) We were also able to prove this theorem without appealing to results on homotopy groups of $BU(n)$. Using spectral sequence arguments, we have computed the first few stages of the Moore-Postnikov tower of the map $BSU(3) \longrightarrow BSU(5)$. This computation, being extremely lengthy, is not given here (see [Matt]).

Now we move on to the product $S^6 \times S^8$.

THEOREM 9.7. *For the product $S^6 \times S^8$, one has*

$$K_+(S^6 \times S^8) = K_c(S^6 \times S^8) = K_\gamma(S^6 \times S^8).$$

The latter is described in Theorem 7.1.

Proof. By Lundell's tables [Lun] (see also [Mim]), one has

$$\pi_{13}(BU(4)) \cong \mathbf{Z}/60 \quad \text{and} \quad \pi_{13}(BU(6)) \cong \mathbf{Z}/720.$$

Corollary 8.3 shows that $[x_1, x_2]$ is of order 60 in $\pi_{13}(BU(6))$. By naturality, the map $j_* = \pi_{13}(j)$, induced by $j: BU(4) \longrightarrow BU(6)$, takes the Whitehead product $[x_1, x_2] \in \pi_{13}(BU(4))$ to $[x_1, x_2] \in \pi_{13}(BU(6))$. This implies that $[x_1, x_2]$ is of order 60 in $\pi_{13}(BU(4))$ too, and that $[ax_1, bx_2]$ vanishes in $\pi_{13}(BU(4))$ precisely when it is zero in $\pi_{13}(BU(6))$. Together with Theorem 8.2, this completes the proof. \square

REMARK 9.8. The proof shows that $[x_1, x_2]$ is a generator of the group $\pi_{13}(BU(4)) \cong \mathbf{Z}/60$ and that the map $j_*: \pi_{13}(BU(4)) \longrightarrow \pi_{13}(BU(6))$ is injective.

10. “GAPS IN COHOMOLOGY” AND THE γ -CONE

In the present section, we are interested in spaces having a “gap in cohomology”, more precisely we look at spaces obtained by attaching a single large-dimensional cell to a finite CW-complex Y . For such spaces, the integral cohomology is zero between the dimension of Y and the top-dimensional class. The products $S^n \times S^m$ are typical examples (see Section 8). For this kind of spaces, the c -cone obviously cannot give information in the