

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 47 (2001)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PROJECTIVE GEOMETRY OF POLYGONS AND DISCRETE 4-VERTEX AND 6-VERTEX THEOREMS
Autor: Ovsienko, V / Tabachnikov, S.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-65426>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

PROJECTIVE GEOMETRY OF POLYGONS AND DISCRETE 4-VERTEX AND 6-VERTEX THEOREMS

by V. OVSIENKO and S. TABACHNIKOV

ABSTRACT. This paper is concerned with discrete versions of three well-known results from projective differential geometry: the four-vertex theorem, the theorem on six affine vertices, and Ghys' theorem on four zeroes of the Schwarzian derivative. We study the geometry of closed polygonal lines in \mathbf{RP}^d and prove that polygons satisfying a certain convexity condition have at least $d + 1$ flattenings. This result provides a new approach to the classical theorems mentioned.

1. INTRODUCTION

A *vertex* of a smooth plane curve is a point where the curve has 4th-order contact with a circle (at a generic point the osculating circle has 3rd-order contact with the curve). An *affine vertex* (or *sextactic point*) of a smooth plane curve is a point of 6th-order contact with a conic. In 1909 S. Mukhopadhyaya [10] published two celebrated theorems, which are joined in the following statement:

Every closed smooth convex plane curve has at least 4 distinct vertices and at least 6 distinct affine vertices.

These results generated an extensive literature. From a modern point of view they are related, among other subjects, to the global singularity theory of wave fronts and to Sturm theory – see e.g. [1, 2, 4, 8, 17, 18] and references therein.

A recent and unexpected result along these lines is the following theorem due to E. Ghys [7]:

The Schwarzian derivative of every diffeomorphism of the projective line has at least 4 distinct zeroes.

(see also [11, 15, 6]). The Schwarzian derivative vanishes when the 3rd jet of the diffeomorphism coincides with that of a projective transformation (at a generic point a diffeomorphism can be approximated by a projective transformation up to the 2nd derivative). Ghys' theorem can be interpreted as the 4-vertex theorem in Lorentzian geometry (cf. references above).

The goal of this note is to study polygonal analogues of the above results. In our opinion, such a discretization of smooth formulations is interesting for the following reasons. Firstly, a discrete theorem is *a priori* stronger; it becomes, in the limit, a smooth one, thus providing a new proof of the latter. An important feature of the discrete approach is the availability of mathematical induction, which can considerably simplify the proofs. Secondly, the very operation of discretization is non-trivial: a single smooth theorem may lead to non-equivalent discrete ones. An example of this phenomenon is provided by two recent versions of the 4-vertex theorem for convex plane polygons [12, 13, 19, 16] – see Remark 2.4 below. To the best of our knowledge, these results are the only available discrete versions of the 4-vertex theorem.

In this regard we would like to draw attention to a famous lemma of Cauchy (1813):

Given two convex (plane or spherical) polygons whose respective sides are congruent, the cyclic sequence of the differences of respective angles of the polygons changes sign at least 4 times.

This result plays a crucial role in the proof of rigidity for convex polyhedra (see [5] for a survey). The Cauchy lemma implies, in the limit, the smooth 4-vertex theorem and can be viewed as the first result in the area under discussion.