

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE RATIONAL FORMS OF NILPOTENT LIE ALGEBRAS AND LATTICES IN NILPOTENT LIE GROUPS  
**Autor:** Semenov, Yu. S.  
**Kapitel:** 2.1 HEISENBERG ALGEBRAS  
**DOI:** <https://doi.org/10.5169/seals-66073>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**THEOREM 2.** *Let  $\mathfrak{g} = \mathfrak{f}_c(p, \mathbf{R})$  be a free nilpotent Lie algebra of class  $c \geq 2$  on  $p$  generators. Then  $\mathfrak{g} \oplus \mathfrak{g}$  and  $\mathfrak{g} \otimes_{\mathbf{R}} \mathbf{C} = \mathfrak{f}_c(p, \mathbf{C})$  (regarded over  $\mathbf{R}$ ) have infinitely many non-isomorphic rational forms.*

In Theorem 3 we also classify all rational forms for three 6-dimensional real nilpotent Lie algebras  $\mathfrak{g}$  (two of them appear in Theorem 2 for  $p = c = 2$ ) which are of class 2 and have 2-dimensional centre coinciding with the derived subalgebra.

In conclusion let us mention a direct way to prove that two given lattices in a nilpotent Lie group are not commensurable. For example, let  $G = UT_3(\mathbf{R})$  be the Lie group of upper triangular  $3 \times 3$ -matrices with 1 on the diagonal,  $\mathfrak{g} = \mathfrak{f}_2(2, \mathbf{R})$  be Lie algebra of  $G$ . Consider  $G \times G$  and its Lie algebra  $\mathfrak{g} \oplus \mathfrak{g}$  which has infinitely many non-isomorphic rational forms  $\mathfrak{h}_m$  ( $m \geq 1$  is a square-free integer), in view and in the notation of Theorems 2, 3 (see Section 4 for more details).

Let  $\Gamma_m$  and  $\Gamma_n$  be corresponding lattices in  $G \times G$  for distinct  $m, n$ . One can prove that the ratio of the covolumes of  $\Gamma_m$  and  $\Gamma_n$  with respect to a Haar measure on  $G \times G$  equals  $m\sqrt{m}/n\sqrt{n}$  up to a rational factor. Hence the lattices are not commensurable. Note that by Proposition 1.1 and Theorem 3  $\Gamma_m$  and  $\Gamma_n$  are not commensurable in any sense.

**ACKNOWLEDGEMENTS.** The author is grateful to Pierre de la Harpe who attracted his attention to problems of commensurability and to the University of Geneva for its hospitality. He thanks also Goulmira Arzhantseva and Thierry Vust for helpful remarks.

## 2. NILPOTENT LIE ALGEBRAS WITH A UNIQUE RATIONAL FORM UP TO ISOMORPHISM

### 2.1 HEISENBERG ALGEBRAS

Let us begin with the following considerations that we will use here and in the next sections (see [2, Chapter 5] for more details). Suppose that a real Lie algebra  $\mathfrak{g}$  has a  $\mathbf{Q}$ -form  $\mathfrak{h}$  and  $\mathfrak{i}$  (resp.  $\mathfrak{a}$ ) is an ideal (resp. a subalgebra) of  $\mathfrak{g}$ . We say that  $\mathfrak{i}$  (resp.  $\mathfrak{a}$ ) is rational if  $\mathfrak{i} \cap \mathfrak{h}$  (resp.  $\mathfrak{a} \cap \mathfrak{h}$ ) is a rational form of  $\mathfrak{i}$  (resp.  $\mathfrak{a}$ ). For instance, the terms  $C^k \mathfrak{g}$  of the lower central series of  $\mathfrak{g}$  are rational as well as centralizers of rational subalgebras or ideals. It is not hard to see that  $\mathfrak{h}/\mathfrak{i} \cap \mathfrak{h}$  is a rational form of the quotient Lie algebra  $\mathfrak{g}/\mathfrak{i}$ .

Let

$$(2.1) \quad \mathfrak{g} = \mathfrak{i}_1 > \mathfrak{i}_2 > \cdots > \mathfrak{i}_{k+1} = 0$$

be a descending series of rational ideals of  $\mathfrak{g}$ . We say that a basis  $X = \{x_1, \dots, x_d\}$  of a rational form  $\mathfrak{h}$  is based on (2.1) if  $x_1, \dots, x_{p_1}$  generate  $\mathfrak{g} \bmod \mathfrak{i}_2$ ,  $x_1, \dots, x_{p_2}$  generate  $\mathfrak{g} \bmod \mathfrak{i}_3$  and so on. It can be shown that such a basis exists for any series (2.1). In the sequel we will use these kinds of bases for a suitable descending series dealing, for instance, with Heisenberg algebras.

Recall that the (generalized) Heisenberg algebra  $\mathfrak{hei}_{2k+1}(\mathbf{R})$  has an  $\mathbf{R}$ -basis  $H_1, \dots, H_{2k+1}$  in which

$$(2.2) \quad [H_1, H_2] = [H_3, H_4] = \cdots = [H_{2k-1}, H_{2k}] = H_{2k+1},$$

other brackets being trivial. Here the 1-dimensional centre is spanned by  $H_{2k+1}$ .

Given an extension (1.1) one can attach to it a 2-cocycle  $\omega: \Lambda^2 \mathfrak{a} \rightarrow \mathbf{R}$  in the usual way. Also  $\omega$  can be regarded as a symplectic form on  $\mathfrak{a}$ . If  $\mathfrak{b} = \mathfrak{hei}_{2k+1}(\mathbf{R})$  then  $\omega$  is the canonical non-degenerate symplectic form with respect to the basis  $H_1, \dots, H_{2k} \pmod{\mathbf{R} \cdot H_{2k+1}}$ .

Let  $d = \dim_{\mathbf{R}} \mathfrak{a}$  and let  $m = d - \text{rank}(\omega)$  be the codimension of the kernel of  $\omega$ . It is not hard to see (cf. the proof of the proposition below) that the Lie algebra  $\mathfrak{b}$  is uniquely defined up to  $\mathbf{R}$ -isomorphism by  $d$  and  $m$ . Namely,

$$\mathfrak{b} \cong \mathfrak{hei}_{d+1-m}(\mathbf{R}) \oplus \mathbf{R}^m.$$

This implies that the centre of  $\mathfrak{b}$  is  $(m+1)$ -dimensional. Thus, two Lie algebras  $\mathfrak{b}_1$  and  $\mathfrak{b}_2$  ( $\dim_{\mathbf{R}} \mathfrak{b}_1 = \dim_{\mathbf{R}} \mathfrak{b}_2$ ) of type (1.1) are not isomorphic if  $m_1 \neq m_2$ .

Evidently,  $\mathfrak{hei}_{2k+1}(\mathbf{Q})$  is a rational form for  $\mathfrak{hei}_{2k+1}(\mathbf{R})$ .

The following proposition holds.

**PROPOSITION 2.1.** *In the above notation let  $\mathfrak{h}$  be a rational form of  $\mathfrak{b}$ . Let  $d = \dim(\mathfrak{b}) - 1$ ,  $m = \dim[\mathfrak{b}, \mathfrak{b}]$  and let  $\mathbf{Q}^m$  denote the abelian Lie  $\mathbf{Q}$ -algebra of dimension  $m$ . Then*

$$\mathfrak{h} \cong \mathfrak{hei}_{d+1-m}(\mathbf{Q}) \oplus \mathbf{Q}^m$$

over  $\mathbf{Q}$ , i.e., there is a unique rational form for  $\mathfrak{b}$  up to isomorphism.

*Proof.* Choose a  $\mathbf{Q}$ -basis  $B_1, \dots, B_{d+1}$  for  $\mathfrak{h}$ . Either all brackets  $[B_i, B_j] = 0$ , and then  $\mathfrak{h} \cong \mathbf{Q}^{d+1}$ , or there are  $i, j$  such that  $[B_i, B_j] = C \neq 0$ .

We may suppose that  $C = B_{d+1}$ . Thus the derived subalgebra of  $\mathfrak{h}$  is spanned by  $B_{d+1}$ . The corresponding symplectic form  $\omega$  is represented by a

skew-symmetric  $d \times d$  matrix  $M = (\mu_{ij})$  with respect to the basis  $B_1, \dots, B_d$  (mod  $[\mathfrak{h}, \mathfrak{h}]$ ). Namely,  $[B_i, B_j] = \mu_{ij} B_{d+1}$ . Over  $\mathbf{Q}$  one can choose a canonical symplectic basis  $\widehat{B}_1, \dots, \widehat{B}_d$  (mod  $[\mathfrak{h}, \mathfrak{h}]$ ) so that the matrix  $\widehat{M}$  representing  $\omega$  has  $l$  blocks of type

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

standing on the diagonal, the other entries being trivial. The rank of  $\omega$  is equal to  $2l$  and  $2l = d - m$ . In the basis  $B_1, \dots, B_{d+1}$  (we omit the 'hats') of  $\mathfrak{h}$

$$[B_1, B_2] = [B_3, B_4] = \dots = [B_{2l-1}, B_{2l}] = B_{d+1},$$

all the other brackets being trivial. This completes the proof.

## 2.2 EXAMPLE OF A FREE NILPOTENT ALGEBRA

Let  $\mathfrak{f}_c(n, \mathbf{R})$  be the free nilpotent Lie algebra of class  $c$  on  $n$  generators. Then  $\mathfrak{f}_c(n, \mathbf{R})$  has a unique rational form  $\mathfrak{f}_c(n, \mathbf{Q})$  up to isomorphism (cf. Theorem 2).

Indeed, let  $\mathfrak{h} = \langle x_1, \dots, x_n, \dots \rangle$  be a rational form of  $\mathfrak{f}_c(n, \mathbf{R})$ . We may suppose that  $x_1, \dots, x_n$  span (modulo the derived subalgebra)  $\mathfrak{h}/[\mathfrak{h}, \mathfrak{h}] \cong \mathbf{Q}^n$ . Consequently,  $\mathfrak{h}$  is generated by  $\{x_1, \dots, x_n\}$  as a Lie algebra. There exists an epimorphism  $\pi: \mathfrak{f}_c(n, \mathbf{Q}) \rightarrow \mathfrak{h}$  because  $\mathfrak{f}_c(n, \mathbf{Q})$  is free. It must be an isomorphism since the dimension of  $\mathfrak{h}$  equals the dimension (not depending on the ground field) of a free nilpotent Lie algebra of class  $c$  on  $n$  generators.

## 2.3 MORE EXAMPLES

The purpose of this subsection is to sketch two more examples of Lie algebras with a unique rational form up to isomorphism.

Let  $\mathfrak{g}_t$ ,  $t \in \mathbf{R}$ , be a family of real 6-dimensional Lie algebras with a basis  $\{x_1, \dots, x_6\}$  such that

$$\begin{aligned} [x_1, x_2] &= x_3, & [x_1, x_3] &= tx_5, & [x_1, x_5] &= x_6, \\ [x_2, x_3] &= x_4, & [x_2, x_4] &= x_5, & [x_3, x_4] &= x_6, \end{aligned}$$

other brackets being trivial. One can show that

1.  $C^k \mathfrak{g}_t = \langle x_{k+1}, \dots, x_6 \rangle$ ,  $k = 2, \dots, 5$ , where  $C^k \mathfrak{g}_t$  are the terms of the lower central series of  $\mathfrak{g}_t$ .
2. The centralizer  $\mathfrak{C}$  of  $C^4 \mathfrak{g}_t$ , that is,  $\mathfrak{C} = \{c \in \mathfrak{g}_t \mid [c, C^4 \mathfrak{g}_t] = 0\}$  is spanned by  $x_2, \dots, x_6$ .