

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	48 (2002)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ON THE RATIONAL FORMS OF NILPOTENT LIE ALGEBRAS AND LATTICES IN NILPOTENT LIE GROUPS
Autor:	Semenov, Yu. S.
Kapitel:	3. Malcev's example
DOI:	https://doi.org/10.5169/seals-66073

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. Real Lie algebras \mathfrak{g}_0 and \mathfrak{g}_1 are not isomorphic but $\forall t \neq 0 \quad \mathfrak{g}_t \cong \mathfrak{g}_1$.
4. If $t \in \mathbf{Q} \setminus \{0\}$ then the rational algebra $\mathfrak{g}_t \cong \mathfrak{g}_1$ over \mathbf{Q} .
5. \mathfrak{g}_0 and \mathfrak{g}_1 are two Lie algebras with a unique rational form up to isomorphism.
6. Let \mathfrak{g} be a split real simple Lie algebra of type G_2 , \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} and $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$ be the triangular decomposition of \mathfrak{g} with respect to \mathfrak{h} . Then \mathfrak{n}_+ is isomorphic to \mathfrak{g}_0 .

3. MALCEV'S EXAMPLE

In this Section we develop Malcev's example and prove Theorem 1.

Suppose that there is a \mathbf{Q} -isomorphism between \mathfrak{g}_t and \mathfrak{g}_s . It must be written in the following form (cf. [5]) since $C^2\mathfrak{g}_t = \langle x_4, x_5, x_6 \rangle$, $C^3\mathfrak{g}_t = \langle x_5, x_6 \rangle$ and the centralizer \mathfrak{c} of $C^2\mathfrak{g}$, which is an ideal in this case, is spanned by x_3, \dots, x_6 .

$$\begin{cases} y_1 = a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + a_{14}x_4 + \dots \\ y_2 = a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + a_{24}x_4 + \dots \\ y_3 = \quad \quad \quad a_{33}x_3 + a_{34}x_4 + \dots \\ y_4 = \quad \quad \quad a_{44}x_4 + \dots \end{cases}$$

We do not explicit the expressions for y_5, y_6 . Here y_1, \dots, y_6 are basis elements of \mathfrak{g}_s satisfying the relations (1.2).

We obtain after straightforward computations that

$$[y_1, y_2] = y_4 = \Delta x_4 + \dots,$$

$\Delta = a_{11}a_{22} - a_{12}a_{21} = a_{44} \neq 0$. On the other hand,

$$\begin{cases} y_5 = [y_1, y_4] = \Delta(a_{11}x_5 + a_{12}x_6), \\ y_6 = [y_2, y_4] = \Delta(a_{21}x_5 + a_{22}x_6). \end{cases}$$

Hence,

$$(3.1) \quad \begin{cases} x_5 = (a_{22}y_5 - a_{12}y_6)/\Delta^2, \\ x_6 = (a_{11}y_6 - a_{21}y_5)/\Delta^2. \end{cases}$$

We need to compute the remaining two brackets. First of all,

$$(3.2) \quad \begin{aligned} [y_1, y_3] &= a_{11}a_{33}[x_1, x_3] + a_{12}a_{33}[x_2, x_3] + a_{11}a_{34}[x_1, x_4] + a_{12}a_{34}[x_2, x_4] \\ &= a_{11}a_{33}x_6 + a_{12}a_{33}(x_5 + tx_6) + a_{11}a_{34}x_5 + a_{12}a_{34}x_6 \\ &= (a_{12}a_{33} + a_{11}a_{34})x_5 + (a_{11}a_{33} + a_{12}a_{34} + ta_{12}a_{33})x_6 = y_6. \end{aligned}$$

Let $u = a_{12}a_{33} + a_{11}a_{34}$, $v = a_{11}a_{33} + a_{12}a_{34} + ta_{12}a_{33}$. In view of (3.1) and (3.2) we have

$$(a_{22}y_5 - a_{12}y_6)u/\Delta^2 + (a_{11}y_6 - a_{21}y_5)v/\Delta^2 = y_6,$$

whence

$$(3.3) \quad \begin{cases} va_{11} - ua_{12} = \Delta^2, \\ va_{21} - ua_{22} = 0. \end{cases}$$

It follows that

$$(3.4) \quad \begin{cases} u = a_{21}\Delta, \\ v = a_{22}\Delta. \end{cases}$$

In addition,

$$(3.5) \quad \begin{aligned} [y_2, y_3] &= a_{21}a_{33}[x_1, x_3] + a_{22}a_{33}[x_2, x_3] + a_{21}a_{34}[x_1, x_4] + a_{22}a_{34}[x_2, x_4] \\ &= a_{21}a_{33}x_6 + a_{22}a_{33}(x_5 + tx_6) + a_{21}a_{34}x_5 + a_{22}a_{34}x_6 \\ &= (a_{22}a_{33} + a_{21}a_{34})x_5 + (a_{21}a_{33} + a_{22}a_{34} + ta_{22}a_{33})x_6 = y_5 + sy_6. \end{aligned}$$

Let $p = a_{22}a_{33} + a_{21}a_{34}$, $q = a_{21}a_{33} + a_{22}a_{34} + ta_{22}a_{33}$. In view of (3.1), (3.5)

$$(a_{22}y_5 - a_{12}y_6)p/\Delta^2 + (a_{11}y_6 - a_{21}y_5)q/\Delta^2 = y_5 + sy_6.$$

This implies that

$$(3.6) \quad \begin{cases} qa_{11} - pa_{12} = s\Delta^2, \\ qa_{21} - pa_{22} = -\Delta^2. \end{cases}$$

Consequently,

$$(3.7) \quad \begin{cases} p = (sa_{21} + a_{11})\Delta, \\ q = (sa_{22} + a_{12})\Delta. \end{cases}$$

Substituting u, v, p, q by the expressions given in (3.4), (3.7) we conclude that

$$(3.8) \quad \begin{cases} a_{11}a_{34} + a_{12}a_{33} = a_{21}\Delta, \\ a_{11}a_{33} + a_{12}(a_{34} + ta_{33}) = a_{22}\Delta, \\ a_{21}a_{34} + a_{22}a_{33} = (a_{11} + sa_{21})\Delta, \\ a_{21}a_{33} + a_{22}(a_{34} + ta_{33}) = (a_{12} + sa_{22})\Delta. \end{cases}$$

The first and the third equations of (3.8) yield

$$(3.9) \quad \begin{cases} a_{34} = a_{21}a_{22} - a_{12}(a_{11} + sa_{21}), \\ a_{33} = a_{11}(a_{11} + sa_{21}) - a_{21}a_{22}. \end{cases}$$

The two remaining ones yield

$$(3.10) \quad \begin{cases} a_{33} = a_{22}a_{22} - a_{12}(a_{12} + sa_{22}), \\ a_{34} + ta_{33} = a_{11}(a_{12} + sa_{22}) - a_{21}a_{21}, \end{cases}$$

whence

$$(3.11) \quad \begin{cases} a_{11}^2 + sa_{11}a_{21} - a_{21}^2 = a_{22}^2 - sa_{22}a_{12} - a_{12}^2 \neq 0, \\ 2(a_{11}a_{12} - a_{21}a_{22}) + s(a_{11}a_{22} - a_{21}a_{12}) = t(a_{11}^2 + sa_{11}a_{21} - a_{21}^2). \end{cases}$$

Let

$$\begin{cases} x_{11} = a_{11} + sa_{21}/2, \\ x_{12} = a_{12} + sa_{22}/2, \\ x_{21} = a_{21}, \\ x_{22} = a_{22}. \end{cases}$$

The system (3.11) can be rewritten in the form

$$(3.12) \quad \begin{cases} x_{11}^2 - (1 + s^2/4)x_{21}^2 = -(x_{12}^2 - (1 + s^2/4)x_{22}^2) \neq 0, \\ 2(x_{11}x_{12} - (1 + s^2/4)x_{21}x_{22}) = t(x_{11}^2 - (1 + s^2/4)x_{21}^2). \end{cases}$$

Thus we may conclude that $\mathfrak{g}_t \cong \mathfrak{g}_s$ if and only if (3.12) has a rational solution such that $x_{11}x_{22} - x_{12}x_{21} \neq 0$. We state the following lemma in order to obtain less sophisticated conditions on s, t .

LEMMA 3.1. *Let $s, t \in \mathbf{Q}$. Then two conditions are equivalent:*

i) *there exists a matrix*

$$M = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in \mathrm{GL}_2(\mathbf{Q})$$

such that x, y, z, w satisfy the system

$$(3.13) \quad \begin{cases} x^2 - (1 + s^2/4)z^2 = -y^2 + (1 + s^2/4)w^2 \neq 0, \\ 2(xy - (1 + s^2/4)zw) = t(x^2 - (1 + s^2/4)z^2). \end{cases}$$

ii) *there exists $q \in \mathbf{Q}$ such that*

$$(3.14) \quad (t^2 + 4)(s^2 + 4) = q^2.$$

Proof. Let $p = 1 + s^2/4$, $r = 1 + t^2/4$. The system (3.13) yields

$$(3.15) \quad \begin{cases} x^2 + y^2 = p(z^2 + w^2), \\ 2xy - tx^2 = p(2zw - tz^2). \end{cases}$$

After the change of variables

$$x = x_0, \quad y = \frac{1}{2}(y_0 + tx_0), \quad z = \frac{z_0}{p}, \quad w = \frac{1}{2}(w_0 + \frac{t}{p}z_0)$$

the system (3.15) can be rewritten as

$$(3.16) \quad \begin{cases} rx_0^2 + \frac{1}{4}y_0^2 = \frac{r}{p}z_0^2 + \frac{p}{4}w_0^2, \\ x_0y_0 = z_0w_0. \end{cases}$$

Geometrically, the system (3.16) defines the intersection I of two quadrics in the projective space \mathbf{P}^3 . Let $\sigma: \mathbf{P}^1 \times \mathbf{P}^1 \rightarrow \mathbf{P}^3$ be the Segre map. In homogeneous coordinates $(a : b ; \alpha : \beta)$ in $\mathbf{P}^1 \times \mathbf{P}^1$, σ is defined by $x_0 = a\alpha$, $y_0 = b\beta$, $z_0 = a\beta$, $w_0 = b\alpha$, and the image $\sigma(\mathbf{P}^1 \times \mathbf{P}^1)$ is the zero locus of the polynomial $x_0y_0 - z_0w_0$.

It is not hard to verify that in coordinates $(a : b ; \alpha : \beta)$ the preimage $\sigma^{-1}(I)$ is given by the following equation (corresponding to the first one of (3.16)):

$$(4ra^2 - pb^2)(p\alpha^2 - \beta^2) = 0.$$

Thus $\sigma^{-1}(I)$ is the union of two pairs of lines (over \mathbf{R}). The second pair defined by the equation $p\alpha^2 - \beta^2 = 0$ yields $xw - zy = \det(M) = 0$. It follows that (3.15) has a rational solution if and only if the equation $4ra^2 - pb^2 = 0$ has one, i.e., $p/4r$ is the square of a rational number. This is equivalent to (3.14). Note that the condition $x^2 - pz^2 \neq 0$ in (3.13) is not very restrictive. This completes the proof of the lemma and of Theorem 1.

COROLLARY 3.2. *There are infinitely many non-isomorphic Lie algebras of the type \mathfrak{g}_s over \mathbf{Q} .*

Proof. Let $s_1 = p_{11}$ be an odd prime. Consider $s_1^2 + 4 = p_1^2 + 4 = p_{21}^{n_{21}} \dots$ It is clear that $s_1^2 + 4$ is not a square (this means that at least one of the n_{2j} is odd) and is not divisible by p_{11} , whence all the $p_{2j} \neq p_{11}$. Let $s_2 = p_{11}p_{21} \dots$ It follows that

$$s_2^2 + 4 = p_{31}^{n_{31}} \dots$$

is not a square and is not divisible by p_{ij} where $i \leq 2$. Then we set $s_3 = p_{11}p_{21} \dots p_{31} \dots$ and so on.

In such a way we obtain an infinite sequence of numbers $s_1, s_2 \dots$ Let $i < j$. Note that $(s_i^2 + 4)(s_j^2 + 4) \neq q^2$, $q \in \mathbf{Q}$. Indeed, by the construction $(s_i^2 + 4)$ is divisible by some p and not divisible by p^2 . Also, p divides s_j . Consequently, it does not divide $s_j^2 + 4$. This means that $(s_i^2 + 4)(s_j^2 + 4)$ is divisible by p but not by p^2 , and this completes the proof.