Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 48 (2002)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MM-SPACES AND GROUP ACTIONS

Autor: Pestov, Vladimir

Kapitel: 2. Some concepts of asymptotic geometric analysis

DOI: https://doi.org/10.5169/seals-66074

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

theory, cf. papers by Gromov [Gr1], Milman [M2,M3], and some others [A-M,G1,P2,P3,G-P,G1-W]. However, it is safe to say that there is still a long way to go towards the full understanding of the picture.

Here we aim at providing a readable introduction into this circle of ideas.

2. Some concepts of asymptotic geometric analysis

DEFINITION 1. A space with metric and measure, or an mm-space, is a triple (X, d, μ) , where d is a metric on a set X and μ is a finite Borel measure on the metric space (X, d). It will be convenient to assume throughout that μ is a probability measure, that is, normalized to one.

DEFINITION 2. The concentration function α_X of an mm-space $X=(X,d,\mu)$ is defined for non-negative real ε as follows:

$$\alpha_X(\varepsilon) = \begin{cases} \frac{1}{2} & \text{if } \varepsilon = 0, \\ 1 - \inf\{\mu(A_{\varepsilon}) \colon A \subseteq X \text{ is Borel, } \mu(A) \ge \frac{1}{2} \} & \text{if } \varepsilon > 0. \end{cases}$$

Here A_{ε} denotes the ε -neighbourhood (ε -fattening, ε -thickening) of A.

EXERCISE 1. Prove that $\alpha(\varepsilon) \to 0$ as $\varepsilon \to \infty$. (For spaces of finite diameter this is of course obvious.)

DEFINITION 3. An infinite family of mm-spaces, $(X_n, d_n, \mu_n)_{n=1}^{\infty}$, is called a *Lévy family* if the concentration functions α_n of X_n converge to zero pointwise on $(0, \infty)$:

$$\forall \varepsilon > 0, \ \alpha_n(\varepsilon) \to 0 \text{ as } n \to \infty.$$

EXERCISE 2. Prove that the above condition is equivalent to the following. Let $A_n \subseteq X_n$ be Borel subsets with the property that

$$\liminf_{n\to\infty}\mu_n(A_n)>0.$$

Then

$$\forall \varepsilon > 0, \lim_{n \to \infty} \mu_n((A_n)_{\varepsilon}) = 1.$$

The following are some of the most common examples of Lévy families.

EXAMPLE 1. Unit spheres S^n in the Euclidean spaces R^{n+1} , equipped with the Euclidean (or geodesic) distances and the normalized Haar measures (that is, the unique rotation-invariant probability measures). This result is due to Paul Lévy [Lév], though his proof, based on the isoperimetric inequality, was only made rigorous much later by Gromov [Gr2]. (Nowadays simpler proofs, using the Brunn–Minkowski inequality, are known, cf. [Gr-M2, Sch].)

EXAMPLE 2. The special orthogonal groups SO(n), equipped with the normalized Haar measure and the uniform operator metric,

$$d(T,S) := ||T - S||,$$

induced from $\mathcal{B}(\mathbf{R}^n) \cong M_n$. This was established by Gromov and Milman [Gr-M1]. The same argument holds for the special unitary groups.

EXAMPLE 3. The family of finite permutation groups (S_n) , equipped with the uniform (normalized counting) measure and the Hamming distance:

$$d(\sigma,\tau) = \frac{1}{n} |\{i \colon \sigma(i) \neq \tau(i)\}|.$$

The result is due to Maurey [Ma], see also [Ta1].

EXAMPLE 4. The Hamming cubes $\{0,1\}^n$ equipped with the normalized counting measure and the Hamming distance $d(x,y) = \frac{1}{n} |\{i : x_i \neq y_i\}|$ form a Lévy family [Sch,M-S].

REMARK 1. All of the above are *normal* Lévy families, meaning that the concentration functions α_n admit Gaussian upper bounds:

$$\alpha_n(\varepsilon) \leq C_1 \exp(-C_2 n \varepsilon^2)$$

for some $C_1, C_2 > 0$.

It should be noted that this is not always the case for 'naturally occurring' Lévy families. For instance, the groups $SL(2, \mathbf{F}_p)$, where p are prime numbers, equipped with the normalized counting measure and the word metric given by a fixed system of generators in $SL(2, \mathbf{Z})$, form a Lévy family with $\alpha_p(\varepsilon) \leq C_1 \exp(-C_2\sqrt{p}\,\varepsilon)$, [A-M, M4]. (Recall in this connection that the n-th prime number $p_n \sim n \log n$.)

REMARK 2. In Example 4, replace $\{0,1\}$ with any probability measure space, $X=(X,\mu)$. Equip every finite power X^n with the product measure $\mu^{\otimes n}$ and the normalized Hamming distance $d(x,y)=\frac{1}{n}|\{i\colon x_i\neq y_i\}|$. Unless X is purely atomic, the measures $\mu^{\otimes n}$ are not Borel, and thus X^n aren't even mm-spaces in the sense of our definition. At the same time, if in the definition of the concentration function we only restrict ourselves to measurable subsets A such that A_{ε} are also measurable, it can be shown that $X^n, n \in \mathbb{N}$ form a Lévy family in a very reasonable sense. (See [Ta1,Ta3] for far-reaching variations.) If anything, this shows that the full formalization of the subject has not yet been achieved and nothing is cast in stone.

Notice that the *mm*-spaces from the above Examples 1–4 are at the same time (phase spaces of) topological transformation groups, with both metrics and measures being invariant under group actions. In Example 1 it is the action of the orthogonal — or the unitary — group on the sphere, while in Examples 2–4 the groups act upon themselves on the left.

3. A TRANSFORMATION GROUP FRAMEWORK

Here is the idea of what kind of interaction between concentration phenomenon and group actions one should expect. The following example is borrowed from a paper by Vitali Milman [M4].

Suppose a group G acts on an mm-space (X,d,μ) by measure-preserving isometries. Assume that the mm-space X strongly concentrates, that is, the function $\alpha_X(\varepsilon)$ drops off sharply already for small values of ε . Let us assume, for instance, that the concentration is so strong that, whenever $\mu(A) \geq \frac{1}{7}$, the measure of the $\frac{1}{10}$ -neighbourhood of A is strictly greater than 0.99. (Cf. Exercise 2.)

If now we partition X into seven pieces, and pick at random one hundred elements $g_1,g_2,\ldots,g_{100}\in G$, then at least one of the pieces, say A, has the property that all one hundred translates, of $\frac{1}{10}$ -neighbourhoods of A by our elements g_i have a point, x^* , in common. Equivalently, x^* is 'close' (closer than $\frac{1}{10}$) to each of the one hundred translates of A.

The above effect becomes more pronounced the higher the level of concentration is. Partition a concentrated ('high-dimensional') *mm*-space into a small number of subsets, and at least one of them is hard to move.

In order to set up a formal framework, we assume all topological spaces and topological groups appearing in this article to be metrizable, for the reasons