

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MM-SPACES AND GROUP ACTIONS
Autor: Pestov, Vladimir
Kapitel: 6.5 The Urysohn metric space
DOI: <https://doi.org/10.5169/seals-66074>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

EXERCISE 11. Prove that the action of S_∞ on LO is continuous and minimal (that is, the orbit of each linear order is everywhere dense in LO).

Recall that a linear order \prec is called *dense* if it has no gaps. A dense linear order without least and greatest elements is said to be of type η . The collection LO_η of all linear orders of type η on \mathbf{Z} can be identified with the factor space $S_\infty / \text{Aut}(\prec)$ through the correspondence $\sigma \mapsto \sigma \prec$. Here \prec is some chosen linear order of type η on \mathbf{Z} and $\text{Aut}(\prec)$ stands for the group of order-preserving self-bijections of (\mathbf{Z}, \prec) , acting on the space of orders in a natural way: $(x \sigma \prec y) \Leftrightarrow \sigma^{-1}x \prec \sigma^{-1}y$.

EXERCISE 12. Show that under the above identification the uniform structure on LO_η , induced from the compact space LO, is the finest uniform structure making the quotient map $S_\infty \rightarrow S_\infty / \text{Aut}(\prec) \cong LO_\eta$ right uniformly continuous.

Let now X be a compact S_∞ -space. The topological subgroup $\text{Aut}(\prec)$ of S_∞ has a fixed point in X , say x' (Exercise 10). The mapping $S_\infty \ni \sigma \mapsto \sigma(x') \in X$ is constant on the left $\text{Aut}(\prec)$ -cosets and thus gives rise to a mapping $\varphi: LO_\eta \rightarrow X$. Using Exercise 12, it is easy to see that φ is right uniformly continuous and thus extends to a morphism of S_∞ -spaces $LO \rightarrow X$. We have established the following result.

THEOREM 6 (Glasner and Weiss [Gl-W]). *The compact space LO forms the universal minimal S_∞ -space.*

6.5 THE URYSOHN METRIC SPACE

The *universal Urysohn metric space* \mathbf{U} [Ur] is determined uniquely (up to an isometry) by the following conditions:

- (i) \mathbf{U} is a complete separable metric space;
- (ii) \mathbf{U} is ω -homogeneous, that is, every isometry between two finite subspaces of \mathbf{U} extends to an isometry of \mathbf{U} ;
- (iii) \mathbf{U} contains an isometric copy of every separable metric space.

A probabilistic description of this space was given by Vershik [Ver]: the completion of the space of integers equipped with a ‘sufficiently random’ metric is almost surely isometric to \mathbf{U} .

The group of isometries $\text{Iso}(\mathbf{U})$ with the compact-open topology is a Polish (complete metric separable) topological group, which also possesses

a universality property: it contains an isomorphic copy of every separable metric group [Usp]. See also [Gr3].

Using concentration of measure, one can prove that the group $\text{Iso}(\mathbf{U})$ is extremely amenable. The Ramsey–Dvoretzky–Milman property leads to the following Ramsey-type result:

Let F be a finite metric space, and let all isometric embeddings of F into \mathbf{U} be coloured using finitely many colours. Then for every finite metric space G and every $\varepsilon > 0$ there is an isometric copy $G' \subset \mathbf{U}$ of G such that all isometric embeddings of F into \mathbf{U} that factor through G are monochromatic to within ε .




FIGURE 5
A Ramsey-type result for metric spaces

Here we say that a set A is *monochromatic to within ε* if there is a monochromatic set A' at a Hausdorff distance $< \varepsilon$ from A . In our case, the Hausdorff distance is formed with regard to the uniform metric on \mathbf{U}^F .

One can also obtain similar results, for example, for the separable Hilbert space ℓ_2 and for the unit sphere S^∞ in ℓ_2 [P3].

7. CONCENTRATION TO A NON-TRIVIAL SPACE

Let f be a Borel measurable real-valued function on an mm -space $X = (X, d, \mu)$. A number $M = M_f$ is called a *median* (or *Lévy mean*) of f if both $f^{-1}[M, +\infty)$ and $f^{-1}(-\infty, M]$ have measure $\geq \frac{1}{2}$.

EXERCISE 13. Show that the median M_f always exists, though it need not be unique.