

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	48 (2002)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF $\sharp_{i=1}^b (S^2 \times S^5)$
Autor:	Schmitt, Alexander
Kapitel:	2.2 Manifolds with trivial cup form δ_x
DOI:	https://doi.org/10.5169/seals-66077

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2.2 MANIFOLDS WITH TRIVIAL CUP FORM δ_X

In addition to describing the explicit geometric meaning of the system of invariants of an E-manifold X with $w_2(X) = 0$, we will describe those manifolds X for which the cup form δ_X vanishes.

For any $b > 0$, let $\mathcal{FC}_b^{\text{PL}(\mathcal{C}^\infty)}$ be the group of isotopy classes of piecewise linear (smooth) embeddings of b disjoint copies of $S^5 \times D^3$ into S^8 . The following result will be established in Section 3.7.

PROPOSITION 2.3. $\text{FL}_b := \mathcal{FC}_b^{\mathcal{C}^\infty} \cong \mathcal{FC}_b^{\text{PL}}$.

Given an element $[l] \in \text{FL}_b$, we can perform surgery along the link l and get a smooth based E-manifold $(X(l), \underline{x}(l))$ with $w_2(X) = 0$ and $b_4(X) = 0$, which is well defined up to smooth isomorphy of based manifolds.

We will also use the following notation: Fix a pair $(\gamma, p) \in Z(0, b')$ which satisfies relation (2) (and (3)) and denote by $\mathcal{I}^{\text{PL}(\mathcal{C}^\infty)}(b, \gamma, p)$ the set of isomorphy classes of based piecewise linear (smooth) E-manifolds $(X, \underline{x}, \underline{y})$ with $w_2(X) = 0$, $b_2(X) = b$, $\gamma_X = \gamma$, and $p_1(X) = p$. Pick a three-connected piecewise linear (smooth) based E-manifold (X_0, \underline{y}_0) with $\gamma_{X_0} = \gamma$ and $p_1(X_0) = p$. In the smooth case, let $\vartheta^8 \cong \mathbf{Z}_2$ [17] be the group of exotic 8-spheres, and set $\vartheta(X_0) := \vartheta^8$, if X_0 is not diffeomorphic to $X_0 \# \Sigma$, Σ a generator for ϑ^8 , and $\vartheta(X_0) := \{[S^8]\} \subset \vartheta^8$ otherwise. Now, we define maps

$$\begin{aligned} K^{\text{PL}}(b, \gamma, p): \text{FL}_b &\longrightarrow \mathcal{I}^{\text{PL}}(b, \gamma, p) \\ [l] &\longmapsto \left[X(l) \# X_0, \underline{x}(l), \underline{y}_0 \right] \end{aligned}$$

and

$$\begin{aligned} K_{X_0}^{\mathcal{C}^\infty}(b, \gamma, p): \text{FL}_b \oplus \vartheta(X_0) &\longrightarrow \mathcal{I}^{\mathcal{C}^\infty}(b, \gamma, p) \\ ([l], [\Sigma]) &\longmapsto \left[X(l) \# X_0 \# \Sigma, \underline{x}(l), \underline{y}_0 \right]. \end{aligned}$$

In $\mathcal{I}^{\text{PL}(\mathcal{C}^\infty)}(b, \gamma, p)$, we mark the class $\left[(\#_{i=1}^b (S^2 \times S^6)) \# X_0, \underline{x}, \underline{y}_0 \right]$, where \underline{x} comes from the natural basis of $H^2(\#_{i=1}^b (S^2 \times S^6), \mathbf{Z})$. Then our main result is the following

THEOREM 2.4. i) For every $b > 0$ and every pair (γ, p) which satisfies the relation (2),

$$\begin{aligned} \{1\} &\longrightarrow \text{FL}_b & K^{\text{PL}}(b, \gamma, p) &\longrightarrow \text{Hom}(S^2 \mathbf{Z}^b, Z^{b'}) \\ 1 &\longmapsto [\text{trivial link}] & [X, \underline{x}, \underline{y}] &\longmapsto \delta_X \end{aligned}$$

is an exact sequence of pointed sets.

ii) For every $b > 0$ and every pair (γ, p) which satisfies the relations (2) and (3),

$$\begin{aligned} \{1\} &\longrightarrow \text{FL}_b \oplus \vartheta(X_0) & K_{X_0}^{\mathcal{C}^\infty}(b, \gamma, p) &\longrightarrow \text{Hom}(S^2 \mathbf{Z}^b, Z^{b'}) \\ 1 &\longmapsto [\text{trivial link}] & [X, \underline{x}, \underline{y}] &\longmapsto \delta_X \end{aligned}$$

is an exact sequence of pointed sets.

The proof will be given in Sections 4.2 and 4.3.

REMARK 2.5. i) In the PL setting, we will show that the inclusion of FL_b into $\mathfrak{I}^{\text{PL}}(b, \gamma, \varphi)$ extends to an action of FL_b on $\mathfrak{I}^{\text{PL}}(b, \gamma, \varphi)$, such that the orbits are the fibres of the map $[X, \underline{x}, \underline{y}] \longmapsto \delta_X$.

ii) On all the sets occurring in Theorem 2.4 there are natural $(\text{GL}_b(\mathbf{Z}) \times \text{GL}_{b'}(\mathbf{Z}))$ -actions, and the maps are equivariant for these actions. Therefore, by forming the equivalence classes with respect to these actions, we get the classification of E-manifolds with vanishing second Stiefel-Whitney class up to orientation preserving piecewise linear (smooth) isomorphy.

iii) We will discuss in Section 3.7 the structure of the group FL_b . It turns out that it is finite if and only if $b = 1$. It follows easily that the set of $\text{GL}_b(\mathbf{Z})$ -equivalence classes in FL_b is infinite for $b \geq 2$. Thus, the cohomology ring and the characteristic classes classify E-manifolds of dimension eight up to finite indeterminacy only if the second Betti number is at most one.

The starting point of our proof of the above results will be the theory of minimal handle decompositions of Smale which states that X can be obtained from D^8 by first attaching $b_2(X)$ 2-handles, then $b_4(X)$ 4-handles, then $b_2(X)$ 6-handles and finally one 8-handle. At each step, the attachment will be determined by the isotopy class of a certain framed link in a 7-manifold, and we will first explain how to read off the isotopy class of the attaching links for the 2- and 4-handles from the invariants.