

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF  $S^1 \times S^5$   
**Autor:** Schmitt, Alexander  
**Kapitel:** 3.3 Homotopy vs. isotopy  
**DOI:** <https://doi.org/10.5169/seals-66077>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

iii) If  $X$  comes with an orientation, we may assume that  $D^m$  is orientation preservingly embedded and that all attaching maps are orientation reversing.

### 3.2 CONSEQUENCES FOR E-MANIFOLDS OF DIMENSION EIGHT WITH $w_2 = 0$

Let  $X$  be a piecewise linear (smooth) E-manifold of dimension eight with  $w_2(X) = 0$ . The first observation concerns the structure of  $W_2$ .

LEMMA 3.5. *One has  $W_2 \cong \#_{i=1}^b(S^2 \times D^6)$ .*

*Proof.* The manifold  $W_2$  is an  $(8, 1)$ -handle body and as such homeomorphic to the boundary connected sum of  $b$   $D^6$ -bundles over  $S^2$  ([17], §11, p.115). As  $\pi_1(\text{SO}(4)) \cong \mathbf{Z}_2$  and we have requested  $w_2(X) = 0$ , the claim follows.  $\square$

The next consequence is

*The manifold  $W_4$  is determined by a framed link of  $b_4(X)$  three-dimensional spheres in  $\#_{i=1}^b(S^2 \times S^5)$ .*

We shall look into the classification of such links below. The third consequence is

LEMMA 3.6. i)  $\partial W_4 \cong \#_{i=1}^b(S^2 \times S^5)$ .

ii) *The manifold  $X$  is of the form  $W_4 \cup_f \#_{i=1}^b(S^2 \times D^6)$  where*

$$f: \#_{i=1}^b(S^2 \times S^5) \longrightarrow \partial W_4$$

*is a piecewise linear (smooth) isomorphism, such that  $f_*$  maps the canonical basis of  $H_2(\#_{i=1}^b(S^2 \times S^5), \mathbf{Z})$  to the preferred basis of  $H_2(\partial W_4, \mathbf{Z})$ .*

*Proof.* i) This follows because  $\partial W_4 \cong \partial \bar{W}_2$ . ii) follows because  $X = W_4 \cup \bar{W}_2$ , and  $\bar{W}_2 \cong \#_{i=1}^b(S^2 \times D^6)$ , by Lemma 3.5.  $\square$

### 3.3 HOMOTOPY VS. ISOTOPY

By Theorem 3.3, the manifold is determined by the ambient isotopy classes of the attaching maps. However, the topological invariants of the manifold give us at best their homotopy classes. It is, therefore, important to have theorems granting that this is enough. In the setting of differentiable manifolds, we have

**THEOREM 3.7** (Haefliger [6], [7]). *Let  $S$  be a closed differentiable manifold of dimension  $n$  and  $M$  an  $m$ -dimensional differentiable manifold without boundary. Let  $f: S \rightarrow M$  be a continuous map and  $k \geq 0$ , such that*

$$\pi_i(f): \pi_i(S) \rightarrow \pi_i(M)$$

*is an isomorphism for  $0 \leq i \leq k$  and surjective for  $i = k + 1$ . Then the following is satisfied:*

1. *If  $m \geq 2n - k$  and  $n > 2k + 2$ , then  $f$  is homotopic to a differentiable embedding.*
2. *If  $m > 2n - k$  and  $n \geq 2k + 2$ , then two differentiable embeddings of  $S$  into  $M$  which are homotopic are also ambient isotopic.*

In the setting of piecewise linear manifolds, similar results hold true. We refer to Haefliger's survey article [9]. For our purposes, the result stated below will be sufficient.

**THEOREM 3.8.** *Suppose  $S$  is a closed  $n$ -dimensional manifold,  $M$  an  $m$ -dimensional manifold without boundary, and  $f: S \rightarrow M$  a continuous map. Assume one has*

- $m - n \geq 3$ ;
- $S$  is  $(2n - m + 1)$ -connected;
- $M$  is  $(2n - m + 2)$ -connected.

*Then:*

- (1)  *$f$  is homotopic to an embedding;*
- (2) *two embeddings which are homotopic to  $f$  are ambient isotopic.*

*Proof.* The theorem of Irwin ([27], Thm. 7.12 and Ex. 7.14, [13], Thm. 8.1) yields (1) and that  $f_1$  and  $f_2$  as in (2) are concordant. But, since  $m - n \geq 3$ , concordance implies ambient isotopy ([13], Chap. IX).  $\square$

**COROLLARY 3.9.** *Let  $S := S^3$  and  $M$  a simply connected differentiable or piecewise linear manifold of dimension 7 without boundary. Then  $\pi_3(M)$  classifies differentiable and piecewise linear embeddings, respectively, of  $S^3$  into  $M$  up to ambient isotopy.*

**COROLLARY 3.10** (Zeeman's unknotting theorem [39]). *For every  $m, n$  with  $m - n \geq 3$ , any piecewise linear embedding of  $S^n$  into  $S^m$  is isotopic to the standard embedding.*