

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF  $\sharp_{i=1}^b (S^2 \times S^5)$   
**Autor:** Schmitt, Alexander  
**Kapitel:** 3.6 Links of 3-spheres in  $\sharp_{i=1}^b (S^2 \times S^5)$   
**DOI:** <https://doi.org/10.5169/seals-66077>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

### 3.5 PONTRJAGIN CLASSES AND $\pi_3(\text{SO}(4))$

Vector bundles of rank 4 over  $S^4$  are classified by elements in  $\pi_3(\text{SO}(4))$ . In our setting, such vector bundles will appear as normal bundles. We recall, therefore, the description of that group and relate it to Pontrjagin classes and self intersection numbers.

First, look at the natural map  $\pi_3(\text{SO}(4)) \longrightarrow \pi_3(\text{SO}(4)/\text{SO}(3)) = \pi_3(S^3)$ . This map has a splitting ([32], §22.6) which induces an isomorphism

$$\pi_3(\text{SO}(4)) = \pi_3(\text{SO}(3)) \oplus \pi_3(S^3).$$

Let  $\alpha_3$  be the generator for  $\pi_3(\text{SO}(3)) \cong \mathbf{Z}$  from [32], §22.3, and  $\beta_3 := [\text{id}_{S^3}] \in \pi_3(S^3)$ , so that we obtain the isomorphism  $\mathbf{Z} \oplus \mathbf{Z} \longrightarrow \pi_3(\text{SO}(4))$ ,  $(k_1, k_2) \longmapsto k_1\alpha_3 + k_2\beta_3$ . Finally, the kernel of the map  $\pi_3(\text{SO}(4)) \longrightarrow \pi_3(\text{SO})$  to the stable homotopy group is generated by  $-\alpha_3 + 2\beta_3$  ([32], §23.6), whence [23], (20.9), implies

**PROPOSITION 3.13.** *Let  $E$  be the vector bundle over  $S^4$  defined by the element  $k_1\alpha_3 + k_2\beta_3 \in \pi_3(\text{SO}(4))$ . Then*

$$p_1(E) = \pm(2k_1 + 4k_2).$$

**COROLLARY 3.14.** *Let  $f: S^4 \longrightarrow M$  be a differentiable embedding of  $S^4$  into the differentiable 8-manifold  $M$ . Let  $E := f^*T_M/T_{S^4}$  be the normal bundle. Then the self intersection number  $s$  of  $f(S^4)$  in  $M$  satisfies*

$$2s \equiv p_1(E) \pmod{4}.$$

*Proof.* If  $E$  is given by the element  $k_1\alpha_3 + k_2\beta_3 \in \pi_3(\text{SO}(4))$ , then  $s = k_2$  ([17], (5.4), p.72). Since  $p_1(E) = \pm(2k_2 + 4k_1)$ , the claim follows.  $\square$

### 3.6 LINKS OF 3-SPHERES IN $\#_{i=1}^b(S^2 \times S^5)$

If  $X$  is a closed E-manifold of dimension 8 with  $w_2(X) = 0$ , then  $W_2 := \#_{i=1}^b(S^2 \times D^6)$ ,  $b = b_2(X)$ , by Lemma 3.5. Thus,  $W_4$  is determined by a framed link of 3-spheres in  $\partial W_2 = \#_{i=1}^b(S^2 \times S^5)$ . Therefore, we will now classify such links.

So, let  $W := \#_{i=1}^b(S^2 \times S^5)$  be a  $b$ -fold connected sum. We can choose  $b$  disjoint 2-spheres  $S_i^2$ ,  $i = 1, \dots, b$ , embedded in  $W$  and representing the natural basis of  $H_2(W, \mathbf{Z})$ . One checks that the homotopy type of  $W$  is given up to dimension 4 by the  $b$ -fold wedge product  $S^2 \vee \dots \vee S^2$ . Suppose we are given a link of  $b'$  three-dimensional spheres, i.e., we are given  $b'$  differentiable embeddings  $g_i: S^3 \longrightarrow W$ ,  $i = 1, \dots, b'$ , with  $g_i(S^3) \cap g_j(S^3) = \emptyset$  for  $i \neq j$ .

By the transversality theorem ([17], IV.(2.4)), one sees that we may assume  $S_i^2 \cap g_j(S^3) = \emptyset$  for all  $i$  and  $j$ .

By Corollary 3.9, the ambient isotopy class of the embedding  $g_k$  is determined by the element  $\varphi_k := [g_k] \in \pi_3(W_k)$ ,  $W_k := W \setminus \bigcup_{j \neq k} g_j(S^3)$ ,  $k = 1, \dots, b'$ . We clearly have (compare [8])

$$\pi_3(W_k) = \pi_3 \left( \underbrace{S^2 \vee \dots \vee S^2}_{b \times} \vee \underbrace{S^3 \vee \dots \vee S^3}_{(b'-1) \times} \right),$$

so that the Hilton-Milnor theorem yields

$$\pi_3(W_k) = \bigoplus_{i=1}^b \pi_3(S^2) \oplus \bigoplus_{1 \leq i < j \leq b} \pi_3(S^3) \oplus \bigoplus_{j \neq k} \pi_3(S^3).$$

Hence, we write  $\varphi_k$  as a tuple of integers:

$$\varphi_k = (l_i^k, i = 1, \dots, b; l_{ij}^k, 1 \leq i < j \leq b; \lambda_{kj}, j \neq k).$$

Observe that, for  $j \neq k$ ,  $\varphi_k$  is mapped under the natural homomorphism

$$\pi_3(W_k) \longrightarrow H_3(W_k, \mathbf{Z}) \longrightarrow H_3(W \setminus g_j(S^3), \mathbf{Z}) (\cong \mathbf{Z})$$

to the image of the fundamental class of  $S^3$  under  $g_{j*}$ . Thus,  $\lambda_{kj}$  is just the ‘usual’ linking number of the spheres  $g_k(S^3)$  and  $g_j(S^3)$  in  $W$  (compare [8]).

### 3.7 LINKS OF 5-SPHERES IN $S^8$

Let  $\mathcal{FC}_b^{\text{PL}(C^\infty)}$  be as before, and let  $\mathcal{C}_b^{\text{PL}(C^\infty)}$  be the group of isotopy classes of piecewise linear (smooth) embeddings of  $b$  disjoint copies of  $S^5$  into  $S^8$ . For  $b = 1$ , these groups are studied in [10], [19], and [20]. A brief summary with references of results in the case  $b > 1$  is contained in Section 2.6 of [11]. We will review some of this material below.

PROPOSITION 3.15. *We have  $\mathcal{FC}_1^{C^\infty} \cong \mathcal{FC}_1^{\text{PL}} \cong \mathbf{Z}_2$ .*

*Proof.* Since  $\pi_5(\text{SO}(3)) \cong \mathbf{Z}_2$ , the standard embedding of  $S^5$  into  $S^8$  with its two possible framings provides an injection of  $\mathbf{Z}_2$  into  $\mathcal{FC}_1^{\text{PL}(C^\infty)}$ . By Zeeman’s unknotting theorem 3.10, the map  $\mathbf{Z}_2 \longrightarrow \mathcal{FC}_1^{\text{PL}}$  is an isomorphism. As remarked in Section 2.6 of [11],  $\mathcal{FC}_1^{\text{PL}}$  is isomorphic to  $\mathcal{F}\vartheta$ , the group of h-cobordism classes of framed submanifolds of  $S^8$  which are homotopy 5-spheres. Moreover, by [10] and [19], there is an exact sequence

$$\dots \longrightarrow \vartheta^6 \longrightarrow \mathcal{FC}_1^{C^\infty} \longrightarrow \mathcal{F}\vartheta \longrightarrow \vartheta^5 \longrightarrow \dots$$

As the groups  $\vartheta^5$  and  $\vartheta^6$  of exotic 5- and 6-spheres are trivial [17], our claim is settled.  $\square$