

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF $\sharp_{i=1}^b (S^2 \times S^5)$
Autor: Schmitt, Alexander
Kapitel: 4. Proof of Theorem 2.2 and Theorem 2.4
DOI: <https://doi.org/10.5169/seals-66077>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. PROOF OF THEOREM 2.2 AND THEOREM 2.4

From now on, X stands for an eight-dimensional E-manifold with $w_2(X) = 0$.

4.1 PROOF OF THEOREM 2.2

The classification result for 3-connected E-manifolds of dimension eight is a special case of a result of Wall's [36] and can be easily obtained with the methods described in [17], VII, §12. Let us recall the details, because we will need them later on.

We fix a basis \underline{b} for $H_4(X, \mathbf{Z})$ and let \underline{y} be the dual basis of $H^4(X, \mathbf{Z})$. Then there is a handle presentation $X = D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4 \cup D^8$ with \underline{b} as the preferred basis. The manifold $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$ is determined by the ambient isotopy class of a framed link of 3-spheres in S^7 , having b' components. Let us first look at such a link, forgetting the framing, i.e., suppose we are given embeddings $g_i: S^3 \rightarrow S^7$ with $S_i \cap S_j = \emptyset$ for $i \neq j$, $S_i := g_i(S^3)$, $i = 1, \dots, b'$. By 3.9, we may assume that the g_i are differentiable. Observe that the normal bundles of the S_i are trivial.

We equip S_i with the orientation induced via g_i by the standard orientation of S^3 and the normal bundle of S_i with the orientation which is determined by requiring that the orientation of S_i followed by that of its normal bundle coincide with the orientation of S^7 . Therefore, a 3-sphere F_i which bounds the fibre of a tubular neighborhood of S_i in S^7 inherits an orientation and thus provides a generator e_i for $H_3(S^7 \setminus S_i, \mathbf{Z}) \cong \mathbf{Z}$, $i = 1, \dots, b'$. For $i \neq j$, the image of the fundamental class $[S_i]$ in $H_3(S^7 \setminus S_j, \mathbf{Z})$ is of the form $\lambda_{ij} \cdot e_j$. The integer λ_{ij} is called *the linking number of S_i and S_j* .

For $i = 1, \dots, b'$, the manifold $S^7 \setminus \bigcup_{j \neq i} S_j$ is up to dimension 5 homotopy equivalent to $\bigvee_{j \neq i} F_j$, and

$$\pi_3(S^7 \setminus \bigcup_{j \neq i} S_j) \cong \pi_3(\bigvee_{j \neq i} F_j) \cong \bigoplus_{j \neq i} H_3(S^7 \setminus S_j, \mathbf{Z}).$$

Under this identification, we have $[g_i] = \sum_{j \neq i} \lambda_{ij} \cdot e_j$. The $[g_i]$ determine the ambient isotopy class of the given link (3.9), and we deduce

PROPOSITION 4.1. *The linking numbers λ_{ij} , $1 \leq i < j \leq b'$, determine the given link up to ambient isotopy.*

The sphere S_i bounds a 4-dimensional disc D_i^- in D^8 , $i = 1, \dots, b'$, which we equip with the induced orientation. We may, furthermore, assume

that the D_i^- intersect transversely in the interior of D^8 . Then the λ_{ij} coincide with the intersection numbers $D_i^- \cdot D_j^-$, $1 \leq i < j \leq b'$. For an intuitive proof (in dimension 4), see [28], p. 67. Now, every disc D_i^- is completed by the core disc D_i^+ of the i^{th} 4-handle to an embedded 4-sphere Σ_i in T , $i = 1, \dots, b'$, and, since all the core discs are pairwise disjoint, the λ_{ij} coincide with the intersection numbers $\Sigma_i \cdot \Sigma_j$, $1 \leq i < j \leq b'$. Finally, X is obtained by gluing an 8-disc to T along ∂T , and the spheres Σ_i represent the elements of the chosen basis \underline{b} of $H_4(X, \mathbb{Z})$. Identifying the intersection ring with the cohomology ring of X via Poincaré-duality, we see

COROLLARY 4.2. *The linking numbers λ_{ij} coincide with the cup products $(y_i \cup y_j)[X]$, $1 \leq i < j \leq b'$, i.e., the link of the attaching spheres is determined up to ambient isotopy by the basis \underline{b} and the cup products.*

As we have remarked before, the normal bundles of the S_i are trivial, whence there exist embeddings $f_i^0: S^3 \times D^4 \rightarrow S^7$ with $f_i^0|_{S^3 \times \{0\}} = g_i$, $i = 1, \dots, b'$. From the uniqueness of tubular (in differential topology) or regular (in piecewise linear topology) neighbourhoods, every other embedding $f_i: S^3 \times D^4 \rightarrow S^7$ with $f_i|_{S^3 \times \{0\}} = g_i$ is ambient isotopic to one of the form $f_i^{[h_i]} := ((x, y) \mapsto (x, h_i \cdot y))$, $[h_i] \in \pi_3(\text{SO}(4))$, $i = 1, \dots, b'$. Corollary 3.14 implies that we can choose the f_i^0 , $i = 1, \dots, b'$, in such a way that the following holds:

LEMMA 4.3. *Suppose T is obtained by attaching 4-handles along $f_i^{[h_i]}$ with $[h_i] = k_1^i \alpha_3 + k_2^i \beta_3$, $i = 1, \dots, b'$, then*

$$\Sigma_i \cdot \Sigma_i = k_2^i \quad \text{and} \quad p_1(T|_{\Sigma_i}) = \pm(2k_2^i + 4k_1^i).$$

This shows that also the framed link used for constructing T and X is determined by the system of invariants associated to (X, \underline{y}) , proving the injectivity in Part i) of the theorem. Moreover, the assertion about the fibres in Part ii) is clear.

Conversely, given a system Z of invariants in $Z(0, b')$, satisfying relation (2), there exists a based 3-connected manifold (X, \underline{y}) realizing Z . Indeed, by the above identification of the invariants, Z determines a framed link in S^7 and thus the manifold $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$. The boundary of T is a 7-dimensional homotopy sphere ([17], (12.2), p. 119) and, therefore, piecewise linearly homeomorphic to S^7 . Hence, $X = T \cup_{S^7} D^8$ is a piecewise linear manifold with the desired system of invariants, settling Part i). If, in

addition, relation (3) holds, then [18] ensures that X will carry a smooth structure (compare Theorem A.4 of [24]), finishing the proof of Part ii). \square

4.2 THE DETERMINATION OF W_4 IN THE GENERAL CASE

We have a handle decomposition $W_0 \subset W_2 \subset W_4 \subset W_6 \subset X$ of X providing preferred bases \underline{b} of $H_2(X, \mathbf{Z})$ and \underline{c} of $H_4(X, \mathbf{Z})$, respectively. Let \underline{x} and \underline{y} be the dual bases of $H^2(X, \mathbf{Z})$ and $H^4(X, \mathbf{Z})$, respectively. Finally, let \underline{y}^* be the basis of $H^4(X, \mathbf{Z})$ which is dual to \underline{y} via γ_X .

We find $\partial W_2 \cong \#_{i=1}^{b'}(S^2 \times S^5)$, and W_4 is determined by the ambient isotopy class of a framed link of 3-spheres in ∂W_2 with b' components. Let $f_k: S^3 \times D^4 \rightarrow \partial W_2$ be the k^{th} component of that link and $g_k := f_k|_{S^3 \times \{0\}}$, $k = 1, \dots, b'$. In the notation of Section 3.6, we write $[g_k] \in \pi_3(\partial W_2 \setminus \bigcup_{k \neq j} S_j)$ in the form $(l_i^k, i = 1, \dots, b, l_{ij}^k, 1 \leq i < j \leq b; \lambda_{kj}, j \neq k)$, $k = 1, \dots, b'$. To see the significance of the l_i^k and l_{ij}^k , note that, by Remark 3.4, $W_2 \cup H_k^4 \subset X$ is homotopy equivalent to $(\bigvee_{i=1}^b S^2) \cup_{g_k} D^4$. The cohomology ring of that complex has been computed in Proposition 3.11, so that the naturality of the cup product implies the following formulae for the cup products in X :

$$x_i \cup x_j = \sum_{k=1}^{b'} l_{ij}^k \cdot y_k^*, \quad i \neq j,$$

$$x_i \cup x_i = \sum_{k=1}^{b'} l_i^k \cdot y_k^*, \quad i = 1, \dots, b.$$

Therefore, the l_i^k and l_{ij}^k are determined by δ_X and γ_X (used to compute \underline{y}^*), in fact $l_i^k = \gamma_X(\delta(x_i \otimes x_i) \otimes y_k)$ and $l_{ij}^k = \gamma_X(\delta(x_i \otimes x_j) \otimes y_k)$.

To determine the λ_{ij} and the framings, we proceed as follows: Look at the embedding $\#_{i=1}^{b'}(S^2 \times S^5) \hookrightarrow X$. There exist b embedded 2-spheres S_1^2, \dots, S_b^2 which represent the basis \underline{b} and which do not meet the given link. Finally, $\#_{i=1}^{b'}(S^2 \times S^5)$ obviously possesses a regular neighborhood in X which is homeomorphic to $\#_{i=1}^{b'}(S^2 \times S^5) \times D^1$. Thus, we can perform "surgery in pairs" as described in Section 3.1. The result is a 3-connected manifold X^* containing S^7 . It is by construction the manifold obtained from the framed link in S^7 derived from the given one in $\#_{i=1}^{b'}(S^2 \times S^5)$ (cf. Section 4.1). We will be finished, once we are able to compare the invariants of X to those of X^* . To do so, we look at the *trace of the surgery*, i.e., at $Y = (X \times I) \cup H_1^5 \cup \dots \cup H_{b'}^5$, the 5-handles being attached along tubular neighborhoods of the $S_i \times \{1\}$ in $X \times \{1\}$. Then $\partial Y = X \sqcup \bar{X}^*$.

The Mayer-Vietoris sequence provides the isomorphisms

$$H_4(X, \mathbf{Z}) \cong H_4\left(X \setminus \bigsqcup_{i=1}^{b'} (S_i \times D^6), \mathbf{Z}\right) \cong H_4(X^*, \mathbf{Z}).$$

Set $H := H_4\left(X \setminus \bigsqcup_{i=1}^{b'} (S_i \times D^6), \mathbf{Z}\right)$. By Lefschetz duality ([5], (28.18)), there is for each $q \in \mathbf{N}$ a diagram (omitting \mathbf{Z} -coefficients)

$$(4) \quad \begin{array}{ccccccc} H^{q-1}(Y) & \longrightarrow & H^{q-1}(\partial Y) & \longrightarrow & H^q(Y, \partial Y) & \longrightarrow & H^q(Y) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ H_{10-q}(Y, \partial Y) & \longrightarrow & H_{9-q}(\partial Y) & \longrightarrow & H_{9-q}(Y) & \longrightarrow & H_{9-q}(Y, \partial Y) \end{array}$$

where the left square commutes up to the sign $(-1)^{q-1}$ and the other two commute. We first use it in the case $q = 5$. Look at the commutative diagram

$$\begin{array}{ccc} H & \xrightarrow{\cong} & H_4(X^*, \mathbf{Z}) \\ \downarrow \cong & & \downarrow \\ H_4(X, \mathbf{Z}) & \longrightarrow & H_4(Y, \mathbf{Z}), \end{array}$$

in which all arrows are injective, because $H_5(Y, X; \mathbf{Z}) = 0 = H_5(Y, X^*; \mathbf{Z})$ (cf. [17], p. 198). Using the identification $H_4(\partial Y, \mathbf{Z}) = H \oplus H$, we find

$$(5) \quad \text{Im}(H_5(Y, \partial Y; \mathbf{Z})) = \{ (y, -y) \in H \oplus H \}.$$

Similar considerations apply to the case $q = 9$. Taking into account that X^* sits in Y with the reversed orientation, (4) shows that the forms γ_X and γ_{X^*} , both defined with respect to the preferred bases, coincide. In the same manner, the pullbacks of $p_1(Y)$ to $H^4(X, \mathbf{Z})$ and $H^4(X^*, \mathbf{Z})$, respectively, agree. Since X and X^* are the boundary components of Y , these pullbacks are $p_1(X)$ and $p_1(X^*)$, respectively, and we are done. \square

4.3 MANIFOLDS WITH GIVEN INVARIANTS

One might speculate, especially in view of the classification of E-manifolds in dimension 4 and 6, that the invariants δ_X , γ_X , and $p_1(X)$ might suffice to classify E-manifolds with $w_2(X) = 0$ in dimension 8. However,

Lemma 3.6 shows that these invariants determine only W_4 and we still have the choice of an isomorphism in gluing $\#_{i=1}^b(S^2 \times S^5)$ to W_4 , and different gluings may lead to different results. The following example, which was communicated to me by J.-C. Hausmann, illustrates this phenomenon.

EXAMPLE 4.4. One has $\pi_5(\mathrm{SO}(3)) \cong \mathbf{Z}_2$ [32]. Therefore, there are two different S^2 -bundles over S^6 , call them $X := S^6 \times S^2$ and $X' := S^6 \tilde{\times} S^2$. Obviously, X and X' are spin-manifolds with trivial invariants, but one computes $\pi_5(X) \cong \mathbf{Z}_2$ and $\pi_5(X') = \{0\}$.

Fix b, b' , and a system Z of invariants in the image of the map $Z^{\mathrm{PL}(C^\infty)}(b, b')$. As we have seen, Z determines a certain manifold W_4 whose boundary is diffeomorphic to $\#_{i=1}^b(S^2 \times S^5)$ together with a basis \underline{b} for $H_2(\partial W_4, \mathbf{Z})$. Let \underline{b}_0 be the natural basis for $H_2(\#_{i=1}^b(S^2 \times S^5), \mathbf{Z})$, and denote by $\mathrm{Iso}_0^{\mathrm{PL}(C^\infty)}$ the set of piecewise linear (smooth) isomorphisms $f: \#_{i=1}^b(S^2 \times S^5) \rightarrow \partial W_4$ with $f_*(\underline{b}_0) = \underline{b}$. Our results show that every based piecewise linear (smooth) manifold $(X, \underline{x}, \underline{y})$ with system of invariants Z is piecewise linearly (smoothly) isomorphic to a manifold of the form

$$X(f) := \partial W_4 \cup_f \#_{i=1}^b(S^2 \times S^5) \quad \text{for some } f \in \mathrm{Iso}_0^{\mathrm{PL}(C^\infty)}$$

with its given bases for $H^2(X(f), \mathbf{Z})$ and $H^4(X(f), \mathbf{Z})$. Conversely, every manifold of the form $X(f)$ is a piecewise linear (smooth) based E-manifold with invariants Z .

Now, suppose we are given $f, f' \in \mathrm{Iso}_0^{\mathrm{PL}(C^\infty)}$, such that $X(f)$ and $X(f')$ are isomorphic as piecewise linear (smooth) based manifolds. We claim that we can find an isomorphism $\varphi: X(f) \rightarrow X(f')$ with $\varphi(W_4) = W_4$. For this, look at the handle decomposition $W_0 \subset W_2 \subset W_4$. Since W_0 is just an embedded 8-disc in $X(f)$ and $X(f')$, respectively, we can choose φ with $\varphi(W_0) = W_0$. Let $l \subset \partial W_0$ be the framed link for attaching the 2-handles. Then $\varphi(l)$ and l are isotopic. Therefore, we can find a level preserving diffeomorphism $\tilde{\psi}: [-1, 1] \times \partial W_0 \rightarrow [-1, 1] \times \partial W_0$ with $\tilde{\psi}|_{\{\pm 1\} \times \partial W_0} = \mathrm{id}_{\partial W_0}$ and $\tilde{\psi}|_{\{0\} \times \partial W_0}(\varphi(l)) = l$. If we choose a tubular neighborhood ($\cong [-1, 1] \times \partial W_0$) of ∂W_0 in $X(f')$, we can use $\tilde{\psi}$ to define an automorphism $\psi: X(f') \rightarrow X(f')$ with $\psi(\varphi(l)) = l$. Thus, $\psi \circ \varphi$ maps W_2 onto W_2 . A similar argument shows that we can achieve $\varphi(W_4) = W_4$.

Let $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6))$ be the group of piecewise linear (smooth) automorphisms g of $\#_{i=1}^b(S^2 \times D^6)$ with $H^2(g, \mathbf{Z}) = \text{id}$ and similarly define $\text{Aut}_0^{\text{PL}(C^\infty)}(W_4)$. Then we have just established

PROPOSITION 4.5. *The set of isomorphism classes of based piecewise linear (smooth) E-manifolds with invariants \mathbf{Z} is in bijection to the set of equivalence classes in $\text{Iso}_0^{\text{PL}(C^\infty)}$ with respect to the equivalence relation coming from the group action*

$$\begin{aligned} \text{Aut}_0^{\text{PL}(C^\infty)}(W_4) \times \text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6)) \times \text{Iso}_0^{\text{PL}(C^\infty)} &\longrightarrow \text{Iso}_0^{\text{PL}(C^\infty)} \\ (h, g, f) &\longmapsto h|_{\partial W_4} \circ f \circ g|_{\#_{i=1}^b(S^2 \times S^5)}^{-1}. \end{aligned}$$

We shall see in Lemma 5.1 that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ contains the commutator subgroup of $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$.

COROLLARY 4.6. *The set of isomorphism classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the abelian group*

$$\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6)).$$

I have been informed by experts that the structure of the groups $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times S^5))$ and $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6))$ has not yet been determined and that this would be a rather difficult task. Therefore, we choose the viewpoint of framed links in order to finish our considerations. In Theorem 5.2, we will then use this viewpoint to compute the group $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$.

As above, let $(X, \underline{x}, \underline{y})$ be a based piecewise linear (smooth) E-manifold with zero second Stiefel-Whitney class and system of invariants $\mathbf{Z}_{(X, \underline{x}, \underline{y})} = (\delta, \gamma, p)$. We have seen that we can find a framed link l_X of 2-spheres in X which represents the basis \underline{x} and perform surgery along this link in order to get a 3-connected piecewise linear (smooth) based manifold (X^*, \underline{y}) together with a framed link $l_{X^*}^*$ of 5-spheres in it. If $(X', \underline{x}', \underline{y}', l_{X'})$ is another such object where $(X', \underline{x}', \underline{y}')$ is isomorphic to $(X, \underline{x}, \underline{y})$, then clearly we can find an isomorphism $\varphi: (X, \underline{x}, \underline{y}) \longrightarrow (X', \underline{x}', \underline{y}')$ with $\varphi(l_X) = l_{X'}$. Such an isomorphism φ yields, after surgery, an isomorphism $\varphi^*: (X^*, \underline{y}) \longrightarrow (X'^*, \underline{y}')$ with $\varphi^*(l_{X^*}^*) = l_{X'^*}^*$. In particular, the manifold (X^*, \underline{y}) is determined up to piecewise linear (smooth) isomorphism. We call it the *type* of $(X, \underline{x}, \underline{y})$. Note that this notion matters only in the smooth case, by Theorem 2.2.

To summarize, we have

PROPOSITION 4.7. *The set of isomorphism classes of based piecewise linear (smooth) E-manifolds of type (X^*, \underline{y}) is in bijection to the set of equivalence classes of framed links of 5-spheres in X^* where two such links l and l' are considered equivalent, if there is a piecewise linear (smooth) automorphism $\varphi^*: (X^*, \underline{y}) \rightarrow (X^*, \underline{y})$ with $\varphi^*(l) = l'$.*

EXAMPLE 4.8. The group $\mathbb{Z}_2^{\oplus b}$ acts freely on the set of isotopy classes of framed links of b spheres of dimension 5 in X^* by altering the framings of the components. Note that the two possible framings of the trivial bundle on a 5-sphere are distinguished by the fact that one extends over D^6 and the other does not. This property is preserved under piecewise linear homeomorphisms, so that we conclude that $\mathbb{Z}_2^{\oplus b}$ acts also freely on the set of equivalence classes of framed links of b spheres of dimension 5 in X^* .

Note that this completes the classification of Spin-E-manifolds of dimension eight with second Betti number one.

Let us look at manifolds of type S^8 . We claim that two framed links l and l' of 5-spheres are equivalent in the above sense, if and only if they are isotopic. Clearly, after replacing l and l' by isotopic links, we may assume that both of them are contained in the Southern hemisphere and that φ^* is the identity on the Northern hemisphere. Now, choose a representative φ^\dagger for the isotopy class of φ^{*-1} which is the identity on the Southern hemisphere. Then $\varphi^\dagger \circ \varphi^*$ is isotopic to the identity and carries l into l' .

For differentiable manifolds, the operation $X \mapsto X \# \Sigma$, Σ an exotic 8-sphere, establishes a bijection between the set of isomorphism classes of based smooth E-manifolds of type S^8 and the set of isomorphism classes of based smooth E-manifolds of type Σ . We conclude

COROLLARY 4.9. i) *The set of isomorphism classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the group $\text{FL}_b = \text{L}_b \oplus \bigoplus_{i=1}^b \mathbb{Z}_2$.*

ii) *The set of isomorphism classes of based smooth E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the group $\vartheta^8 \oplus \text{FL}_b$.*

Finally, we have to deal with those manifolds for which the cup form δ is trivial. Our investigations in Sections 3.6 and 4.2 show that the framed link of 3-spheres in ∂W_2 can be chosen to be contained in a small disc.

In other words, a manifold X with $\delta_X \equiv 0$ is piecewise linear (smoothly) isomorphic $X^\dagger \# X^*$ where X^* is the type of X and $b_4(X^\dagger) = 0$. As our surgery arguments above reveal, an isomorphism between $X^\dagger \# X^*$ and $X'^\dagger \# X'^*$ can be chosen of the form $\varphi^\dagger \# \varphi^*$ where $\varphi^\dagger: X^\dagger \rightarrow X'^\dagger$ and $\varphi^*: X^* \rightarrow X'^*$ are isomorphisms. Therefore, the set of isomorphism classes of based piecewise linear E-manifolds of type X^* with $b_2 = b$ is in bijection to the set of isomorphism classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$. The same goes for differentiable manifolds of type X^* , if X^* is not diffeomorphic to $X^* \# \Sigma$, Σ an exotic 8-sphere. Otherwise, we have to divide by the action of ϑ^8 . This observation together with Corollary 4.9 settles Theorem 2.4. \square

5. STRUCTURE OF THE GROUP $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$

In this section we prove that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is an abelian group which is, moreover, isomorphic to the group FL_b defined before. This result should be of some independent interest, especially because the group FL_b is quite well understood by Haefliger's work. For $b = 1$, we refer to [20] for more specific information.

We begin with the elementary

LEMMA 5.1. *Let $k \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ be a commutator. Then k extends to an automorphism of $\#_{i=1}^b(S^2 \times D^6)$.*

Proof. For the proof, we depict $\#_{i=1}^b(S^2 \times S^5)$ as follows: Let V_i , $i = 1, \dots, b$, be b copies of $S^2 \times D^6$, and we join V_i and V_{i+1} by a tube $T_i \cong [-1, 1] \times D^7$, $i = 1, \dots, b-1$. The result is a manifold W whose boundary is isomorphic to $\#_{i=1}^b(S^2 \times S^5)$. We make the following normalizations: Write ∂V_i as $(S^2 \times D_+^i) \cup (S^2 \times D_-^i)$, let n_i and s_i be the centers of D_+^i and D_-^i , respectively, and set $S_+^i := S^2 \times n_i$ and $S_-^i := S^2 \times s_i$, $i = 1, \dots, b$. Choose furthermore points $e_i \neq w_i$ in $(S^2 \times D_+^i) \cap (S^2 \times D_-^i)$, $i = 1, \dots, b$, and suppose that $\{-1\} \times D^7 \subset T_i$ is attached to a disc around w_i in ∂V_i and $\{1\} \times D^7 \subset T_i$ to a disc around e_{i+1} in ∂V_{i+1} , $i = 1, \dots, b-1$. Set $T := \bigsqcup_{i=1}^{b-1} T_i$.

Now, let $k = f \circ g \circ f^{-1} \circ g^{-1}$ with $f, g \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$. As $H_2(h, \mathbf{Z})$ is the identity for every element $h \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ and S_\pm^i , $i = 1, \dots, b$, both represent the same basis for $H_2(\partial W, \mathbf{Z})$, h is isotopic to a map h' which satisfies either assumption (A) or (B) below.