

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF  $\sharp_{i=1}^b (S^2 \times S^5)$   
**Autor:** Schmitt, Alexander  
**Kapitel:** 4.1 Proof of Theorem 2.2  
**DOI:** <https://doi.org/10.5169/seals-66077>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## 4. PROOF OF THEOREM 2.2 AND THEOREM 2.4

From now on,  $X$  stands for an eight-dimensional E-manifold with  $w_2(X) = 0$ .

## 4.1 PROOF OF THEOREM 2.2

The classification result for 3-connected E-manifolds of dimension eight is a special case of a result of Wall's [36] and can be easily obtained with the methods described in [17], VII, §12. Let us recall the details, because we will need them later on.

We fix a basis  $\underline{b}$  for  $H_4(X, \mathbf{Z})$  and let  $\underline{y}$  be the dual basis of  $H^4(X, \mathbf{Z})$ . Then there is a handle presentation  $X = D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4 \cup D^8$  with  $\underline{b}$  as the preferred basis. The manifold  $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$  is determined by the ambient isotopy class of a framed link of 3-spheres in  $S^7$ , having  $b'$  components. Let us first look at such a link, forgetting the framing, i.e., suppose we are given embeddings  $g_i: S^3 \rightarrow S^7$  with  $S_i \cap S_j = \emptyset$  for  $i \neq j$ ,  $S_i := g_i(S^3)$ ,  $i = 1, \dots, b'$ . By 3.9, we may assume that the  $g_i$  are differentiable. Observe that the normal bundles of the  $S_i$  are trivial.

We equip  $S_i$  with the orientation induced via  $g_i$  by the standard orientation of  $S^3$  and the normal bundle of  $S_i$  with the orientation which is determined by requiring that the orientation of  $S_i$  followed by that of its normal bundle coincide with the orientation of  $S^7$ . Therefore, a 3-sphere  $F_i$  which bounds the fibre of a tubular neighborhood of  $S_i$  in  $S^7$  inherits an orientation and thus provides a generator  $e_i$  for  $H_3(S^7 \setminus S_i, \mathbf{Z}) \cong \mathbf{Z}$ ,  $i = 1, \dots, b'$ . For  $i \neq j$ , the image of the fundamental class  $[S_i]$  in  $H_3(S^7 \setminus S_j, \mathbf{Z})$  is of the form  $\lambda_{ij} \cdot e_j$ . The integer  $\lambda_{ij}$  is called *the linking number of  $S_i$  and  $S_j$* .

For  $i = 1, \dots, b'$ , the manifold  $S^7 \setminus \bigcup_{j \neq i} S_j$  is up to dimension 5 homotopy equivalent to  $\bigvee_{j \neq i} F_j$ , and

$$\pi_3(S^7 \setminus \bigcup_{j \neq i} S_j) \cong \pi_3(\bigvee_{j \neq i} F_j) \cong \bigoplus_{j \neq i} H_3(S^7 \setminus S_j, \mathbf{Z}).$$

Under this identification, we have  $[g_i] = \sum_{j \neq i} \lambda_{ij} \cdot e_j$ . The  $[g_i]$  determine the ambient isotopy class of the given link (3.9), and we deduce

**PROPOSITION 4.1.** *The linking numbers  $\lambda_{ij}$ ,  $1 \leq i < j \leq b'$ , determine the given link up to ambient isotopy.*

The sphere  $S_i$  bounds a 4-dimensional disc  $D_i^-$  in  $D^8$ ,  $i = 1, \dots, b'$ , which we equip with the induced orientation. We may, furthermore, assume

that the  $D_i^-$  intersect transversely in the interior of  $D^8$ . Then the  $\lambda_{ij}$  coincide with the intersection numbers  $D_i^- \cdot D_j^-$ ,  $1 \leq i < j \leq b'$ . For an intuitive proof (in dimension 4), see [28], p. 67. Now, every disc  $D_i^-$  is completed by the core disc  $D_i^+$  of the  $i^{\text{th}}$  4-handle to an embedded 4-sphere  $\Sigma_i$  in  $T$ ,  $i = 1, \dots, b'$ , and, since all the core discs are pairwise disjoint, the  $\lambda_{ij}$  coincide with the intersection numbers  $\Sigma_i \cdot \Sigma_j$ ,  $1 \leq i < j \leq b'$ . Finally,  $X$  is obtained by gluing an 8-disc to  $T$  along  $\partial T$ , and the spheres  $\Sigma_i$  represent the elements of the chosen basis  $\underline{b}$  of  $H_4(X, \mathbb{Z})$ . Identifying the intersection ring with the cohomology ring of  $X$  via Poincaré-duality, we see

**COROLLARY 4.2.** *The linking numbers  $\lambda_{ij}$  coincide with the cup products  $(y_i \cup y_j)[X]$ ,  $1 \leq i < j \leq b'$ , i.e., the link of the attaching spheres is determined up to ambient isotopy by the basis  $\underline{b}$  and the cup products.*

As we have remarked before, the normal bundles of the  $S_i$  are trivial, whence there exist embeddings  $f_i^0: S^3 \times D^4 \rightarrow S^7$  with  $f_i^0|_{S^3 \times \{0\}} = g_i$ ,  $i = 1, \dots, b'$ . From the uniqueness of tubular (in differential topology) or regular (in piecewise linear topology) neighbourhoods, every other embedding  $f_i: S^3 \times D^4 \rightarrow S^7$  with  $f_i|_{S^3 \times \{0\}} = g_i$  is ambient isotopic to one of the form  $f_i^{[h_i]} := ((x, y) \mapsto (x, h_i \cdot y))$ ,  $[h_i] \in \pi_3(\text{SO}(4))$ ,  $i = 1, \dots, b'$ . Corollary 3.14 implies that we can choose the  $f_i^0$ ,  $i = 1, \dots, b'$ , in such a way that the following holds:

**LEMMA 4.3.** *Suppose  $T$  is obtained by attaching 4-handles along  $f_i^{[h_i]}$  with  $[h_i] = k_1^i \alpha_3 + k_2^i \beta_3$ ,  $i = 1, \dots, b'$ , then*

$$\Sigma_i \cdot \Sigma_i = k_2^i \quad \text{and} \quad p_1(T|_{\Sigma_i}) = \pm(2k_2^i + 4k_1^i).$$

This shows that also the framed link used for constructing  $T$  and  $X$  is determined by the system of invariants associated to  $(X, \underline{y})$ , proving the injectivity in Part i) of the theorem. Moreover, the assertion about the fibres in Part ii) is clear.

Conversely, given a system  $Z$  of invariants in  $Z(0, b')$ , satisfying relation (2), there exists a based 3-connected manifold  $(X, \underline{y})$  realizing  $Z$ . Indeed, by the above identification of the invariants,  $Z$  determines a framed link in  $S^7$  and thus the manifold  $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$ . The boundary of  $T$  is a 7-dimensional homotopy sphere ([17], (12.2), p. 119) and, therefore, piecewise linearly homeomorphic to  $S^7$ . Hence,  $X = T \cup_{S^7} D^8$  is a piecewise linear manifold with the desired system of invariants, settling Part i). If, in

addition, relation (3) holds, then [18] ensures that  $X$  will carry a smooth structure (compare Theorem A.4 of [24]), finishing the proof of Part ii).  $\square$

#### 4.2 THE DETERMINATION OF $W_4$ IN THE GENERAL CASE

We have a handle decomposition  $W_0 \subset W_2 \subset W_4 \subset W_6 \subset X$  of  $X$  providing preferred bases  $\underline{b}$  of  $H_2(X, \mathbf{Z})$  and  $\underline{c}$  of  $H_4(X, \mathbf{Z})$ , respectively. Let  $\underline{x}$  and  $\underline{y}$  be the dual bases of  $H^2(X, \mathbf{Z})$  and  $H^4(X, \mathbf{Z})$ , respectively. Finally, let  $\underline{y}^*$  be the basis of  $H^4(X, \mathbf{Z})$  which is dual to  $\underline{y}$  via  $\gamma_X$ .

We find  $\partial W_2 \cong \#_{i=1}^{b'}(S^2 \times S^5)$ , and  $W_4$  is determined by the ambient isotopy class of a framed link of 3-spheres in  $\partial W_2$  with  $b'$  components. Let  $f_k: S^3 \times D^4 \rightarrow \partial W_2$  be the  $k^{\text{th}}$  component of that link and  $g_k := f_k|_{S^3 \times \{0\}}$ ,  $k = 1, \dots, b'$ . In the notation of Section 3.6, we write  $[g_k] \in \pi_3(\partial W_2 \setminus \bigcup_{k \neq j} S_j)$  in the form  $(l_i^k, i = 1, \dots, b, l_{ij}^k, 1 \leq i < j \leq b; \lambda_{kj}, j \neq k)$ ,  $k = 1, \dots, b'$ . To see the significance of the  $l_i^k$  and  $l_{ij}^k$ , note that, by Remark 3.4,  $W_2 \cup H_k^4 \subset X$  is homotopy equivalent to  $(\bigvee_{i=1}^b S^2) \cup_{g_k} D^4$ . The cohomology ring of that complex has been computed in Proposition 3.11, so that the naturality of the cup product implies the following formulae for the cup products in  $X$ :

$$x_i \cup x_j = \sum_{k=1}^{b'} l_{ij}^k \cdot y_k^*, \quad i \neq j,$$

$$x_i \cup x_i = \sum_{k=1}^{b'} l_i^k \cdot y_k^*, \quad i = 1, \dots, b.$$

Therefore, the  $l_i^k$  and  $l_{ij}^k$  are determined by  $\delta_X$  and  $\gamma_X$  (used to compute  $\underline{y}^*$ ), in fact  $l_i^k = \gamma_X(\delta(x_i \otimes x_i) \otimes y_k)$  and  $l_{ij}^k = \gamma_X(\delta(x_i \otimes x_j) \otimes y_k)$ .

To determine the  $\lambda_{ij}$  and the framings, we proceed as follows: Look at the embedding  $\#_{i=1}^{b'}(S^2 \times S^5) \hookrightarrow X$ . There exist  $b$  embedded 2-spheres  $S_1^2, \dots, S_b^2$  which represent the basis  $\underline{b}$  and which do not meet the given link. Finally,  $\#_{i=1}^{b'}(S^2 \times S^5)$  obviously possesses a regular neighborhood in  $X$  which is homeomorphic to  $\#_{i=1}^{b'}(S^2 \times S^5) \times D^1$ . Thus, we can perform "surgery in pairs" as described in Section 3.1. The result is a 3-connected manifold  $X^*$  containing  $S^7$ . It is by construction the manifold obtained from the framed link in  $S^7$  derived from the given one in  $\#_{i=1}^{b'}(S^2 \times S^5)$  (cf. Section 4.1). We will be finished, once we are able to compare the invariants of  $X$  to those of  $X^*$ . To do so, we look at the *trace of the surgery*, i.e., at  $Y = (X \times I) \cup H_1^5 \cup \dots \cup H_{b'}^5$ , the 5-handles being attached along tubular neighborhoods of the  $S_i \times \{1\}$  in  $X \times \{1\}$ . Then  $\partial Y = X \sqcup \bar{X}^*$ .