

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TORSION NUMBERS OF AUGMENTED GROUPS WITH APPLICATIONS TO KNOTS AND LINKS

Autor: Silver, Daniel S. / Williams, Susan G.

Kurzfassung

DOI: <https://doi.org/10.5169/seals-66079>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 20.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

TORSION NUMBERS OF AUGMENTED GROUPS WITH APPLICATIONS TO KNOTS AND LINKS

by Daniel S. SILVER and Susan G. WILLIAMS *)

Dedicated to the memory of Arnold E. Ross

ABSTRACT. Torsion and Betti numbers for knots are special cases of more general invariants b_r and β_r , respectively, associated to a finitely generated group G and epimorphism $\chi: G \rightarrow \mathbf{Z}$. The sequence of Betti numbers is always periodic; under mild hypotheses about (G, χ) , the sequence b_r satisfies a linear homogeneous recurrence relation with constant coefficients. Generally, b_r exhibits exponential growth rate. However, again under mild hypotheses, the p -part of b_r has trivial growth for any prime p . Applications to branched cover homology for knots and links are presented.

1. INTRODUCTION

A *knot* is a simple closed curve in the 3-sphere S^3 . Knots are *equivalent* if there is an orientation-preserving homeomorphism of S^3 that carries one into the other. Equivalent knots are regarded as the same. An *invariant* is a well-defined quantity that depends only on a knot equivalence class. Two knots for which some invariant differs are necessarily distinct.

Associated to any knot k and natural number r there is a compact, oriented 3-manifold M_r , the r -fold cyclic cover of S^3 branched over k . A precise definition can be found in [Li97] or [Ro76], for example. Topological invariants of M_r are invariants of k . Two such invariants, the first Betti number β_r and the order b_r of the torsion subgroup of $H_1(M_r; \mathbf{Z})$, were first considered by J. Alexander and G. Briggs [Al28], [AB27] and by O. Zariski [Za32]. The continuing interest in these invariants is witnessed by numerous papers (e.g., [Go72], [Me80], [We80], [Ri90] and [GS91]). We call b_r the r^{th} *torsion*

*) First author partially supported by ÉGIDE at CMI, Université de Provence. Second author partially supported by CNRS at Institut de Mathématiques de Luminy. Both authors partially supported by NSF grant DMS-0071004.